

Small RNA biology is systems biology

Daniel Jost, Andrzej Nowojewski & Erel Levine*

Department of Physics, FAS Center for Systems Biology, Harvard University, Cambridge MA 02138, USA

During the last decade small regulatory RNA (srRNA) emerged as central players in the regulation of gene expression in all kingdoms of life. Multiple pathways for srRNA biogenesis and diverse mechanisms of gene regulation may indicate that srRNA regulation evolved independently multiple times. However, small RNA pathways share numerous properties, including the ability of a single srRNA to regulate multiple targets. Some of the mechanisms of gene regulation by srRNAs have significant effect on the abundance of free srRNAs that are ready to interact with new targets. This results in indirect interactions among seemingly unrelated genes, as well as in a crosstalk between different srRNA pathways. Here we briefly review and compare the major srRNA pathways, and argue that the impact of srRNA is always at the system level. We demonstrate how a simple mathematical model can ease the discussion of governing principles. To demonstrate these points we review a few examples from bacteria and animals. (BMB reports 2011; 44(1): 11-21)

INTRODUCTION

The role of RNA molecules in the regulation of gene expression was first suggested by Jacob and Monod 50 years ago (1). Still, for over 4 decades evidence for RNA-based regulation was limited to the control of mobile elements (plasmids and transposons) and isolated examples of endogenous bacterial small RNAs in *Escherichia coli*. But at the turn of the millennium evidence for the real impact of small RNA were accumulating: the first animal microRNA, the heterochronic gene *lin-4* of the nematode *Caenorhabditis elegans*, was discovered (2, 3); microarray experiments, comparative genomics and bioinformatics (4-6) revealed dozens of small RNAs in *E. coli*; and the ability of double-stranded RNA to completely silence endogenous gene expression indicated the existence of the RNAi pathway (7). But full appreciation for the role of RNA in con-

trolling gene regulation came only when another *C. elegans* microRNA, *let-7*, was shown to have homologs in species across the animal kingdom, including mouse and human (8, 9). Since then, the accumulation of genome sequences that allow bioinformatic identification of small RNA, and in particular the emergence of Next-Generation Sequencing methods enabling deep sequencing of the transcriptome, led to identification of countless microRNAs and multiple small RNA pathways (For recent reviews see 10, 11).

Small RNAs in Animals

Multiple srRNA pathways exist in animals, including the well studied microRNAs (miRNAs), endogenous small-interfering RNAs (siRNAs) in the RNAi pathway, and Piwi-interacting RNAs (piRNAs) (Fig. 1). MiRNAs have been shown to play significant roles in all fields of biology, including development (reviewed in 12, 13), stem cell differentiation (14), germline maintenance (15, 16), cancer biology (17) and stress response (18). piRNAs have been implicated in maintenance of genomic integrity and suppression of transposon activity (19). The role endogenous siRNAs is even more diverse and less




Fig. 1. Outline of three srRNA pathways in animals (miRNA and siRNA) and bacteria (Hfq-dependent sRNAs).

*Corresponding author. Tel: +1-617-495-3735; E-mail: elvine@fas.harvard.edu
DOI 10.5483/BMBRep.2011.44.1.11

Received 12 January 2011

Keywords: Mathematical modeling, Post-transcriptional regulation, Small regulatory RNAs, Systems biology

understood.

Small RNA pathways differ in the biogenesis of the small RNAs. MiRNAs are derived from distinctive hairpin precursors that are genomically encoded, transcribed by RNA polymerase II and undergo several processing steps (20, 21). Endogenous siRNAs and piRNAs in animals come from multiple sources, including transposons, long RNA transcripts, and mRNAs (22-31); siRNAs are processed from double stranded RNAs, while piRNAs are not. Different sets of proteins are involved in biogenesis of srRNAs in different pathways, although some of these enzymes can work in multiple pathways (see below).

The mechanism of gene silencing is also divergent: siRNAs lead to direct cleavage of the bound mRNA (32, 33), while the mechanism of miRNA silencing is less direct, and may involve both translation inhibition and mRNA destabilization (reviewed in 11). siRNAs require perfect base-pairing with their target, whereas miRNA hybridization to their target is more complex, and involves a perfectly matched "seed" 6-8 nt long and some mismatches outside of it. In some animals (and in plants), siRNA or piRNA binding to its target can trigger the synthesis of secondary siRNAs (34-37), thus amplifying the silencing signal. In nematodes (and in plants) this allows for systemic silencing, where the amplified silencing signal can travel across cells and tissues (38).

At the same time, much is common to the diverse RNA pathways. Importantly, specificity is attributed to Watson-Crick basepairing between the srRNA and its targets. Dicer is involved in srRNA processing in multiple pathways. All known srRNAs bind to a member of the Argonaute protein family, which confers stability to the small RNA molecule, facilitates its binding with its mRNA target, and is involved in recruiting additional factors required for silencing.

Small RNAs in Bacteria

Bacterial small regulatory RNAs (recently reviewed in 39, 40) are longer than their eukaryotic counterparts, and are typically between 60-150 nt long. This can be attributed to the rarity of RNA processing in bacteria, as animal miRNAs are also synthesized as longer molecules that are processed to their final small size post-transcriptionally. Bacterial srRNAs typically bind the 5'UTR of their target in the vicinity of the ribosomal binding site (RBS), resulting in modulation of translation efficiency and stability of the mRNA molecules. While RNA regulation in eukaryotes typically results in gene silencing, bacterial small RNAs can both repress or activate their targets.

The physiological role of bacterial small RNAs is diverse. Notably, bacterial small RNAs are abundant in stress response pathways (41), and have been implicated in the regulation of virulence genes (42-47). While the molecular interaction between a bacterial small RNA and its target may be more complex than that of the short srRNA in eukaryotes, recognition is still typically limited to a short (7-10 nt) sequence of almost perfect complementarity (40).

Small RNA Recycling Determines its Functional Properties

Post-transcriptional silencing of a gene requires targeting the multiple mRNA molecules that reside in the cytoplasm. These molecules need to be efficiently found by the srRNA, bound and silenced. In the RNAi pathway, a siRNA - bound to an Argonaute protein in an RNA-induced silencing complex (RISC) (48) - can be recycled and undergo multiple rounds of silencing (49-51). In some cases (as mentioned above) the interaction of the siRNA with its target can even lead to the synthesis of secondary siRNAs. With these two properties, a relatively small dsRNA trigger can result in efficient - and sometimes systemic - gene silencing.

Gene silencing by miRNAs, however, is believed to be more complex than simple direct cleavage of the target, leading to some titration of the miRNA to its target. MiRNAs therefore act in a more stoichiometric way, and their levels should be better matched with that of their targets (52). Indeed, in cases where a miRNA is known to strongly suppress its target, the miRNA is highly expressed.

An abundant class of bacterial small RNAs marks the other extreme. These srRNAs require the bacterial RNA chaperone Hfq for efficient gene silencing (40). Hfq binds to these srRNAs and stabilizes them, thus forming the bacterial equivalent of RISC. Small RNAs of this class may be directly affected by the interaction with their target. In such cases, a small RNA directs cleavage of the srRNA:mRNA complex, thus reducing the stability of the mRNA target while promoting its own degradation (53, 54).

Recent quantitative studies of small RNA regulation focused on the implications of incomplete (or absent) srRNA recycling on the properties of target repression (55-59). Using simple kinetic models (Box 1) it has been demonstrated that srRNAs can impose a threshold for the activation of gene expression: a significant gene silencing is achieved when the rate of mature mRNA production is smaller than the threshold, and is effectively relieved above it (Fig. 2). This mechanism for silencing has been shown to exhibit reduced level of fluctuations, as compared with a more catalytic reaction (such as transcriptional repression). Dynamically, stoichiometric srRNA-target interactions result in a delayed response to abrupt changes in the activity of either gene, allowing filtering of environmental noise. These models suggest that the efficiency of srRNA regulation (*i.e.* the sharpness of the transition, noise reduction etc) is controlled by a single effective parameter which lumps together the biochemical properties of the srRNA and its target (Box 1 and Fig. 2).

Coordinated Response in Bacteria (1): srRNA Regulation of Master Regulators

Two of the better characterized srRNAs in bacteria are regulating multiple mRNA species including global transcriptional

Box 1. Minimal Model for Small RNA Regulation

The motivation behind composing a generic mathematical model for small RNA based regulation comes from two “big” questions. First, on the face of it small RNA regulation and transcriptional regulation seem redundant. Nevertheless, the modes of regulation coexist almost in all organisms, suggesting that the two complement each other in some non-trivial way. It was therefore suggested that small RNA regulation might exhibit quantitative features that make it different from protein-based transcriptional regulation. Second, given that small RNAs have been missed over decades of molecular biology, it has been suggested that their role in regulation is minor, idea that was supported by the fact that deletion of almost all miRNAs in *C. elegans*, one by one, has no effect on development or viability. It was therefore suggested that small RNAs have a more quantitative role in tuning gene expression.

In order to answer such high-level questions, one needs to construct a model that is stripped down to the essential ingredients, without losing the essence of the modeled system. When done right, details can either be lumped together into effective parameters, or can be accounted for by functions whose properties - albeit not detailed form - can be postulated.

Generic models for small RNA regulation typically focus on a small RNA and one (or a small number of) targets, and account for synthesis of the RNA molecules (with rates denoted below by α); bare degradation (denoted by β) or active degradation (by χ) of the molecules; and the two-species interaction (denoted by k). Recycling of the small RNA can be parameterized via a probability parameter q . In its simplest form, the model is represented by a set of mass-action equations.

$$\frac{dm_i}{dt} = \alpha_i - \beta_i m_i - (k_i^+ m_i s - k_i^- c_i)$$

$$\frac{ds_i}{dt} = \alpha_s - \beta_s s - \sum_i [(k_i^+ m_i s - (k_i^- - q\chi_i) c_i)]$$

$$\frac{dc_i}{dt} = (k_i^+ m_i s - k_i^- c_i) - \chi_i c_i$$

Here m_i is the concentration of mRNAs of species i , s is the concentration of the srRNA, and c_i the concentration of srRNA:mRNA complex. To complete the model one needs to specify the process of translation. For example, in cases where translation is inhibited directly by binding of RISC, the concentration of protein i will be proportional to that of its free mRNA, m_i . However if RISC binding only facilitates degradation of the molecules (e.g. in the siRNA pathway) then protein concentration is proportional to $(m_i + c_i)$.

More generally, other linear combinations of the two are possible.

Even in this simplified scheme, these are highly coupled non-linear differential equations, and one cannot solve them exactly. However, if one focuses on steady state properties, all time derivatives (the left-hand side of these equations) are set to zero, and one remains with a simpler set of algebraic equations. These are readily solved for the case of one or two targets. In this case the kinetics of the interaction between the srRNA and a target is given by the binding rate $k = \chi k^+ / (k^- + \chi) \approx k^+$. The last approximation holds in cases where the complex is efficiently degraded (as in the bacterial case and in the RNAi pathway).

A key feature of this model is a non-linear relation between the transcription rate of a gene and its steady-state mRNA concentration (Fig. 2). In this “threshold-linear” response, the gene is efficiently silenced when its transcription rate is below a prescribed threshold. Above the threshold, the mRNA level increases linearly with the transcription rate. The behavior near the threshold is determined by a combination of all other model parameters, that measures the efficiency of srRNA binding and regulation. Efficient regulation (e.g. large k and/or small β 's) makes the transition sharp and well defined; less efficient regulation makes it broader and leaky.

The “threshold-linear” picture already provides one way to see the cross-talk between different targets of an srRNA. From the perspective of one target (say, $i=1$), the threshold in α_i is given by $\alpha_s - \alpha_2$. Thus, at one level of α_i the first gene may be either in the silenced or active state, depending on the rate of transcription of the second target. Alternatively, one can either use numerical methods, or group all “other” targets into a single effective species.

Increasing the complexity of the mathematical description can reveal more quantitative features. To study the dynamics of regulatory response, one retains the temporal derivatives. While limited analytical progress can be made, integrating these equations numerically is straightforward, and one can identify scenarios where the temporal response may be faster or slower than the expected response in transcriptional regulatory circuits. The model can be developed further to account of the discreteness in molecules number and the burstiness in transcription and translation processes, showing that small RNA acts efficiently to suppress intrinsic noise in the silenced state.

Without major modifications, this model can integrate more details of the pathway and account for more players, assuming that these act to modify the kinetic parameters. This approach is taken in the text to account for enzymes of the srRNA pathways (such as the bacterial RNA chaperone Hfq, and the eukaryotic Dicer and Argonaute proteins).

regulators. One of the two is the *E. coli* srRNA DsrA. DsrA is repressing the expression of H-NS proteins by binding at the ribosome binding site (RBS) of the Hns mRNA, resulting in repression of translation and destabilization of the mRNA molecule (60). H-NS is a nucleoid-associated DNA-binding protein abundant in enteric bacteria. H-NS binds to high-affinity sites and spreads along adjacent AT-rich DNA to silence transcription, allowing it to repress the expression of foreign DNA with higher AT-content than the resident genome (61). H-NS has long been regarded as a global modulator of gene expression in response to pH, temperature, osmolarity and growth phase (62, 63).

DsrA is also an activator of a global transcription regulator, the major enterobacterial stress sigma factor, σ^S , encoded by *rpoS* (64). In the absence of DsrA, the long 5'UTR of the RpoS mRNA forms a secondary structure that obscures its RBS, thus

attenuating translation. In the presence of the small RNA, an alternative secondary structure is stabilized to promote ribosome binding and increasing translation. We note in passing that RpoS is also regulated by two other small RNAs, RprA and OxyS (65-68).

The *dsrA* promoter is active only at low temperatures ($<30^\circ\text{C}$) (69, 70). Consequently, this small RNA conveys global transcriptional response to low temperature stress via two major transcription regulators. The quantitative model described above suggests that this regulation takes the form of a threshold response. Above the threshold temperature, transcription of *dsrA* is smaller than that of its targets, and its presence is ignored; as temperature falls below the threshold the targets become significantly affected.

The Qrr sRNAs of the *Vibrio* species form the second example. These small RNAs have been identified as inter-




Fig. 2. The threshold-linear model for small RNA regulation. Target is strongly suppressed only when synthesis rate of the small RNA exceeds that of the target. Efficient silencing (e.g. strong binding) results in sharpening of the threshold. Here binding is 10 times stronger in the blue curve, dashed line for the unregulated gene.

mediates in the quorum sensing signal transduction pathway (71-73). The Quorum sensing mechanism integrates multiple chemical signals to sense the density of like cells in the environment (74). Based on these signals, a master regulator is activated and goes on to regulate multiple response pathways. Interestingly, the (redundant multi-copied) Qrr sRNAs are positioned downstream of signal integration (by transcriptional response regulators) and up-stream of the master regulator.

The function of this pathway can be described in terms of a computational device called a classifier. A classifier takes multiple inputs, and needs to classify them into one of several possible classes. In the case of the quorum sensing pathways, a linear “weighted” sum of the input signals are passed through the thresholding filter provided by the sRNAs to performs a classification between “high” and “low” cell density (71). This classification scheme is known as a linear classifier, and is one of the basic ingredients in the theory of machine learning.

These two examples show how small RNAs can play a role in global regulation by targeting just a few strategic targets. The role of the sRNA in this case is to link between the input signal (or signals) and the response by setting a threshold for the response. It is easy to see how global regulation would benefit from the quantitative features of sRNA regulation, in particular the tight repression and noise suppression in the silenced state.

The Many Targets of Small RNAs

The sequence that is responsible for specificity of sRNA-target interaction is limited to around 6-8 nt. It is therefore no surprise that naïve attempts at predicting targets of a small RNA yield astounding number of potential targets per sRNA. As target prediction algorithms improve to incorporate target struc-

ture, thermodynamic predictions, and - most importantly - evolutionary conservation (75), the idea that most miRNA have multiple potential targets becomes substantiated. Indeed, some miRNAs are predicted to have even hundreds of targets (76, 77).

MiRNAs certainly affect the evolution of mRNA 3'UTR (78, 79), such that binding targets for a particular miRNA would be specifically absent from genes that are co-expressed with the miRNA. Thus, in many cases putative miRNA binding sites have no real biological implication, as they are present in mRNAs that are rarely co-expressed with the targeting miRNA.

Still, the association of multiple targets with one miRNA raises the possibility that a small RNA rewires the genetic network at the system level. Incomplete recycling of the small RNA results in indirect interactions among targets of a small RNA, as the expression of each target affects the shared srRNA. From a quantitative standpoint, these interactions can be accounted for without significantly complicating the mathematical model (Box 1). This model suggests that this crosstalk may in fact be highly efficient and sensitive (55, 56).

In the reminder of this review we discuss a few examples for the system-level behavior of small RNA, starting from a small pathway and working our way up towards the cell-level system.

Coordinated Response (2): Recycling-dependent Cross Talk between Targets of a Small RNA

Targets of a common srRNA can interact indirectly by modulating srRNA abundance. This is exemplified clearly by a conserved bacterial small RNA, ChiX, which acts in the chitin metabolism network (54, 80). Chitin is an abundant organic polymer, whose main degradation product is chitobiose (N-acetylglucosamine dimer).

Two conserved targets of ChiX are *chb* and *chiP*. The *chb* operon in *E. coli* and *Salmonella* encodes genes for the transport and degradation of chitobiose. This operon also encodes its own regulator, *chbR*, which represses transcription in the absence of chitobiose and activates it in its presence. The chitoporin ChiP was recently found to be absolutely required for *Salmonella* to grow on chitotriose, but is not essential for uptake of chitobiose. While the *chb* operon is transcriptionally active only in the presence of the inducer, *chiP* is transcribed constitutively.

The small RNA ChiX is also constitutively expressed. In the absence of the inducer, it strongly suppresses the accumulation of ChiP mRNA, but is not co-degraded. Conversely, in the presence of chitobiose the Chb mRNA binds the sRNA and degrades it (presumably with little or no effect on the expression of genes in the operon). Thus, transcription of *chb* allows ChiP mRNA (and proteins) to accumulate.

Interestingly, *ChbR*, that up-regulates the transcription of *chb* (81), also activates (moderately) the transcription of *chiP*. What is the need, then, for the indirect activation through ChiX? A possible answer comes from the quantitative model

(Box 1). Analysis of this model (82) suggests a hierarchy between the two targets. In this scenario, sudden increase in chitobiose is met with immediate and gradual response by *chb*. Conversely, activation of *chiP* is delayed and rapid, and occurs only if *chb* activity crosses a prescribed threshold. Possibly, this hierarchy reflects the relative abundance of the different degradation products of chitin.

This simple model allows querying other aspects of the relationship between the targets (82). For example, one may expect that efficient depletion of ChiX would require a preference for binding to *chb* rather than to ChiP. Model results, however, indicates that once the binding rates to both targets are at the same range, no significant improvement is achieved from preferential attachment to *chb*. This is due to the cooperative non-linearity of this mode of action.

Coordinated Response (3): The Oxidative Stress Hypothesis

The fact that a small RNA is not only affecting its targets but is also affected by them offers an attractive interpretation for the role of a small RNA in coordinating stress response. For clarity we focus on RyhB, a small RNA that is involved in iron metabolism in enteric bacteria (83-87). The ferric uptake repressor Fur, the key regulator of genes involved in iron uptake and metabolism, transcriptionally controls RyhB. Transcription of the *RyhB* gene is activated under iron limitation. Multiple genes were found to be under the control of RyhB, including genes encoding for non-essential iron-using proteins, ferritins and importantly the superoxide dismutase (SOD) SodB (83, 85). RyhB is also a positive regulator of ShiA, a permease of shikimate, a compound believed to function in the biosynthesis of siderophores that acquire extracellular iron (88). Fur itself is also repressed by RyhB (89).

Given that most targets of RyhB utilize iron, it has been suggested that RyhB assists in controlling the flux of iron in the cell, along with other regulators (85). Quantitative analysis of iron metabolism in *E. coli* suggested a fairly robust control system (90), however it failed to explain why a small RNA is required in this position.

We propose here an alternative, system-level view of the role of RyhB. We note that one target, SodB, stands out as a major target of RyhB. First, the SodB mRNA has the longest conserved seed sequence complementarity among all putative targets of RyhB. Second, the mRNA level of SodB has been shown to be most strongly affected by a *RyhB* deletion (83, 85). Finally, some enteric bacteria carry multiple copies of RyhB, some of which conserve only the base-pairing with SodB (A Nowojewski, unpublished data). SodB, which encodes for the Fe-SOD, also stands out for its function. While it indeed obeys the definition of an iron-carrying gene that is not essential under iron starvation, SodB is the only RyhB target that functions as a stress response gene.

Superoxide is only mildly reactive physiologically. However, iron interacts with this species to generate a highly reactive

and extremely damaging hydroxyl radical. *In vivo* superoxide concentrations are considered too low to cause iron reduction, but can be sufficiently high to damage the exposed (4Fe-4S) clusters of iron-sulfur proteins, some of which are targets of RyhB (91). Iron-mediated sensitivity to superoxide stress is also overcome by increasing the levels of iron chelators and the storage capacity through ferritins.

Taken together, we suggest that the targets of RyhB can be classified in two classes: those targets that are regulated by RyhB, and those that modulate RyhB. SodB is our major candidate for the first class. The levels of all other genes, together with transcriptional regulation by Fur, make the effective accumulation rate of RyhB a faithful gauge for the iron-related severity of superoxide-stress, that goes beyond the instantaneous iron concentration (82). Targets of the second class also play a role in further reducing fluctuations in the main target (82). The ability of utilizing RyhB to estimate future levels of free iron may also be useful during pathogenesis, as the bacterial colony engages in a battle for iron with the host.

Might similar target classification be relevant to animal microRNA? One evidence in this direction comes from the two founding fathers of the miRNAs family, the miRNAs *let-7* and *lin-4* of *C. elegans*. Each of these miRNA is predicted to have tens of targets. However, a compensating mutation in only one gene (*lin-41* and *lin-14*, respectively) is enough to suppress a mutation in the miRNAs sequence (92, 93). It is possible, then, that these are the "class-1" targets of these miRNAs, while other targets serve to regulate (or buffer) the miRNA.

Competition for the Enzymes of the srRNA Pathways

Many of the enzymes that make the srRNA pathways are shared among different pathways. This feature, however, is not conserved. Mammals and *C. elegans*, for example, have only one Dicer that is shared between the miRNA and siRNA pathways (94, 95), while in *D. melanogaster* two enzymes, Dicer-1 and Dicer-2, are used exclusively in the miRNA and siRNA pathways, respectively (96). *C. elegans*, on the other hand, has at least 27 Argonaute proteins, each with a unique function, as opposed to humans where one of 4 Argonaute proteins, Ago2, functions in the miRNA and multiple siRNA pathways (97). Therefore, under conditions where one or more of these - and other enzymes in the srRNA pathways - become rate limiting, shared enzymes may introduce cross-talk among different pathways.

Due to the high efficiency of RNAi regulation, synthetic siRNAs or miRNAs have been widely developed for genomic studies or potential therapeutic treatments. To ensure complete silencing of the target gene one is typically motivated to use high levels of siRNA. However, many side effects have been noticed when introducing highly concentrated synthetic sRNAs necessary to silence the desired target gene (98). In particular, the down-regulation of many off-targets of the introduced sRNA is observed (98, 99) and more surprisingly, the

global up-regulation of transcripts which do not share complementary site with the sRNA (100). These changes in the regulatory pathways could potentially lead to toxic or lethal modifications (101). While the down-regulation effect is straightforward and could simply be explained by the presence of extra-regulatory molecules in the cell, it has been advanced that the up-regulation effect could be due to the saturation of the endogenous RNAi machinery by a high number of exogenous sRNAs that compete for the enzymes (102).

In studying the function of endogenous miRNAs (or siRNAs) it is a common practice to study both loss-of-function and gain-of-function mutants. One way of achieving the latter is creating a transgenic line that over expressed the endogenous miRNA. Once again, strong over expression (e.g. through the use of a high-copy plasmid) may outcompete other sRNAs, expressed at endogenous levels, for RNAi enzymes.

To account for competition over enzymes, the simplified model introduced above needs to be augmented with a quantitative relation between the kinetic parameters of the model (e.g. synthesis and binding rates) and the concentrations of the enzymes. For example, if the rate limiting enzyme are components of the RISC (e.g. Ago2, as believed to be the case in RNAi experiments (102,103)) the siRNA:mRNA complex formation rate k would need to be specified as $k = k_0 [RISC] / (K + [srRNA]_{Total})$. Here $[srRNA]_{Total}$ accounts for the total concentration of all free srRNAs that compete for the enzyme, and may include srRNAs from different pathways. It is therefore organism dependent. The parameter K sets the scale for the total srRNA concentration at which the $[RISC]$ component becomes saturated. Straightforward algebraic manipulation yields a characteristic dependence of the expression level of a silenced target on $[srRNA]_{Total}$ as $(A + [srRNA]_{Total}) / (A + B + [srRNA]_{Total})$. The positive amplitudes A and B depend on the model parameters, in a way that becomes messier as one introduces more features into the model. In any case, the underlying parameters are not expected to be known for a particular case. Still, this functional form can easily be identified (or ruled out) if one plots the level of the miRNA target as a function of the concentration of exogenous siRNAs on a double-log scale (Fig. 3).

Alternatively, the rate-limiting enzyme may be Dicer. In this case, one modified the expression for the biogenesis rate of mature miRNA in the model as $\alpha_{miRNA} = \alpha_0 [Dicer] / (K + [srRNA]_{Total})$. This time, the functional form for the dependence of the miRNA target concentration on $[srRNA]_{Total}$ is more complex and non-linear than in the RISC case, a reminiscent of the threshold-linear behavior described above. Plotting this dependence on a double-log scale is quantitatively different, in a way that allows one to discriminate between the two even without knowing any of the underlying parameters (Fig. 3, and unpublished data from our lab).

De-repression by Exogenous Targets - miRNA “Sponges”

Not only saturating levels of srRNA can affect the activity of




Fig. 3. Response of a miRNA target to saturating the level of a key enzyme by over-expressing other srRNAs. The response to limiting Ago takes the form of a ratio between linear polynomials (Hill function with $n=1$). The response to limiting Dicer is much sharper, and cannot be fitted to such function (even with higher n).

RNAi enzymes, but also high levels of targets, as in cases where the srRNA:target complex is long-lived. This is particularly true for bacterial srRNAs that rely on the RNA chaperone Hfq for regulation (40, 104-106). Hfq binds not only the small RNA (making the bacterial equivalent of RISC) but also to the mRNA. Binding of Hfq to the target accelerates the kinetics of srRNA:mRNA binding, and in some cases may play a role in stabilizing the mRNA molecule *in vivo*, perhaps by protecting it from RNaseE-dependent degradation by sequestering its initiation site. Hfq has been reported to exist in the *E.coli* cell at extremely high numbers (on the order of 10,000 copies of hexameric Hfq, similar to the abundance of ribosomes in the fast growing cell) (107). However recent single-molecule data suggest numbers that are order of magnitude smaller (108). Contradicting evidence about the level of Hfq at stationary state and under stress conditions (107), when many srRNAs are activated, make it even harder to determine the relevant level of Hfq. Indeed over-expression an Hfq-binding target can alleviate srRNA repression globally (109). It is straightforward to account for Hfq in the mathematical model (Box 1), in a way similar to our treatment of RISC above. One readily concludes that srRNA targets that experience “efficient” srRNA regulation (due for example to fast mRNA:srRNA binding) would be more sensitive to the level of Hfq. If the abundance of Hfq changes significantly during the life cycle of a bacterial population (e.g. between vegetative growth and steady-state), a regulating small RNA may couple the expression of such genes to the global state of the cell.

In animals, the centrality of miRNAs in many biological pathways motivated the need for miRNA loss-of-function approaches. One such approach is the introduction of a “sponge” mRNA, which contains multiple target sites complementary to the miRNA of interest (110). When the sponge is

expressed at high levels, it specifically inhibits the activity of a whole family of miRNAs that share a common seed. This approach has major advantages over genetic knockout; among which is the fact that srRNA are frequently redundant, and family members may appear in multiple genetic loci.

The efficiency of miRNA sponges has been shown to depend not only on the affinity and avidity of binding sites, but also on the concentration of sponge RNAs relative to the concentration of the miRNA (52). To maximize sponge expression, sponges are cloned behind strong promoters, and high-copy number plasmids are used for transfections.

Introduction of highly abundant miRNA target affects not only the specific miRNA. It may also titrate limiting components of the srRNA pathways, such as Argonaute proteins, resulting in global deregulation of multiple endogenous miRNA and siRNA targets (e.g. in humans, where Ago2 is used in both pathways), as discussed above. It is therefore of merit to limit the expression of the sponge RNA.

The same prototypical model discussed above (Box 1) can be used to consider the effect of a miRNA sponge. First it is noted that the miRNA sponge should compete for all miRNAs, not only the free ones. This distinction may sometimes be elusive both in modeling and in interpreting experimental results, but is critical in cases where a large fraction of the miRNA population is titrated to targets. Effective de-repression requires a match between the synthesis rates of the miRNA sponge and the mature miRNA, not their abundance.

The quantitative model allows one to investigate ways to trade expression rates with other properties. For example, within this simple model, one can readily show that the effect of a miRNA sponge relies on the product ($k_{\text{endogenous target}}/k_{\text{sponge}} \times \alpha_{\text{miRNA}}/\alpha_{\text{sponge}}$), where k denotes the binding rate to the miRNA and α the biogenesis rate. Thus one can trade affinity with expression in a linear fashion.

Conclusions: Unifying View of srRNA Systems Biology

In this mini review we toured multiple small RNA pathways in bacteria and animals to point out the system level aspects of small RNA regulation. While these pathways are very different in principles and in details - they also share some common aspects, especially at the coarse-grained and system level.

We suggest that one should always take a systems approach when studying small RNA regulation. This is due to the fact that post-transcriptional regulation has to act on multiple molecules and species at the same time, thus introducing indirect coupling among targets and between pathways. These indirect interactions may be essential part of the story (as in the examples of chitin and iron metabolism in bacteria), or may just complicate the interpretation of experimental data (as in the case of RNAi and miRNA sponges). Moreover, the non-linear response of an mRNA to its small RNA may complicate the results of large scale experiments meant to identify the targets of a particular srRNA, as srRNAs only have significant effect

when a target is expressed at low levels; highly abundant target is only mildly affected by an srRNA (but may have a large effect on the srRNA itself). Consequently, we suggest that some srRNA:target interactions are there to affect the mRNA, while some are placed to modulate the level of the srRNA.

Attempts to answer fundamental questions, that are true across organisms and pathways, require high-level models. For a given question, it is not clear that such models even exist, as it may be possible that the details that have to be ignored to construct it cannot be omitted. Here, however, it seems that under some simplifying assumptions, a fairly simple mathematical model can surface interesting possibilities and generate testable predictions across a wide range of biological realms.

REFERENCES

1. Jacob, F. and Monod, J. (1961) Genetic regulatory mechanisms in the synthesis of proteins. *J. Mol. Biol.* **3**, 318-356.
2. Lee, R. (1993) The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* **175**, 843-854.
3. Wightman, B., Ha, I. and Ruvkun, G. (1993) Posttranscriptional regulation of the heterochronic gene *lin-14* by *lin-4* mediates temporal pattern formation in *C. elegans*. *Cell* **75**, 855-862.
4. Argaman, L., Hershberg, R., Vogel, J., Bejerano, G., Wagner, E. H., Margalit, H. and Altuvia, S. (2001) Novel small RNA-encoding genes in the intergenic regions of *Escherichia coli*. *Current Biology*. **11**, 941-950.
5. Wassarman, K. M., Repoila, F., Rosenow, C., Storz, G. and Gottesman, S. (2001) Identification of novel small RNAs using comparative genomics and microarrays. *Genes & Development* **15**, 1637-1651.
6. Rivas, E., Klein, R. J., Jones, T. A. and Eddy, S. R. (2001) Computational identification of noncoding RNAs in *E. coli* by comparative genomics. *Current Biology* **11**, 1369-1373.
7. Fire, A., Xu, S., Montgomery, M. K., Kostas, S. A., Driver, S. E. and Mello, C. C. (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* **391**, 806-811.
8. Reinhart, B. J., Slack, F. J., Basson, M., Pasquinelli, A. E., Bettinger, J. C., Rougvie, A. E., Horvitz, H. R. and Ruvkun, G. (2000) The 21-nucleotide *let-7* RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* **403**, 901-906.
9. Pasquinelli, A. E., Reinhart, B. J., Slack, F., Martindale, M. Q., Kuroda, M. I., Maller, B., Hayward, D. C., Ball, E. E., Degnan, B., Müller, P., Spring, J., Srinivasan, A., Fishman, M., Finnerty, J., Corbo, J., Levine, M., Leahy, P., Davidson, E. and Ruvkun, G. (2000) Conservation of the sequence and temporal expression of *let-7* heterochronic regulatory RNA. *Nature* **408**, 86-89.
10. Ghildiyal, M. and Zamore, P. D. (2009) Small silencing RNAs: an expanding universe. *Nat. Rev. Genet.* **10**, 94-108.
11. Bartel, D. P. (2009) MicroRNAs: target recognition and

- regulatory functions. *Cell* **136**, 215-233.
12. Alvarez-Garcia, I. and Miska, E. A. (2005) MicroRNA functions in animal development and human disease. *Development* **132**, 4653-4662.
 13. Hagen, J. W. and Lai, E. C. (2008) microRNA control of cell-cell signaling during development and disease. *Cell Cycle* **7**, 2327-2332.
 14. Ivey, K. N., Muth, A., Arnold, J., King, F. W., Yeh, R., Fish, J. E., Hsiao, E. C., Schwartz, R. J., Conklin, B. R., Bernstein, H. S. and Srivastava, D. (2008) MicroRNA regulation of cell lineages in mouse and human embryonic stem cells. *Cell Stem Cell* **2**, 219-229.
 15. Khurana, J. S. and Theurkauf, W. E. (2008) piRNA function in germline development (July 30, 2008), StemBook, ed. The Stem Cell Research Community, StemBook, doi/10.3824/stembook.1.12.1, <http://www.stembook.org>.
 16. Reynolds, S. and Ruohola-Baker, H. (2008) microRNA's role in germline differentiation (September 15, 2008), Stem Book, ed. The Stem Cell Research Community, StemBook, doi/10.3824/stembook.1.17.1, <http://www.stembook.org>.
 17. Croce, C. M. (2009) Causes and consequences of microRNA dysregulation in cancer. *Nat. Rev. Genet* **10**, 704-714.
 18. Leung, A. K. L. and Sharp, P. A. (2010) MicroRNA functions in stress responses. *Mol. Cell* **40**, 205-215.
 19. Aravin, A. A., Hannon, G. J. and Brennecke, J. (2007) The Piwi-piRNA pathway provides an adaptive defense in the transposon arms race. *Science* **318**, 761-764.
 20. Lee, Y., Jeon, K., Lee, J., Kim, S. and Kim, V. N. (2002) MicroRNA maturation: stepwise processing and subcellular localization. *EMBO J.* **21**, 4663-4670.
 21. Lee, Y., Kim, M., Han, J., Yeom, K., Lee, S., Baek, S. H. and Kim, V. N. (2004) MicroRNA genes are transcribed by RNA polymerase II. *EMBO J.* **23**, 4051-4060.
 22. Czech, B., Malone, C. D., Zhou, R., Stark, A., Schlingeheyde, C., Dus, M., Perrimon, N., Kellis, M., Wohlschlegel, J. A., Sachidanandam, R., Hannon, G. J. and Brennecke, J. (2008) An endogenous small interfering RNA pathway in *Drosophila*. *Nature* **453**, 798-802.
 23. Okamura, K., Chung, W., Ruby, J. G., Guo, H., Bartel, D. P. and Lai, E. C. (2008) The *Drosophila* hairpin RNA pathway generates endogenous short interfering RNAs. *Nature* **453**, 803-806.
 24. Kawamura, Y., Saito, K., Kin, T., Ono, Y., Asai, K., Sunohara, T., Okada, T. N., Siomi, M. C. and Siomi, H. (2008) *Drosophila* endogenous small RNAs bind to Argonaute(thinsp)2 in somatic cells. *Nature* **453**, 793-797.
 25. Okamura, K., Balla, S., Martin, R., Liu, N. and Lai, E. C. (2008) Two distinct mechanisms generate endogenous siRNAs from bidirectional transcription in *Drosophila melanogaster*. *Nat. Struct. Mol. Biol.* **15**, 581-590.
 26. Tam, O. H., Aravin, A. A., Stein, P., Girard, A., Murchison, E. P., Cheloufi, S., Hodges, E., Anger, M., Sachidanandam, R., Schultz, R. M. and Hannon, G. J. (2008) Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature* **453**, 534-538.
 27. Watanabe, T., Totoki, Y., Toyoda, A., Kaneda, M., Kura-mochi-Miyagawa, S., Obata, Y., Chiba, H., Kohara, Y., Kono, T., Nakano, T., Surani, M. A., Sakaki, Y. and Sasaki, H. (2008) Endogenous siRNAs from naturally formed dsRNAs regulate transcripts in mouse oocytes. *Nature* **453**, 539-543.
 28. Ketting, R. F., Haverkamp, T. H. A., van Luenen, H. G. A. M. and Plasterk, R. H. A. (1999) mut-7 of *C. elegans*, Required for Transposon Silencing and RNA Interference, Is a Homolog of Werner Syndrome Helicase and RNaseD. *Cell* **99**, 133-141.
 29. Tabara, H., Sarkissian, M., Kelly, W. G., Fleenor, J., Grishok, A., Timmons, L., Fire, A. and Mello, C. C. (1999) The rde-1 Gene, RNA Interference, and Transposon Silencing in *C. elegans*. *Cell* **99**, 123-132.
 30. Chung, W., Okamura, K., Martin, R. and Lai, E. C. (2008) Endogenous RNA Interference Provides a Somatic Defense against *Drosophila* Transposons. *Current Biology* **18**, 795-802.
 31. Ghildiyal, M., Seitz, H., Horwich, M. D., Li, C., Du, T., Lee, S., Xu, J., Kittler, E. L. W., Zapp, M. L., Weng, Z. and Zamore, P. D. (2008) Endogenous siRNAs derived from transposons and mRNAs in *Drosophila* somatic cells. *Science* **320**, 1077-1081.
 32. Hammond, S. M., Bernstein, E., Beach, D. and Hannon, G. J. (2000) An RNA-directed nuclease mediates post-transcriptional gene silencing in *Drosophila* cells. *Nature* **404**, 293-296.
 33. Zamore, P. D., Tuschl, T., Sharp, P. A. and Bartel, D. P. (2000) RNAi: Double-Stranded RNA Directs the ATP-Dependent Cleavage of mRNA at 21 to 23 Nucleotide Intervals. *Cell* **101**, 25-33.
 34. Pak, J. and Fire, A. (2007) Distinct populations of primary and secondary effectors during RNAi in *C. elegans*. *Science* **315**, 241-244.
 35. Sijen, T., Steiner, F. A., Thijssen, K. L. and Plasterk, R. H. A. (2007) Secondary siRNAs result from unprimed RNA synthesis and form a distinct class. *Science* **2007**, 315, 244-247.
 36. Sijen, T., Fleenor, J., Simmer, F., Thijssen, K. L., Parrish, S., Timmons, L., Plasterk, R. H. and Fire, A. (2001) On the Role of RNA Amplification in dsRNA-Triggered Gene Silencing. *Cell* **107**, 465-476.
 37. Brennecke, J., Aravin, A. A., Stark, A., Dus, M., Kellis, M., Sachidanandam, R. and Hannon, G. J. (2007) Discrete Small RNA-Generating Loci as Master Regulators of Transposon Activity in *Drosophila*. *Cell* **128**, 1089-1103.
 38. Hunter, C. P., Winston, W. M., Molodowitch, C., Feinberg, E. H., Shih, J., Sutherlin, M., Wright, A. J. and Fitzgerald, M. C. (2006) Systemic RNAi in *Caenorhabditis elegans*. *Cold Spring Harb. Symp. Quant. Biol.* **71**, 95-100.
 39. Gottesman, S. (2005) Micros for microbes: non-coding regulatory RNAs in bacteria. *Trends in Genetics* **21**, 399-404.
 40. Aiba, H. (2007) Mechanism of RNA silencing by Hfq-binding small RNAs. *Current Opinion in Microbiology* **10**, 134-139.
 41. Gottesman, S., McCullen, C. A., Guillier, M., Vanderpool, C. K., Majdalani, N., Benhammou, J., Thompson, K. M., Fitzgerald, P. C., Sowa, N. A. and Fitzgerald, D. J. (2006) Small RNA regulators and the bacterial response to stress. *Cold Spring Harb. Symp. Quant. Biol.*

- 71, 1-11.
42. Geissmann, T., Possedko, M., Huntzinger, E., Fechter, P., Ehresmann, C. and Romby, P. (2006) Regulatory RNAs as mediators of virulence gene expression in bacteria. *Handb Exp. Pharmacol.* **173**, 9-43.
 43. Murphy, E. R. and Payne, S. M. (2007) RyhB, an iron-responsive small RNA molecule, regulates *Shigella dysenteriae* virulence. *Infect. Immun.* **75**, 3470-3477.
 44. Padalon-Brauch, G., Hershberg, R., Elgrably-Weiss, M., Baruch, K., Rosenshine, I., Margalit, H. and Altuvia, S. (2008) Small RNAs encoded within genetic islands of *Salmonella typhimurium* show host-induced expression and role in virulence. *Nucleic. Acids. Res.* **36**, 1913-1927.
 45. Schiano, C. A., Bellows, L. E. and Lathem, W. W. (2010) The small RNA chaperone Hfq is required for the virulence of *Yersinia pseudotuberculosis*. *Infect. Immun.* **78**, 2034-2044.
 46. Chabelskaya, S., Gaillot, O. and Felden, B. (2010) A *Staphylococcus aureus* small RNA is required for bacterial virulence and regulates the expression of an immune-evasion molecule. *PLoS Pathog.* **6**, e1000927.
 47. Podkaminski, D. and Vogel, J. (2010) Small RNAs promote mRNA stability to activate the synthesis of virulence factors. *Mol. Microbiol.* **78**, 1327-1331.
 48. Hammond, S. M., Bernstein, E., Beach, D. and Hannon, G. J. (2000) An RNA-directed nuclease mediates post-transcriptional gene silencing in *Drosophila* cells. *Nature* **404**, 293-296.
 49. Hutvagner, G. and Zamore, P. D. (2002) A microRNA in a multiple-turnover RNAi enzyme complex. *Science* **297**, 2056-2060.
 50. Liu, J., Carmell, M. A., Rivas, F. V., Marsden, C. G., Thomson, J. M., Song, J., Hammond, S. M., Joshua-Tor, L. and Hannon, G. J. (2004) Argonaute2 is the catalytic engine of mammalian RNAi. *Science* **305**, 1437-1441.
 51. Haley, B. and Zamore, P. D. (2004) Kinetic analysis of the RNAi enzyme complex. *Nat. Struct. Mol. Biol.* **11**, 599-606.
 52. Arvey, A., Larsson, E., Sander, C., Leslie, C. S. and Marks, D. S. (2010) Target mRNA abundance dilutes microRNA and siRNA activity. *Mol. Syst. Biol.* **6**, 363.
 53. Massé, E., Escorcia, F. E. and Gottesman, S. (2003) Coupled degradation of a small regulatory RNA and its mRNA targets in *Escherichia coli*. *Genes Dev.* **17**, 2374-2383.
 54. Figueroa-Bossi, N., Valentini, M., Malleret, L. and Bossi, L. (2009) Caught at its own game: regulatory small RNA inactivated by an inducible transcript mimicking its target. *Genes & Development* **23**, 2004-2015.
 55. Levine, E., Zhang, Z., Kuhlman, T. and Hwa, T. (2007) Quantitative characteristics of gene regulation by small RNA. *PLoS Biol.* **5**, e229.
 56. Mitarai, N., Andersson, A. M., Krishna, S., Semsey, S. and Sneppen, K. (2007) Efficient degradation and expression prioritization with small RNAs. *Phys. Biol.* **4**, 164-171.
 57. Shimonj, Y., Friedlander, G., Hetzroni, G., Niv, G., Altuvia, S., Biham, O. and Margalit, H. (2007) Regulation of gene expression by small non-coding RNAs: a quantitative view. *Mol. Syst. Biol.* **3**, 138.
 58. Mehta, P., Goyal, S. and Wingreen, N. S. (2008) A quantitative comparison of sRNA-based and protein-based gene regulation. *Mol. Syst. Biol.* **4**, 221.
 59. Levine, E., Huang, M., Huang, Y., Kuhlman, T., Zhang, Z. and Hwa, T. On noise and silence in gene regulation by small RNA. In submission.
 60. Lease, R. A. and Belfort, M. (2000) A trans-acting RNA as a control switch in *Escherichia coli*: DsrA modulates function by forming alternative structures. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 9919-9924.
 61. Fang, F. C. and Rimsky, S. (2008) New insights into transcriptional regulation by H-NS. *Current Opinion in Microbiology.* **11**, 113-120.
 62. Amit, R., Oppenheim, A. B. and Stavans, J. (2003) Increased Bending Rigidity of Single DNA Molecules by H-NS, a Temperature and Osmolarity Sensor. *Biophysical Journal* **84**, 2467-2473.
 63. Dorman, C. J. (2007) H-NS, the genome sentinel. *Nat. Rev. Micro.* **5**, 157-161.
 64. Majdalani, N., Cuning, C., Sledjeski, D., Elliott, T. and Gottesman, S. (1998) DsrA RNA regulates translation of RpoS message by an anti-antisense mechanism, independent of its action as an antisilencer of transcription. *Proc. Natl. Acad. Sci. U.S.A.* **95**, 12462-12467.
 65. Zhang, A., Altuvia, S., Tiwari, A., Argaman, L., Hengge-Aronis, R. and Storz, G. (1998) The OxyS regulatory RNA represses rpoS translation and binds the Hfq (HF-I) protein. *EMBO J.* **17**, 6061-6068.
 66. Basineni, S. R., Madhugiri, R., Kolmsee, T., Hengge, R. and Klug, G. (2009) The influence of Hfq and ribonucleases on the stability of the small non-coding RNA OxyS and its target rpoS in *E. coli* is growth phase dependent. *RNA Biol.* **6**, 584-594.
 67. Repoila, F., Majdalani, N. and Gottesman, S. (2003) Small non-coding RNAs, co-ordinators of adaptation processes in *Escherichia coli*: the RpoS paradigm. *Mol. Microbiol.* **48**, 855-861.
 68. Madhugiri, R., Basineni, S. R. and Klug, G. (2010) Turnover of the small non-coding RNA RprA in *E. coli* is influenced by osmolarity. *Mol. Genet. Genomics.* **284**, 307-318.
 69. Repoila, F. and Gottesman, S. (2001) Signal Transduction Cascade for Regulation of RpoS: Temperature Regulation of DsrA. *J. Bacteriol.* **183**, 4012-4023.
 70. Repoila, F. and Gottesman, S. (2003) Temperature Sensing by the dsrA Promoter. *J. Bacteriol.* **185**, 6609-6614.
 71. Lenz, D., Mok, K., Lilley, B., Kulkarni, R., Wingreen, N. and Bassler, B. (2004) The small RNA chaperone Hfq and multiple small RNAs control quorum sensing in *Vibrio harveyi* and *Vibrio cholerae*. *Cell* **118**, 69-82.
 72. Tu, K. C. and Bassler, B. L. (2007) Multiple small RNAs act additively to integrate sensory information and control quorum sensing in *Vibrio harveyi*. *Genes Dev.* **21**, 221-233.
 73. Svenningsen, S. L., Tu, K. C. and Bassler, B. L. (2009) Gene dosage compensation calibrates four regulatory RNAs to control *Vibrio cholerae* quorum sensing. *EMBO*

- J.* **28**, 429-439.
74. Long, T., Tu, K. C., Wang, Y., Mehta, P., Ong, N. P., Bassler, B. L. and Wingreen, N. S. (2009) Quantifying the integration of quorum-sensing signals with single-cell resolution. *PLoS Biol.* **7**, e68.
 75. Thomas, M., Lieberman, J. and Lal, A. (2010) Desperately seeking microRNA targets. *Nat. Struct. Mol. Biol.* **17**, 1169-1174.
 76. Baek, D., Villén, J., Shin, C., Camargo, F. D., Gygi, S. P. and Bartel, D. P. (2008) The impact of microRNAs on protein output. *Nature* **455**, 64-71.
 77. Selbach, M., Schwanhauser, B., Thierfelder, N., Fang, Z., Khanin, R. and Rajewsky, N. (2008) Widespread changes in protein synthesis induced by microRNAs. *Nature* **455**, 58-63.
 78. Stark, A., Brennecke, J., Bushati, N., Russell, R. B. and Cohen, S. M. (2005) Animal MicroRNAs confer robustness to gene expression and have a significant impact on 3'UTR evolution. *Cell* **123**, 1133-1146.
 79. Jan, C. H., Friedman, R. C., Ruby, J. G. and Bartel, D. P. (2010) Formation, regulation and evolution of *Caenorhabditis elegans* 3'UTRs. *Nature* doi:10.1038/nature09616.
 80. Overgaard, M., Johansen, J., Møller-Jensen, J. and Valentin-Hansen, P. (2009) Switching off small RNA regulation with trap-mRNA. *Mol. Microbiol.* **73**, 790-800.
 81. Plumbridge, J. and Pellegrini, O. (2004) Expression of the chitobiose operon of *Escherichia coli* is regulated by three transcription factors: NagC, ChbR and CAP. *Mol. Microbiol.* **52**, 437-449.
 82. Nowojewski, A. and Levine, E. Manuscript in preparation.
 83. Massé, E. and Gottesman, S. (2002) A small RNA regulates the expression of genes involved in iron metabolism in *Escherichia coli*. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 4620-4625.
 84. Mey, A. R., Craig, S. A. and Payne, S. M. (2005) Characterization of *Vibrio cholerae* RyhB: the RyhB regulon and role of RyhB in biofilm formation. *Infect. Immun.* **73**, 5706-5719.
 85. Massé, E., Vanderpool, C. K. and Gottesman, S. (2005) Effect of RyhB small RNA on global iron use in *Escherichia coli*. *J. Bacteriol.* **187**, 6962-6971.
 86. Jacques, J., Jang, S., Prévost, K., Desnoyers, G., Desmarais, M., Imlay, J. and Massé, E. (2006) RyhB small RNA modulates the free intracellular iron pool and is essential for normal growth during iron limitation in *Escherichia coli*. *Mol. Microbiol.* **62**, 1181-1190.
 87. Wyckoff, E. E., Mey, A. R. and Payne, S. M. (2007) Iron acquisition in *Vibrio cholerae*. *Biometals* **20**, 405-416.
 88. Prévost, K., Salvail, H., Desnoyers, G., Jacques, J., Phaneuf, E. and Massé, E. (2007) The small RNA RyhB activates the translation of *shiA* mRNA encoding a permease of shikimate, a compound involved in siderophore synthesis. *Mol. Microbiol.* **64**, 1260-1273.
 89. Vecerek, B., Moll, I. and Bläsi, U. (2007) Control of Fur synthesis by the non-coding RNA RyhB and iron-responsive decoding. *EMBO J.* **26**, 965-975.
 90. Semsey, S., Andersson, A. M. C., Krishna, S., Jensen, M. H., Massé, E. and Snieppen, K. (2006) Genetic regulation of fluxes: iron homeostasis of *Escherichia coli*. *Nucleic Acids. Res.* **34**, 4960-4967.
 91. Andrews, S. C., Robinson, A. K. and Rodríguez-Quiriones, F. (2003) Bacterial iron homeostasis. *FEMS Microbiol. Rev.* **27**, 215-237.
 92. Ambros, V. (1989) A hierarchy of regulatory genes controls a larva-to-adult developmental switch in *C. elegans*. *Cell* **57**, 49-57.
 93. Slack, F. J., Basson, M., Liu, Z., Ambros, V., Horvitz, H. R. and Ruvkun, G. (2000) The *lin-41* RBCC gene acts in the *C. elegans* heterochronic pathway between the *let-7* regulatory RNA and the *LIN-29* transcription factor. *Mol. Cell* **5**, 659-669.
 94. Ketting, R. F., Fischer, S. E., Bernstein, E., Sijen, T., Hannon, G. J. and Plasterk, R. H. (2001) Dicer functions in RNA interference and in synthesis of small RNA involved in developmental timing in *C. elegans*. *Genes & Development* **15**, 2654-2659.
 95. Grishok, A., Pasquinelli, A. E., Conte, D., Li, N., Parrish, S., Ha, I., Baillie, D. L., Fire, A., Ruvkun, G. and Mello, C. C. (2001) Genes and Mechanisms Related to RNA Interference Regulate Expression of the Small Temporal RNAs that Control *C. elegans* Developmental Timing. *Cell* **106**, 23-34.
 96. Lee, Y. S., Nakahara, K., Pham, J. W., Kim, K., He, Z., Sontheimer, E. J. and Carthew, R. W. (2004) Distinct Roles for *Drosophila* Dicer-1 and Dicer-2 in the siRNA/miRNA Silencing Pathways. *Cell* **117**, 69-81.
 97. Tolia, N. H. and Joshua-Tor, L. (2007) Slicer and the Argonautes. *Nat. Chem. Biol.* **3**, 36-43.
 98. Jackson, A. L. and Linsley, P. S. (2010) Recognizing and avoiding siRNA off-target effects for target identification and therapeutic application. *Nat. Rev. Drug Discov.* **9**, 57-67.
 99. Jackson, A. L., Bartz, S. R., Schelter, J., Kobayashi, S. V., Burchard, J., Mao, M., Li, B., Cavet, G. and Linsley, P. S. (2003) Expression profiling reveals off-target gene regulation by RNAi. *Nat. Biotechnol.* **21**, 635-637.
 100. Yi, R., Doehle, B. P., Qin, Y., Macara, I. G. and Cullen, B. R. (2005) Overexpression of Exportin 5 enhances RNA interference mediated by short hairpin RNAs and microRNAs. *RNA* **11**, 220-226.
 101. Grimm, D., Streetz, K. L., Jopling, C. L., Storm, T. A., Pandey, K., Davis, C. R., Marion, P., Salazar, F. and Kay, M. A. (2006) Fatality in mice due to oversaturation of cellular microRNA/short hairpin RNA pathways. *Nature* **441**, 537-541.
 102. Khan, A. A., Betel, D., Miller, M. L., Sander, C., Leslie, C. S. and Marks, D. S. (2009) Transfection of small RNAs globally perturbs gene regulation by endogenous microRNAs. *Nat. Biotech.* **27**, 549-555.
 103. Larsson, E., Sander, C. and Marks, D. (2010) mRNA turnover rate limits siRNA and microRNA efficacy. *Mol. Syst. Biol.* **6**, 433.
 104. Sittka, A., Lucchini, S., Papenfort, K., Sharma, C. M., Rolfe, K., Binnewies, T. T., Hinton, J. C. D. and Vogel, J. (2008) Deep sequencing analysis of small noncoding RNA and mRNA targets of the global post-transcriptional regulator, Hfq. *PLoS Genet.* **4**, e1000163.
 105. Jousselin, A., Metzinger, L. and Felden, B. (2009) On the facultative requirement of the bacterial RNA chaperone, Hfq. *Trends. Microbiol.* **17**, 399-405.

106. Le Derout, J., Boni, I. V., Régnier, P. and Hajsndorf, E. (2010) Hfq affects mRNA levels independently of degradation. *BMC Mol. Biol.* **11**, 17.
107. Valentin-Hansen, P., Eriksen, M. and Udesen, C. (2004) The bacterial Sm-like protein Hfq: a key player in RNA transactions. *Mol. Microbiol.* **51**, 1525-1533.
108. Taniguchi, Y., Choi, P. J., Li, G., Chen, H., Babu, M., Hearn, J., Emili, A. and Xie, X. S. (2010) Quantifying *E. coli* proteome and transcriptome with single-molecule sensitivity in single cells. *Science* **329**, 533-538.
109. Hussein, R. and Lim, H. N. (2010) Disruption of small RNA signaling caused by competition for Hfq. *Proc. Natl. Acad. Sci. U.S.A.* doi:10.1073/pnas.1010082108.
110. Ebert, M. S. and Sharp, P. A. (2010) MicroRNA sponges: Progress and possibilities. *RNA* **16**, 2043-2050.