

A Lichen Genus *Porpidia* (Porpidiaceae) from South Korea

Xin Yu Wang, Yogesh Joshi and Jae-Seoun Hur*

Korean Lichen Research Institute, Suncheon National University, Suncheon 540-742, Korea

(Received January 24, 2011. Accepted February 18, 2011)

Taxonomic study of the crustose lichen *Porpidia* was performed in this paper. Three species are described, including two recorded species and one new record: *Porpidia macrocarpa* (DC.) Hertel & A. J. Schwab. A description of each species is presented along with morphological, anatomic, and chemical characteristics. A key to the identification of species of *Porpidia* is also presented.

KEYWORDS: Lichens, New record, *Porpidia macrocarpa*, Taxonomy

The crustose lichen genus *Porpidia* Körber is a saxicolous lichen that is widely distributed in the mountains of South Korea. The genus was firstly described by Körber [1], and it belonged to the classical family Lecideaceae and classical genus *Lecidea* [2] for a long time. It was later segregated from *Lecidea* in 1975 as the genus name *Huilia* Zahlbr. by Hertel [3], and in 1984, the same author [4] resurrected the genus name *Porpidia* as the correct name for the genus *Huilia*.

The species of *Porpidia* occur mostly on siliceous rocks, although a few are confirmed on bark, lignum, worked timber, and consolidated soil [5]. Due to their growth on rock, *Porpidia* spp. are difficult to collect, such that the number of specimens is relatively fewer than that of macro-lichens. These share some common characteristics: thallus endolithic to epilithic, 0.1 to 1.5 mm thick; continuous thalli almost completely uncracked; thallus gray to ashy gray or orange; apothecia usually sessile on thallus, disk pruinose or not, spores ellipsoid and covered with a hyaline halo, ascus tip with a dark blue tubular structure when stained in iodine solution.

Two species, *P. albocaerulescens* and *P. crustulata*, have been reported in South Korea [6]. Except for a checklist, no taxonomic study on this genus has been carried out, and therefore this work focused on the detailed taxonomic analysis of this genus from South Korea. One hundred and twenty specimens were collected from the main mountains of South Korea during 2003 to 2010. Two species, *P. albocaerulescens* and *P. crustulata*, were found during our study, together with another reported species. In total, three species are included in this paper with a key for the identification of species of *Porpidia*.

Specimens were examined using standard microscopical techniques and hand-sectioned under a Nikon SMZ645

dissecting microscope (Nikon, Tokyo, Japan). All measurements were made on material mounted in water, amyloid reactions were tested with iodine solution, and lactophenol cotton blue was used as a stain. Nikon Coolpix 4500 was used for taking photographs of the species. Thin layer chromatography was performed in solvent system C (toluene : acetic acid = 85 : 15) as described by Elix *et al.* [7] and White and James [8]. The specimens were lodged at the herbarium of the Lichen & Allied Bioresource Center, Korean Lichen Research Institute (KoLRI), Suncheon National University, Korea.

Taxonomic Treatment of the Species

Key to the genus *Porpidia* in South Korea

1. Apothecia disk heavily pruinose, thallus thick (0.5~1.2 mm) *P. albocaerulescens*
1. Apothecia non-pruinose, thallus thin 2
2. Apothecia less than 1 mm in diam., hymenium 60~90 µm high, spores 10~17 µm *P. crustulata*
2. Apothecia 1~3 mm in diam., hymenium 80~120 µm high, spores 16~25 µm *P. macrocarpa*

The Species

Porpidia albocaerulescens (Wulfen) Hertel & Knoph

Diagnostic characters. Thallus grayish green to whitish, sometimes dark to olive green when wet, continuous, slightly cracked when dry, 0.3~1.2 mm thick, surface even, marginal part usually thinner than the thallus center. Prothallus black, rather obvious when two individual thallus abut each other. Apothecia abundant, clustered and sessile, but sunken in thallus when young, 0.5~1.5 mm in diameter when mature, disk black, covered with white pruina. Hymenium 70~110 µm high, epithecium brown to dark brown, subhymenium 20~40 µm high, exciple dark brown

*Corresponding author <E-mail : jshru1@suncheon.ac.kr>

to blackish in marginal part, lighter brown within. Spores ellipsoid, $17\sim 25 \times 6\sim 10 \mu\text{m}$.

Chemistry. Stictic acid (main), cryptostictic acid, constictic acid, and norstictic acid.

Habitat. On HCl-rocks, particularly in shady and humid areas.

South Korean distribution. Rather common, widely distributed all over South Korea, found in altitudes from sea level up to 1,600 m.

World distribution. Eastern to Southeastern Asia, Europe, and North America [5, 9, 10].

Remarks. The species was easy to recognize even in the field. It is the only pruinose *Porpidia* species in South Korea. By having a thick and smooth thallus, as well as large apothecia and thick whitish pruina, it is easy to separate from other crustose lichen species in South Korea.

Representative specimens examined. Mt. Halla, Jeju Island, $33^{\circ}22'47.3''$ N, $126^{\circ}33'43.0''$ E, alt. 1,270 m, Hur090128. Bogil Island Jeonnam Prov., $34^{\circ}08'77.8''$ N, $126^{\circ}32'77.7''$ E, alt. 2 m, Hur100003. Mt. Jiri, Hadong-gun, Gyeongnam Prov., $35^{\circ}18'355''$ N, $127^{\circ}35'214''$ E, alt. 1,473 m, Hur091183. Mt. Duta, Pyeongchang-gun, Gangwon Prov., $37^{\circ}33'94.3''$ N, $128^{\circ}35'02.5''$ E, alt. 355 m, Hur100691. Mt. Suri, Anyang City, Gyeonggi Prov., $37^{\circ}22'004''$ N, $126^{\circ}53'612''$ E, alt. 194 m, Hur101224.

***Porpidia crustulata* (Ach.) Hertel & Knoph**

Remarks. This species was not found during our study, but it has been reported by Moon [11] on Mt. Sorak. Unfortunately, we did not find the species after several field surveys on Mt. Sorak.

The species is characterized by having thin thallus (less than 0.5 mm), short hymenium ($60\sim 90 \mu\text{m}$ high), small spores ($10\sim 17 \mu\text{m}$ long), small and numerous apothecia (less than 1 mm in diameter), and non-pruinose black disk. It is similar with *P. macrocarpa* but differs in having smaller spore size, hymenium height, and apothecia size.

South Korean distribution. Rare, reported only on Mt. Sorak [11].

World distribution. It has been reported all over the world, from temperate to alpine or arctic climates [10].

***Porpidia macrocarpa* (DC.) Hertel & A. J. Schwab (Fig. 1A~1C)**

Diagnostic characters. Thallus greenish gray to whitish gray, sometimes partly orange, epilithic or less frequently




Fig. 1. Morphological characteristics and distribution of *Porpidia macrocarpa*. A, Habit of *Porpidia macrocarpa* (DC.) Hertel & A. J. Schwab; B, Exciple section of *P. macrocarpa*; C, Distribution of *P. macrocarpa* in South Korea.

endolithic, smooth or rugulose in some old parts, rather thin, 0.1 to 0.5 mm in diameter. Clear black prothallus present in the marginal part. Apothecia abundant and clustered, sessile in mature thallus, sunken when young, $0.5\sim 2.0$ (~ 3.0) mm in diameter, black or dark brown, disk non-pruinose, usually flat. Hymenium $70\sim 100 \mu\text{m}$ high, epithecium usually olive brown, subhymenium $20\sim 40 \mu\text{m}$ high, exciple composed of elongated cells radiating from hypothecium, dark brown to blackish in margin and paler within. Spores ellipsoid, $18\sim 23 \times 6\sim 9 \mu\text{m}$.

Chemistry. Stictic acid and cryptostictic acid (all accessory); or no compound.

Habitat. HCl-rock, on exposed but humid area.

South Korean distribution. Rare, it is found only in Gangwon province at an altitude around $800\sim 1,200$ m.

World distribution. Asia, Europe, and North America [9, 10, 12].

Remarks. The species might be confused with *P. crustulata* when its apothecia are small, but it has a higher

hymenium and larger spores (usually around 20 µm long), whereas *P. crustulata* has spores always shorter than 17 µm. Some *P. albocaerulescens* specimens with unclear pruina on disk might be confusing, but *P. macrocarpa* has much thinner thallus, smaller spores, and darker exciple.

Specimens examined. Mt. Eungbok, Hongcheon-gun, Gangwon Prov., 37°51'35.9" N, 128°30'9.74" E, alt. 1,192 m, Hur090664. Mt. Jang, Sangdong-eup, Yeongwol-gun, Gangwon Prov., 37°08'38.7" N, 128°51'04.2" E, alt. 762 m, Hur100866.

Acknowledgements

This work was supported by a grant from the Korea National Research Resource Center Program (Grant 2010-0000660) and the Korean Forest Service Program (KNA 2010) through the Korea National Arboretum.

References

1. Körber GW. Systema Lichenum Germaniae. Breslau: Trendt & Granier; 1855.
2. Zahlbruckner A. Lichenes (Flechten). B. Spezieller Teil. In: Engler A, Prantl K, editors. Die Natürlichen Pflanzenfamilien, Vol. 8. Leipzig: W. Engelmann; 1926. p. 61-270.
3. Hertel H. Beiträge zur Kenntnis der Flechtenfamilie Lecideaceae VI. Herzogia 1975;3:365-406.
4. Hertel H. Über saxicole, lecideoide Flechten der Subantarktis. Nova Hedwigia Beih 1984;79:399-499.
5. Inoue M. Japanese species of *Huilia* (Lichenes) (1-3). Jpn Bot 1983;58:113-28, 161-73, 225-36.
6. Hur JS, Koh YJ, Harada H. A checklist of Korean lichens. Lichenology 2005;4:65-95.
7. Elix JA, Johnston J, Parker JL. A catalogue of standardized thin layer chromatographic data and biosynthetic relationships for lichen substances. 2nd ed. Canberra: Australian National University; 1987.
8. White FJ, James PW. A new guide to microchemical techniques for the identification of lichen substances. Br Lichen Soc Bull (Suppl) 1985;57:1-41.
9. Gowan SP. The lichen genus *Porpidia* (Porpidiaceae) in North America. Bryologist 1989;92:25-59.
10. Hertel H. Gesteinsbewohnende Arten der Sammelgattung *Lecidea* (Lichenes) aus Zentral-, Ost-, und Südasien. Khumbu Himal 1977;6:145-378.
11. Moon KH. Lichens of Mt. Sorak in Korea. J Hattori Bot Lab 1999;86:187-220.
12. Knoph JG. Vorarbeiten zu einer Monographie der euthallinen Arten der Flechtengattung *Porpidia* (Porpidiaceae, Lecanorales) Europas, mit besonderer Berücksichtigung des Alpengebietes. München: Institut für Systematische Botanik der Universität München; 1984.