

# Phospholipase Activities in Clinical and Environmental Isolates of *Acanthamoeba*

Abdul Matin<sup>1</sup> and Suk-Yul Jung<sup>2,\*</sup>

<sup>1</sup>Institute of Biomedical and Genetic Engineering, 24 Mauve Area, Sector: G-9/1, Islamabad, Pakistan; <sup>2</sup>Department of Biomedical Laboratory Science, Molecular Diagnosis Research Institute, Namseoul University, Cheonan 331-707, Korea

**Abstract:** The pathogenesis and pathophysiology of *Acanthamoeba* infections remain incompletely understood. Phospholipases are known to cleave phospholipids, suggesting their possible involvement in the host cell plasma membrane disruption leading to host cell penetration and lysis. The aims of the present study were to determine phospholipase activities in *Acanthamoeba* and to determine their roles in the pathogenesis of *Acanthamoeba*. Using an encephalitis isolate (T1 genotype), a keratitis isolate (T4 genotype), and an environmental isolate (T7 genotype), we demonstrated that *Acanthamoeba* exhibited phospholipase A<sub>2</sub> (PLA<sub>2</sub>) and phospholipase D (PLD) activities in a spectrophotometry-based assay. Interestingly, the encephalitis isolates of *Acanthamoeba* exhibited higher phospholipase activities as compared with the keratitis isolates, but the environmental isolates exhibited the highest phospholipase activities. Moreover, *Acanthamoeba* isolates exhibited higher PLD activities compared with the PLA<sub>2</sub>. *Acanthamoeba* exhibited optimal phospholipase activities at 37°C and at neutral pH indicating their physiological relevance. The functional role of phospholipases was determined by in vitro assays using human brain microvascular endothelial cells (HBMEC), which constitute the blood-brain barrier. We observed that a PLD-specific inhibitor, i.e., compound 48/80, partially inhibited *Acanthamoeba* encephalitis isolate cytotoxicity of the host cells, while PLA<sub>2</sub>-specific inhibitor, i.e., cytidine 5'-diphosphocholine, had no effect on parasite-mediated HBMEC cytotoxicity. Overall, the T7 exhibited higher phospholipase activities as compared to the T4. In contrast, the T7 exhibited minimal binding to, or cytotoxicity of, HBMEC.

**Key words:** *Acanthamoeba*, human brain microvascular endothelial cell, phospholipase A2, phospholipase D, adhesion, cytotoxicity

## INTRODUCTION

*Acanthamoeba* is a free-living protozoan and has been isolated from diverse environments, including air, soil, tap water, or swimming pools [1]. It is recognized as one of the most ubiquitous protozoans [1]. *Acanthamoeba* is an opportunistic human pathogen and can cause a rare form of granulomatous encephalitis which almost always results in death or a common keratitis with consequences of blindness [2-4]. However, the pathogenesis and pathophysiology associated with *Acanthamoeba* infections remain incompletely understood. *Acanthamoeba* has been shown to demonstrate several extracellular toxins, particularly extracellular proteases which might contribute towards its pathogenicity. Previous studies have shown the presence of phospholipases in *Acanthamoeba*; however, our knowl-

edge of the role of phospholipases in the virulence of *Acanthamoeba* is unclear [5-7]. Besides their role in remodelling of the plasma membrane of the parasite itself, based on the fact that phospholipases are known to cleave phospholipids, they may involve in the host cell plasma membrane disruptions leading to penetration of the host cells, and cell lysis [5-7]. This suggests that an understanding of the phospholipases in the biology and pathogenesis of *Acanthamoeba* may identify targets for therapeutic interventions.

Phospholipases are a diverse group of enzymes that hydrolyze ester linkage in glycerophospholipids and are involved in the biosynthesis and degradation of membrane lipids. Thus, the activities of phospholipases can result in membrane dysfunction. The 5 major known phospholipases are A<sub>1</sub>, A<sub>2</sub>, B, C, and D, each having the ability to cleave a specific ester bond in the phospholipid substrate of the target membrane [8]. For example, PLA<sub>1</sub> hydrolyses the fatty acyl ester bond at the *sn*-1 position of the glycerol moiety, while PLA<sub>2</sub> removes the fatty acid at the *sn*-2 position of glycerol [8]. The results of both the PLA<sub>1</sub> and PLA<sub>2</sub> action are free fatty acids or acyl lysophospholipid. The latter are further cleaved by the action of lysophospholi-

• Received 29 September 2010, revised 30 November 2010, accepted 16 December 2010.

\* Corresponding author (syjung@nsu.ac.kr)

© 2011, Korean Society for Parasitology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

pases. Phospholipase B has both hydrolase activities, such as cleaving fatty acids from phospholipids and lysophospholipids as well as acyltransferase activity, for instance, the production of phospholipids by transferring a free fatty acid to a lysophospholipid [8]. Phospholipase C hydrolyses phosphodiester bond to yield 1,2-diacylglycerol depending on the phospholipid class involved, while PLD hydrolyses the phosphodiester bond between the phosphate and the polar head group, producing phosphatidic acid and choline [8-10]. All phospholipases are present in multiple forms. For example, approximately 20 different PLA<sub>2</sub> enzymes have been identified in mammalian tissues and have been broadly classified into 3 families based on their calcium requirement for catalytic activity [8,11,12]. Different parts of the breakages by PLA<sub>2</sub> and PLD are focused on this study. Victoria and Korn [6] observed that PLA of *Acanthamoeba castellanii* was located in the plasma membrane [6]. It may be affected on binding of *Acanthamoeba* to target cells. The aims of the present study were to determine PLA<sub>2</sub> and PLD activities in *Acanthamoeba* and to determine their role in *Acanthamoeba* binding to, and cytotoxicity of, human brain microvascular endothelial cells (HBMEC).

## MATERIALS AND METHODS

### Culture of *Acanthamoeba*

All chemicals were purchased from Sigma Laboratories (Poole, Dorset, UK), unless otherwise stated. The following *Acanthamoeba* isolates of different pathogenicity were used; (1) a clinical isolate of *A. castellanii* belonging to T4 genotype, isolated from a keratitis patient (ATCC 50492) (American Type Culture Collection, Manassas, USA), (2) a clinical isolate of *A. castellanii* belonging to T1 genotype, isolated from an encephalitis patient (ATCC 50494), and (3) an environmental isolate of *Acanthamoeba astronyxis* belonging to T7 genotype, isolated from soil (ATCC 30137). All amoebae were grown in 15 ml of PYG medium (proteose peptone 0.75% w/v, yeast extract 0.75% w/v, and glucose 1.5% w/v) at 30°C and the media was refreshed 17-20 hr prior to experiments as previously described [13]. This resulted in more than 99% amoebae in trophozoite forms, which were used for subsequent experiments.

### Human brain microvascular endothelial cells (HBMEC)

The primary HBMEC were isolated from the human brain tissue and cultured as previously described [14,15]. In brief, HBMEC were purified by fluorescent activated cell sorting (FA-

CS) and tested for endothelial characteristics, such as the expression of endothelial markers, F-VIII, carbonic anhydrase IV, and the uptake of acetylated low density lipoprotein (AcLDL), resulting in more than 99% pure cultures. HBMEC were routinely grown on rat tail collagen-coated dishes in RPMI-1640 containing 10% heat inactivated fetal bovine serum, 10% Nu serum, 2 mM glutamine, 1 mM sodium pyruvate, penicillin (100 U/ml), streptomycin (100 U/ml), non-essential amino acids, and vitamins [14,15].

### Adhesion assays

Adhesion assays were performed as previously described [4]. Briefly, HBMEC were grown to confluency in 24-well plates. *Acanthamoeba* isolates ( $2 \times 10^5$  amoebae/0.5 ml/well) were incubated with cell monolayers in RPMI-1640 containing 2 mM glutamine, 1 mM sodium pyruvate, and non-essential amino acids. The plates were incubated at 37°C in a 5% CO<sub>2</sub> incubator. After 60 min incubations, the numbers of unbound amoebae in the supernatants were determined by hemocytometer counting. The percentage of bound amoebae was calculated as follows:  $\text{Number of unbound amoebae} / \text{Total number of amoebae} \times 100 = \% \text{ unbound amoebae}$ . The numbers of bound amoebae were deduced as follows:  $100 - \% \text{ unbound amoebae} = \% \text{ bound amoebae}$ .

To determine the role of PLA<sub>2</sub> and PLD in *Acanthamoeba* adhesion to the HBMEC, assays were performed in the presence of phospholipase inhibitors. Briefly, amoebae were pre-incubated with the phospholipase inhibitors for 45 min at room temperature. Following this incubation, amoebae plus inhibitors were transferred to HBMEC monolayers for adhesion assays as described above. The following phospholipase inhibitors were used: cytidine 5'-diphosphocholine, i.e., PLA<sub>2</sub> inhibitor; and compound 48/80, i.e., PLC/PLD inhibitor.

### Cytotoxicity assay

To determine the role of phospholipases in *Acanthamoeba*-mediated HBMEC death, cytotoxicity assays were performed as previously described [13]. In brief, amoebae in the presence or absence of phospholipase inhibitors were added to HBMEC monolayers as described for adhesion assays. The plates were observed periodically for monolayer disruptions under a phase-contrast microscope for up to 24 hr. Following this incubation, the supernatants were collected and examined for host cell cytotoxicity by measuring lactate dehydrogenase (LDH) release (cytotoxicity detection kit; Roche Applied Science, Lewes, East

Sussex, UK). In brief, the supernatants of co-cultures of *Acanthamoeba* and the host cells were assessed for the presence of LDH, the release of which is considered as an estimate of cell death. The percentage LDH release was calculated as follows:  $(\text{LDH activity in experimental sample as measured by optical density at 492 nm} - \text{LDH activity in control samples}) / (\text{total LDH activity release} - \text{LDH activity in control samples}) \times 100 = \% \text{ cytotoxicity}$ . Control samples were obtained from host cells or *Acanthamoeba* incubated alone. Total LDH activity release was determined by total host cell lysis with 1% Triton X-100 for 30 min at 37°C.

#### Phospholipase A<sub>2</sub> assay

To determine the activity of phospholipases, the characteristics of L- $\alpha$ -glycerophosphate oxidase for PLA<sub>2</sub> and choline oxidase for PLD were used. *Acanthamoeba* PLA<sub>2</sub> activities were determined using parasite lysates and their conditioned medium in a spectrophotometric-based assay. To prepare lysates, amoebae were lysed using freeze-thaw method as previously described [16]. In brief, various numbers of amoebae were resuspended in 100  $\mu$ l of PBS, followed by freeze-thaw for 3 times. To prepare the conditioned medium, *Acanthamoeba* were incubated in RPMI-1640 for 24 hr. Next day, the cell-free conditioned medium was collected by centrifugation and used for PLA<sub>2</sub> assays. A 100  $\mu$ l of amoebae lysate or conditioned medium was incubated with 900  $\mu$ l of PLA<sub>2</sub> substrate solution (0.2 M D, L- $\alpha$ -glycerophosphate, 0.1% 4-aminoantipyrine, 0.1% phenol, and 0.025% peroxidase dissolved in 0.125 M Tris-HCl buffer, pH 8.0). The reaction mixtures were incubated for 2 hr. After this incubation, the mixtures were centrifuged at 900 g for 5 min and supernatants were collected and their optical density determined at 500 nm. The PLA<sub>2</sub> activity was calculated as follows:  $(\text{OD test} - \text{OD blank}) \times V_t \times df / 13.3 \times 0.5 \times t \times 1.0 \times V_s = \text{enzymatic activity (units per ml)}$ . Here,  $V_t$  is total volume;  $V_s$  is sample volume; 13.3 is the millimolar extinction coefficient of quinoneimine dye under the assay condition; 0.5 is the factor based on the fact that 1 mole of H<sub>2</sub>O<sub>2</sub> produces half a mole of quinoneimine dye;  $t$  is reaction time; 1.0 is light path (cm); and  $df$  is the dilution factor. To determine the optimal activity, PLA<sub>2</sub> assays were performed using various numbers of amoebae at different pH and temperatures.

#### Phospholipase D assay


*Acanthamoeba* PLD activities were determined using parasite lysates and their conditioned medium in a spectrophotomet-

ric-based assay. The lysates and conditioned medium were prepared as described above. A 100  $\mu$ l of amoebae lysate or conditioned medium was incubated with 900  $\mu$ l of PLD substrate solution (2.1% choline chloride, 1% 4-aminoantipyrine, 1% phenol, and 5 mg peroxidase in 0.1 M Tris-HCl buffer, pH 8.0). The reaction mixtures were incubated for 2 hr. After this incubation, the mixtures were centrifuged at 900 g for 5 min, and supernatants were collected and their optical density determined at 500 nm. The PLD activity was calculated as follows:  $(\text{OD test} - \text{OD blank}) / \text{min} \times V_t \times df / 12.0 \times 0.5 \times 1.0 \times V_s = \text{enzymatic activity (units per ml)}$ . Here,  $V_t$  is total volume;  $V_s$  is sample volume; 12.0 is the millimolar extinction coefficient of quinoneimine dye under the assay condition; 0.5 is the factor based on the fact that 1 mole of H<sub>2</sub>O<sub>2</sub> produces half a mole of quinoneimine dye;  $t$  is reaction time; 1.0 is light path (cm); and  $df$  is the dilution factor.

## RESULTS

#### *Acanthamoeba* exhibit PLA<sub>2</sub> activities

Both clinical and environmental isolates of *Acanthamoeba* exhibited PLA<sub>2</sub> activities (Fig. 1). As expected, optimal PLA<sub>2</sub> activities were observed with increasing numbers of *Acanthamoeba*. Interestingly, the environmental isolate belonging to T7 genotype exhibited the maximum PLA<sub>2</sub> activities, followed by the encephalitis isolate belonging to the T1 genotype, while the keratitis isolate belonging to the T4 genotype exhibited the least PLA<sub>2</sub> activities (Fig. 1). Of interest, *Acanthamoeba* condi-




**Fig. 1.** *Acanthamoeba* exhibiting PLA<sub>2</sub> activities. Both the encephalitis (A) and the environmental (C) isolates exhibited optimal, while the keratitis isolate (B) exhibited the least PLA<sub>2</sub> activities. Results are representative of 3 independent experiments performed in triplicate. Bars represent standard errors.

tioned medium did not exhibit PLA<sub>2</sub> activities in the assay conditions tested in the present study, suggesting that PLA<sub>2</sub> are cell-associated and/or not released (data not shown).

### PLD activities

The results revealed that both clinical and environmental isolates of *Acanthamoeba* exhibit PLD activities (Fig. 2). Again, the encephalitis isolate (T1) and environmental isolate (T7) exhibited the highest levels of PLD activities compared with the keratitis isolate (T4) (Fig. 2). It was interesting to note that



**Fig. 2.** Clinical and environmental *Acanthamoeba* isolates exhibiting PLD activities. As for PLA<sub>2</sub>, both the encephalitis (A) and the environmental (C) isolates exhibited optimal, while the keratitis isolate (B) exhibited the least PLD activities. Results are representative of 3 independent experiments performed in triplicate. Bars represent standard errors.

PLD activities in *Acanthamoeba* were significantly greater compared with *Acanthamoeba* PLA<sub>2</sub> activities (Figs. 1, 2). As for PLA<sub>2</sub>, the *Acanthamoeba* conditioned medium did not exhibit PLD activity in the assay conditions tested in the present study, indicating that PLD activities are cell-associated and/or not released (data not shown).

### Optimal phospholipase activities at 37°C



To determine the optimum phospholipase activities, assays were performed at various temperatures from 4°C to 65°C. The optimal phospholipase activities were observed at 37°C suggesting their physiological relevance (Fig. 3). Similar findings were observed both with PLA<sub>2</sub> and PLD activities. Activities were also observed at 65°C, suggesting that *Acanthamoeba* PLA<sub>2</sub> and PLD activities are stable at higher temperatures (Fig. 3).

### Optimal phospholipase activities at neutral pH


Next to determine the pH for optimal phospholipase activities, assays were performed at various pH. Our findings revealed that *Acanthamoeba* exhibit optimal PLA<sub>2</sub> and PLD activities at neutral pH indicating their physiological relevance (Fig. 4). Both clinical and environmental isolates exhibited optimal phospholipase activities at pH 7.5.

### Phospholipase inhibitors blocked the environmental isolate binding to HBMEC, but not clinical isolates


To determine whether phospholipases play a role in *Acan-*



**Fig. 3.** *Acanthamoeba* exhibiting optimal phospholipase activities at 37°C. To determine optimal temperatures of *Acanthamoeba* phospholipases, assays were performed at various temperatures using *Acanthamoeba* lysates ( $5 \times 10^6$  amoebae). Note that all *Acanthamoeba* isolates tested exhibit optimal PLA<sub>2</sub> (A) and PLD (B) activities at 37°C indicating their physiological relevance. Results are representative of 3 independent experiments performed in triplicate. Bars represent standard errors.



**Fig. 4.** *Acanthamoeba* exhibiting optimal phospholipase activities at neutral pH. To determine optimal pH of *Acanthamoeba* phospholipases, assays were performed at various pH using *Acanthamoeba* lysates ( $5 \times 10^6$  amoebae). Note that all *Acanthamoeba* isolates tested exhibit optimal PLA<sub>2</sub> (A) and PLD (B) activities at neutral pH indicating their physiological relevance. Results are representative of 3 independent experiments performed in triplicate. Bars represent standard errors.






**Fig. 5.** Phospholipase inhibitors blocking the environmental isolate-mediated binding to HBMEC but not the clinical isolates-mediated binding. The results revealed that both phospholipase inhibitors restricted environmental isolate (C)-mediated binding to HBMEC. However, neither inhibitors had any effects on the clinical isolates (A, T1; B, T4). Results are representative of 3 independent experiments performed in duplicates. Bars represent standard errors.

*thamoeba* binding to the host cells, adhesion assays were performed using the PLA<sub>2</sub> inhibitor, i.e., cytidine 5'-diphosphocholine, and the PLC/PLD inhibitor, i.e., compound 48/80. Our findings revealed that neither phospholipase inhibitor had any significant effects on the adhesion of clinical isolates of *Acanthamoeba* to HBMEC (Fig. 5). In contrast, adhesion of the envi-

ronmental isolate of *Acanthamoeba* was completely abolished in the presence of phospholipase inhibitors (Fig. 5).

Phospholipase A<sub>2</sub> inhibitor had no effect on *Acanthamoeba*-mediated HBMEC cytotoxicity

To determine the role of PLA<sub>2</sub> in *Acanthamoeba*-mediated



**Fig. 6.** Compound 48/80 partially inhibiting encephalitis isolate-mediated HBMEC cytotoxicity. The results revealed that cytidine 5'-diphosphocholine had no significant effect on *Acanthamoeba*-mediated HBMEC cytotoxicity. Similarly, compound 48/80 had no effect on *Acanthamoeba* keratitis isolate (B) but partially inhibited encephalitis isolate (A)-mediated HBMEC cytotoxicity. C is a environmental isolate of T7. Results are representative of 3 independent experiments performed in duplicates. Bars represent standard errors.

HBMEC death, cytotoxicity assays were performed in the presence of cytidine 5'-diphosphocholine, a PLA<sub>2</sub>-specific inhibitor. The results revealed that cytidine 5'-diphosphocholine had no significant effect on parasite-mediated HBMEC cytotoxicity in all *Acanthamoeba* isolates tested (Fig. 6).

Phospholipase C/D inhibitor inhibited HBMEC cytotoxicities mediated by the encephalitis isolate but not by the keratitis isolate

To determine whether PLC/PLD plays a role in *Acanthamoeba*-mediated HBMEC death, cytotoxicity assays were performed in the presence of compound 48/80. Interestingly, compound 48/80 partially blocked the encephalitis isolate-mediated HBMEC cytotoxicity, i.e., 49% cell death in the presence of the inhibitor compared with 73% in the absence of the inhibitor (Fig. 6). However, it had no effect on the keratitis isolate-mediated HBMEC cytotoxicity (Fig. 6). In contrast, the effects of phospholipase inhibitors on the environmental isolate-mediated HBMEC cytotoxicity remained unclear. This was due to the fact that the environmental isolate exhibited minimal HBMEC cytotoxicity even in the absence of inhibitors.

## DISCUSSION

Infections due to *Acanthamoeba* have increased over the years. This is most likely due to the increasing numbers of contact lens wearers, increasing populations of immunocompromised patients, and global warming. This has led to a remarkable interest in the field of *Acanthamoeba* pathogenesis and to design strategies to control and prevent *Acanthamoeba* infections. In particular, over the last 2 decades, research in *Acanthamoeba* virulence factors has gained significant attention to identify novel approaches for therapeutic interventions. *Acanthamoeba* pathogenesis is attributed to their ability to bind to the host cells, induce host cell apoptosis, secrete proteases, exhibit ecto-ATPase activities, and switch phenotypes. The present study was undertaken to determine the roles of phospholipases (i.e., PLA<sub>2</sub> and PLD) in *Acanthamoeba* and *Acanthamoeba*-mediated binding to, and cytotoxicity of HBMEC in vitro. Our findings revealed that *Acanthamoeba* show PLA<sub>2</sub> and PLD activities. This is consistent with previous studies that showed the presence of phospholipase activities in *Acanthamoeba* [5-7].

It was interesting to note that the environmental isolate (belonging to the T7 genotype) exhibited higher phospholipase

activities as compared to the keratitis isolate (belonging to the T4 genotype). In contrast, the environmental isolate exhibited minimal binding to or cytotoxicity of HBMEC. This is not a surprising finding. Being a free-living amoeba, the primary functions of phospholipases in *Acanthamoeba* may be predominantly in the phospholipids turnover for membrane maintenance and remodelling. The fact that environmental isolate is significantly larger in size as compared to the keratitis isolate, thus larger plasma membranes, may explain its optimal PLA<sub>2</sub> and PLD activities, and thus rationalize these findings. In support, the size of keratitis isolate tested here is approximately 7 µm. The size of the encephalitis isolate is approximately 10 µm and the environmental isolate being the largest, approximately 17 µm. Another explanation may be that the environmental isolate exhibited distinct types or isoforms of phospholipases compared with the clinical isolates. For example, Sissons et al. [17] have shown that both clinical and environmental isolates of *Acanthamoeba* exhibit ecto-ATPase activities, however, clinical isolates exhibit distinct ecto-ATPases. This may be the case for phospholipases, which are normally required for routine cellular functions but pathogenic isolates may possess distinct phospholipases. The fact that compound 48/80 had no effect on keratitis isolate-mediated host cell death but partially inhibited encephalitis isolate-mediated HBMEC cytotoxicity further suggests possible differences in phospholipases in *Acanthamoeba* belonging to different genotypes. Future studies are in progress to determine the molecular basis of these differences.

Of interest was the observation that compound 48/80, only partially inhibited encephalitis isolate-mediated HBMEC cytotoxicity. It is noteworthy that *Acanthamoeba* pathogenesis is a multifactorial process involving adherence, phagocytosis, apoptosis, proteases, ecto-ATPases, and the inhibition of a single putative factor may not be sufficient to block their ability to produce host cell cytotoxicity. In support, previous studies have shown that inhibition of *Acanthamoeba* binding to host cells is not adequate to block host cell cytotoxicity [18]. At present, the inability of compound 48/80 to block keratitis isolate-mediated host cell death is unclear. To this end, the source of amoeba isolates, i.e., keratitis and encephalitis patients may be indicative of these differences. This possibly means that phospholipases from the encephalitis isolate may play a role in the pathogenesis of AGE. However, it is a subject which needs additional research and study.

The finding that phospholipase inhibitors did not clearly

block *Acanthamoeba*-mediated HBMEC cytotoxicity does not rule out their involvement in *Acanthamoeba* pathogenesis. It is possible that cytotoxicity is a delayed event and that phospholipases are involved in the early events. Recent studies have shown that phospholipases may be involved in interference with the intracellular signalling pathways. For example, phospholipases generate lipids and lipid-derived products that act as mediators and second messengers, which may act as the modulators of signal transduction pathway [19,20]. Oishi et al. [21] showed that lysophospholipids, by-end products of PLA<sub>2</sub> and PLB, induce activation of protein kinase C, which has diverse functions in host cell signalling pathways [21]. Phospholipase C of *Clostridium perfringens* induces expression of IL-8 synthesis in endothelial cells [22,23]. Phospholipase A facilitate host cell penetration of *Toxoplasma gondii* [24] and *Entamoeba histolytica* [25], while PLD has been shown to be an important virulence determinant of *Corynebacterium pseudotuberculosis* in the persistence and spread of bacteria within the host [26].

Overall, these studies suggest that *Acanthamoeba* phospholipases may play a role in causing host cell damage or affect other cellular functions, such as inducing inflammatory responses and thus facilitating the virulence of *Acanthamoeba*. The fact that phospholipases cleave phospholipids suggests their possible role in membrane disruptions and penetration into host cells. Future studies are needed to identify and characterize *Acanthamoeba* phospholipases, which should help determine their potential role for therapeutic interventions and in differentiation of *Acanthamoeba* isolates belonging to different genotypes. Some previous studies have shown that phospholipase C from *C. perfringens* induced protection against *C. perfringens*-mediated gas gangrene [27]. In addition, targeting phospholipases using synthetic compounds are shown to have the potential to prevent *Candida* infections [28].

## ACKNOWLEDGEMENTS

Funding for this paper was provided by Namseoul University.

## REFERENCES

1. Khan NA. *Acanthamoeba*: biology and increasing importance in human health. FEMS Microbiol Rev 2006; 30: 564-595.
2. Marciano-Cabral F, Cabral G. *Acanthamoeba* spp. as agents of disease in humans. Clin Microbiol Rev 2003; 16: 273-307.
3. Schuster FL, Visvesvara GS. Free-living amoebae as opportunistic and non-opportunistic pathogens of humans and animals. Int J

- Parasitol* 2004; 34: 1001-1027.
4. Jung SY, Alsam S, Kim KS, Khan NA. Pathogen-pathogen interactions: a comparative study of *Escherichia coli* interactions with the clinical and environmental isolates of *Acanthamoeba*. *World J Microbiol Biotechnol* 2008; 24: 2339-2348.
  5. Victoria EJ, Korn ED. Plasma membrane and soluble lysophospholipases of *Acanthamoeba castellanii*. *Arch Biochem Biophys* 1975; 171: 255-258.
  6. Victoria EJ, Korn ED. Enzymes of phospholipid metabolism in the plasma membrane of *Acanthamoeba castellanii*. *J Lipid Res* 1975; 16: 54-60.
  7. Cursons RIM, Brown TJ, Keys EA. Virulence of pathogenic free-living amoebae. *J Parasitol* 1978; 64: 744-745.
  8. Ghannoum MA. Potential role of phospholipases in virulence and fungal pathogenesis. *Clin Microbiol Rev* 2000; 13: 122-143.
  9. Exton JH. Phospholipase D-structure, regulation and function. *Rev Physiol Biochem Pharmacol* 2002; 144: 1-94.
  10. McDermott M, Wakelam MJO, Morris AJ. Phospholipase D. *Biochem Cell Biol* 2004; 82: 225-253.
  11. Akiba S, Sato T. Cellular function of calcium-independent phospholipase A<sub>2</sub>. *Biol Pharm Bull* 2004; 27: 1174-1178.
  12. Kudo I. Diversity of phospholipase A<sub>2</sub> enzymes. Foreword. *Biol Pharm Bull* 2004; 27: 1157.
  13. Sissons J, Kim KS, Stins M, Jayasekera S, Alsam S, Khan NA. *Acanthamoeba castellanii* induces host cell death via a phosphatidylinositol 3-kinase-dependent mechanism. *Infect Immun* 2005; 73: 2704-2708.
  14. Stins ME, Gilles F, Kim KS. Selective expression of adhesion molecules on human brain microvascular endothelial cells. *J Neuroimmunol* 1997; 76: 81-90.
  15. Alsam S, Kim KS, Stins M, Rivas AO, Sissons J, Khan NA. *Acanthamoeba* interactions with human brain microvascular endothelial cells. *Microb Pathog* 2003; 35: 235-241.
  16. Matin A, Stins M, Kim KS, Khan NA. *Balamuthia mandrillaris* exhibits metalloprotease activities. *FEMS Immunol Med Microbiol* 2006; 47: 83-91.
  17. Sissons J, Alsam S, Jayasekera S, Khan NA. Ecto-ATPases of clinical and non-clinical isolates of *Acanthamoeba*. *Microb Pathog* 2004; 37: 231-239.
  18. Leher H, Silvano R, Alizadeh H, Huang J, Niederkorn JY. Mannose induces the release of cytopathic factors from *Acanthamoeba castellanii*. *Infect Immun* 1998; 66: 5-10.
  19. Dennis EA, Rhee SG, Billah MM, Hannun YA. Role of phospholipase in generating lipid second messengers in signal transduction. *FASEB J* 1991; 5: 2068-2077.
  20. Serhan CN, Haeggström JZ, Leslie CC. Lipid mediator networks in cell signaling: update and impact of cytokines. *FASEB J* 1996; 10: 1147-1158.
  21. Oishi K, Raynor RL, Charp PA, Kuo JF. Regulation of protein kinase C by lysophospholipids. Potential role in signal transduction. *J Biol Chem* 1988; 263: 6865-6871.
  22. Bryant AE, Stevens DL. Phospholipase C and perfringolysin O from *Clostridium perfringens* upregulate endothelial cell-leukocyte adherence molecule 1 and intercellular leukocyte adherence molecule 1 expression and induce interleukin-8 synthesis in cultured human umbilical vein endothelial cells. *Infect Immun* 1996; 64: 358-362.
  23. Bunting M, Lorant DE, Bryant AE, Zimmerman GA, McIntyre TM, Stevens DL, Prescott SM. Alpha toxin from *Clostridium perfringens* induces proinflammatory changes in endothelial cells. *J Clin Invest* 1997; 100: 565-574.
  24. Saffer LD, Long Krug SA, Schwartzman JD. The role of phospholipase in host cell penetration by *Toxoplasma gondii*. *Am J Trop Med Hyg* 1989; 40: 145-149.
  25. Long-Krug SA, Fischer KJ, Hysmith RM, Ravdin JI. Phospholipase A enzymes of *Entamoeba histolytica*: description and subcellular localization. *J Infect Dis* 1985; 152: 536-541.
  26. McNamara PJ, Bradley GA, Songer JG. Targeted mutagenesis of the phospholipase D gene results in decreased virulence of *Corynebacterium pseudotuberculosis*. *Mol Microbiol* 1994; 12: 921-930.
  27. Kameyama S, Sato H, Murata R. The role of alpha-toxin of *Clostridium perfringens* in experimental gas gangrene in guinea pigs. *Jpn J Med Sci Biol* 1972; 25: 200.
  28. Hänel H, Kirsch R, Schmidts HL, Kottmann H. New systematically active antimycotics from the beta-blocker category. *Mycoses* 1995; 38: 251-264.