

## Integral Operator of Analytic Functions with Positive Real Part

BASEM AREF FRASIN

*Faculty of Science, Department of Mathematics, Al al-Bayt University, P. O. Box: 130095 Mafraq, Jordan*

*e-mail: bafrasin@yahoo.com*

ABSTRACT. In this paper, we introduce the integral operator  $I_\beta(p_1, \dots, p_n; \alpha_1, \dots, \alpha_n)(z)$  of analytic functions with positive real part. The radius of convexity of this integral operator when  $\beta = 1$  is determined. In particular, we get the radius of starlikeness and convexity of the analytic functions with  $Re \{f(z)/z\} > 0$  and  $Re \{f'(z)\} > 0$ . Furthermore, we derive sufficient condition for the integral operator  $I_\beta(p_1, \dots, p_n; \alpha_1, \dots, \alpha_n)(z)$  to be analytic and univalent in the open unit disc, which leads to univalence of the operators  $\int_0^z (f(t)/t)^\alpha dt$  and  $\int_0^z (f'(t))^\alpha dt$ .

### 1. Introduction and definitions

Let  $\mathcal{A}$  denote the class of functions of the form :

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

which are analytic in the open unit disc  $\mathcal{U} = \{z : |z| < 1\}$ . Further, by  $\mathcal{S}$  we shall denote the class of all functions in  $\mathcal{A}$  which are univalent in  $\mathcal{U}$ . A function  $f(z)$  belonging to  $\mathcal{S}$  is said to be starlike if it satisfies

$$(1.1) \quad \operatorname{Re} \left( \frac{z f'(z)}{f(z)} \right) > 0 \quad (z \in \mathcal{U})$$

We denote by  $\mathcal{S}^*$  the subclass of  $\mathcal{A}$  consisting of functions which are starlike in  $\mathcal{U}$ . Also, a function  $f(z)$  belonging to  $\mathcal{S}$  is said to be convex if it satisfies

$$(1.2) \quad \operatorname{Re} \left( 1 + \frac{z f''(z)}{f'(z)} \right) > 0 \quad (z \in \mathcal{U})$$

---

Received August 3, 2010; accepted September 27, 2010.

2000 Mathematics Subject Classification: 30C45.

Key words and phrases: Analytic and univalent functions, Starlike and convex functions, Functions of positive real part, Integral operator.

We denote by  $\mathcal{K}$  the subclass of  $\mathcal{A}$  consisting of functions which are convex in  $\mathcal{U}$ .

Let  $\alpha_i \in \mathbb{C}$  for all  $i = 1, \dots, n$ ,  $n \in \mathbb{N}$ ,  $\beta \in \mathbb{C}$  with  $\operatorname{Re}(\beta) > 0$ . We let  $I_\beta : A^n \rightarrow A$  be the integral operator defined by

$$(1.3) \quad I_\beta(p_1, \dots, p_n; \alpha_1, \dots, \alpha_n)(z) = \left\{ \int_0^z \beta t^{\beta-1} (p_1(t))^{\alpha_1} \dots (p_n(t))^{\alpha_n} dt \right\}^{\frac{1}{\beta}},$$

where  $p_i(z)$  are analytic in  $\mathcal{U}$  and satisfy  $p_i(0) = 1$  for all  $i = 1, \dots, n$ . Here and throughout in the sequel every many-valued function is taken with the principal branch.

**Remark 1.1.** Note that the integral operator  $I_\beta(p_1, \dots, p_n; \alpha_1, \dots, \alpha_n)(z)$  generalizes many operators introduced and studied by several authors, for example:

(2) For  $p_i(t) = \frac{D_\lambda^{m,\gamma} f_i(t)}{t}$ ;  $1 \leq i \leq n$ , we obtain the following integral operator introduced and studied by Bulut [7]

$$(1.4) \quad I_\beta^{m,\gamma}(f_1, \dots, f_n)(z) = \left\{ \int_0^z \beta t^{\beta-1} \left( \frac{D_\lambda^{m,\gamma} f_1(t)}{t} \right)^{\alpha_1} \dots \left( \frac{D_\lambda^{m,\gamma} f_n(t)}{t} \right)^{\alpha_n} dt \right\}^{\frac{1}{\beta}}$$

where  $D_\lambda^{m,\gamma} f(z) = z + \sum_{n=2}^{\infty} \left( \frac{\Gamma(n+1)\Gamma(2-\gamma)}{\Gamma(n+1-\gamma)} (1 + (n-1)\lambda) \right)^m a_n z^n$ ,  $m \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$  is the generalized Al-Oboudi operator [2].

(3) For  $p_i(t) = \frac{f_i(t)}{t}$ ;  $1 \leq i \leq n$ , we obtain the integral operator

$$(1.5) \quad I_\beta(f_1, \dots, f_n)(z) = \left\{ \int_0^z \beta t^{\beta-1} \left( \frac{f_1(t)}{t} \right)^{\alpha_1} \dots \left( \frac{f_n(t)}{t} \right)^{\alpha_n} dt \right\}^{\frac{1}{\beta}}$$

introduced and studied by Breaz and Breaz [3].

(1) For  $\beta = 1$  and  $p_i(t) = \frac{(f_i * g_i)(t)}{t}$ ;  $1 \leq i \leq n$ , we obtain the integral operator

$$(1.6) \quad I(f_1, \dots, f_n; g_1, \dots, g_n)(z) = \int_0^z \left( \frac{(f_1 * g_1)(t)}{t} \right)^{\alpha_1} \dots \left( \frac{(f_n * g_n)(t)}{t} \right)^{\alpha_n} dt$$

introduced and studied by Frasin [9].

(4) For  $\beta = 1$  and  $p_i(t) = \frac{f_i(t)}{t}$ ;  $1 \leq i \leq n$ , we obtain the integral operator

$$(1.7) \quad F_n(z) = \int_0^z \left( \frac{f_1(t)}{t} \right)^{\alpha_1} \dots \left( \frac{f_n(t)}{t} \right)^{\alpha_n} dt$$

introduced and studied by Breaz and Breaz [3].

(5) For  $\beta = 1$  and  $p_i(t) = f'_i(t)$ ;  $1 \leq i \leq n$ , we obtain the integral operator

$$(1.8) \quad F_{\alpha_1, \dots, \alpha_n}(z) = \int_0^z (f'_1(t))^{\alpha_1} \dots (f'_n(t))^{\alpha_n} dt$$

introduced and studied by Breaz *et al.* [5].

(6) For  $\beta = 1$  and  $p_i(t) = \frac{R^k f_i(t)}{t}$ ;  $1 \leq i \leq n$ , we obtain the integral operator introduced in [11]

$$(1.9) \quad I(f_1, \dots, f_n)(z) = \int_0^z \left( \frac{R^k f_1(t)}{t} \right)^{\alpha_1} \dots \left( \frac{R^k f_n(t)}{t} \right)^{\alpha_n} dt,$$

where  $R^k f(z) = z + \sum_{n=2}^{\infty} C_{k+n-1}^k a_n z^n$ ,  $k \in \mathbb{N}_0$  is Ruscheweyh differential operator [17].

(7) For  $\beta = 1$  and  $p_i(t) = \frac{D^k f_i(t)}{t}$ ;  $1 \leq i \leq n$ , we obtain the integral operator introduced and studied by Breaz *et al.* [4]

$$(1.10) \quad D^k F(z) = \int_0^z \left( \frac{D^k f_1(t)}{t} \right)^{\alpha_1} \dots \left( \frac{D^k f_n(t)}{t} \right)^{\alpha_n} dt,$$

where  $D^k f(z) = z + \sum_{n=2}^{\infty} n^k a_n z^n$ ,  $k \in \mathbb{N}_0$  is Sălăgean differential operator [18].

(8) For  $\beta = 1$  and  $p_i(t) = \frac{D_{\lambda}^k f_i(t)}{t}$ ;  $1 \leq i \leq n$ , we obtain the following integral operator introduced and studied by Bulut [6]

$$(1.11) \quad I_n(f_1, \dots, f_n)(z) = \int_0^z \left( \frac{D_{\lambda}^k f_1(t)}{t} \right)^{\alpha_1} \dots \left( \frac{D_{\lambda}^k f_n(t)}{t} \right)^{\alpha_n} dt,$$

where  $D_{\lambda}^k f(z) = z + \sum_{n=2}^{\infty} [1 + (n-1)\lambda]^k a_n z^n$ ,  $0 \leq \lambda \leq 1$ , is Al-Oboudi differential operator [2].

(9) For  $\beta = 1$  and  $p_i(t) = \frac{L(a,c)f_i(t)}{t}$ ;  $1 \leq i \leq n$ , we obtain the integral operator introduced and studied by Selvaraj and Karthikeyan [19]

$$(1.12) \quad F_{\alpha}(a, c; z) = \int_0^z \left( \frac{L(a, c)f_1(t)}{t} \right)^{\alpha_1} \dots \left( \frac{L(a, c)f_n(t)}{t} \right)^{\alpha_n} dt,$$

where  $L(a, c)f(z) := z + \sum_{n=2}^{\infty} \frac{(a)_{n-1}}{(c)_{n-1}} a_n z^n$  is the Carlson-Shaffer linear operator [8].

(5) For  $\beta = 1$ ,  $n = 1$ ,  $\alpha_1 = \alpha$  and  $p_1(t) = \frac{f(t)}{t}$ , we obtain the integral operator

$$(1.13) \quad F_\alpha(z) = \int_0^z \left( \frac{f(t)}{t} \right)^\alpha dt$$

studied in [13]. In particular, for  $\alpha = 1$ , we obtain Alexander integral operator introduced in [1]

$$(1.14) \quad I(z) = \int_0^z \frac{f(t)}{t} dt$$

(6) For  $\beta = 1$ ,  $n = 1$ ,  $\alpha_1 = \alpha$  and  $p_1(t) = f'(t)$ , we obtain the integral operator

$$(1.15) \quad G_\alpha(z) = \int_0^z (f'(t))^\alpha dt$$

studied in [15] (see also [16]).

In the present paper, the radius of convexity of the integral operator defined by (1.3) when  $\beta = 1$  are determined. In particular, we get the radius of starlikeness and convexity of the analytic functions with  $Re\{f(z)/z\} > 0$  and  $Re\{f'(z)\} > 0$  obtained by MacGregor [12]. Furthermore, we derive sufficient condition for the integral operator  $I_\beta(p_1, \dots, p_n; \alpha_1, \dots, \alpha_n)(z)$  to be analytic and univalent in  $\mathcal{U}$ , which leads to univalence of the operators  $\int_0^z (f(t)/t)^\alpha dt$  and  $\int_0^z (f'(t))^\alpha dt$  obtained by Kim and Merkes [10] and Pfaltzgraff [16].

In the proofs of our main results we need the following lemmas

**Lemma 1.2([12]).** *Let  $p(z) = 1 + c_1z + c_2z^2 + \dots$  be analytic in  $\mathcal{U}$  and satisfy  $p(0) = 1$  with  $Re\{p(z)\} > 0$ , then we have*

$$(1.16) \quad \left| \frac{zp'(z)}{p(z)} \right| < \frac{2|z|}{1-|z|^2}, \quad (z \in \mathcal{U}).$$

**Lemma 1.3([14]).** *Let  $\beta \in \mathbb{C}$  with  $Re(\beta) > 0$ . If  $f \in \mathcal{A}$  satisfies*

$$(1 - |z|^{2Re(\beta)}) \left| \frac{zf''(z)}{f'(z)} \right| \leq Re(\beta),$$

for all  $z \in \mathcal{U}$ , then the integral operator

$$F_\beta(z) = \left\{ \beta \int_0^z t^{\beta-1} f'(t) dt \right\}^{\frac{1}{\beta}}$$

is in the class  $\mathcal{S}$ .

## 2. Convexity of $I_1(p_1, \dots, p_n; \alpha_1, \dots, \alpha_n)(z)$

In this section, we obtain the radius of convexity of the integral operator  $I_\beta(p_1, \dots, p_n; \alpha_1, \dots, \alpha_n)(z)$  defined by (1.3) when  $\beta = 1$ .

**Theorem 2.1.** *Let  $p_i(0) = 1$  and  $\operatorname{Re}\{p_i(z)\} > 0$  for all  $i = 1, \dots, n$ . Then the integral operator defined by*

$$(2.1) \quad I_1(z) = \int_0^z \prod_{i=1}^n (p_i(t))^{\alpha_i} dt$$

is convex in  $|z| = r < \sqrt{\left(\sum_{i=1}^n \alpha_i\right)^2 + 1} - \sum_{i=1}^n \alpha_i$ ; where  $\alpha_i > 0$  for all  $i = 1, \dots, n$ .

*Proof.* From (2.1), it is easy to see that  $I_1(0) = I_1'(0) - 1 = 0$  and

$$(2.2) \quad I_1'(z) = \prod_{i=1}^n (p_i(z))^{\alpha_i}.$$

Differentiate (2.2) logarithmically with respect with  $z$ , we obtain

$$\frac{zI_1''(z)}{I_1'(z)} = \sum_{i=1}^n \alpha_i \left( \frac{zp_i'(z)}{p_i(z)} \right)$$

or, equivalently,

$$(2.3) \quad 1 + \frac{zI_1''(z)}{I_1'(z)} = 1 + \sum_{i=1}^n \alpha_i \left( \frac{zp_i'(z)}{p_i(z)} \right).$$

Now, using the estimate (1.16) in Lemma 1.2, from (2.3) it follows that

$$\begin{aligned} \operatorname{Re} \left\{ 1 + \frac{zI_1''(z)}{I_1'(z)} \right\} &= 1 + \operatorname{Re} \sum_{i=1}^n \alpha_i \left( \frac{zp_i'(z)}{p_i(z)} \right) \\ &\geq 1 - \left| \sum_{i=1}^n \alpha_i \left( \frac{zp_i'(z)}{p_i(z)} \right) \right| \\ &\geq 1 - \left( \frac{2|z|}{1-|z|^2} \right) \sum_{i=1}^n \alpha_i, \quad (|z| < 1). \end{aligned}$$

If  $|z| < \sqrt{\left(\sum_{i=1}^n \alpha_i\right)^2 + 1} - \sum_{i=1}^n \alpha_i$ , then  $\operatorname{Re} \left\{ 1 + \frac{zI_1''(z)}{I_1'(z)} \right\} > 0$ . Therefore,  $I_1(z)$  is convex in  $|z| < \sqrt{\left(\sum_{i=1}^n \alpha_i\right)^2 + 1} - \sum_{i=1}^n \alpha_i$ . This proves the theorem.  $\square$

Letting  $n = 1$ ,  $\alpha_1 = 1$  and  $p_1 = p$  in Theorem 2.1, we have

**Corollary 2.2.** *Let  $p(0) = 1$  and  $\operatorname{Re}\{p(z)\} > 0$ . Then the integral operator  $\int_0^z p(t)dt$  is convex in  $|z| < \sqrt{2} - 1$ .*

Letting  $p(z) = f(z)/z$  and  $p(z) = f'(z)$  in Corollary 2.2, we get the following interesting results due to MacGregor [12].

**Corollary 2.3.** *Let  $f(z) \in \mathcal{A}$ . If  $\operatorname{Re}\{f(z)/z\} > 0$  for  $|z| < 1$ , then  $f(z)$  is starlike in  $|z| < \sqrt{2} - 1$ . The result is sharp for the extremal function  $f(z) = (z + z^2)/(1 - z^2)$ .*

**Corollary 2.4.** *Let  $f(z) \in \mathcal{A}$ . If  $\operatorname{Re}\{f'(z)\} > 0$  for  $|z| < 1$ , then  $f(z)$  is convex in  $|z| < \sqrt{2} - 1$ . The result is sharp for the extremal function  $f(z) = \int_0^z (1+t)/(1-t)dt$ .*

**Remark 2.5.** Taking different choices of  $p_i(z)$  as stated in Section 1, Theorems 2.1 leads to new radius of convexity for the integral operators defined in Section 1 when  $\beta = 1$ .

### 3. Univalence of $I_\beta(p_1, \dots, p_n; \alpha_1, \dots, \alpha_n)(z)$

Next, we obtain the following sufficient condition for the integral operator  $I_\beta(p_1, \dots, p_n; \alpha_1, \dots, \alpha_n)(z)$  to be analytic and univalent in  $\mathcal{U}$ .

**Theorem 3.1.** *Let  $\alpha_i \in \mathbb{C}$  for all  $i = 1, \dots, n$  and  $\beta \in \mathbb{C}$  with  $\operatorname{Re}(\beta) = a$ . If*

$$(3.1) \quad \sum_{i=1}^n |\alpha_i| \leq \begin{cases} a/2 & \text{if } 0 < a \leq 1/2 \\ 1/4 & \text{if } a \geq 1/2. \end{cases}$$

*Then the integral operator  $I_\beta(p_1, \dots, p_n; \alpha_1, \dots, \alpha_n)(z)$  defined by (1.3) is analytic and univalent in  $\mathcal{U}$ , where  $p_i(z)$  are analytic in  $\mathcal{U}$  and satisfy  $p_i(0) = 1$  with  $\operatorname{Re}\{p_i(z)\} > 0$  for all  $i = 1, \dots, n$ .*

*Proof.* Define

$$h(z) = \int_0^z \prod_{i=1}^n (p_i(t))^{\alpha_i} dt,$$

so that, obviously,

$$(3.2) \quad h'(z) = \prod_{i=1}^n (p_i(z))^{\alpha_i}.$$

Making use of the logarithmic differentiation on both sides of (3.2) and multiplying

by  $z$ , we have

$$(3.3) \quad \frac{zh''(z)}{h'(z)} = \sum_{i=1}^n \alpha_i \left( \frac{zp'_i(z)}{p_i(z)} \right).$$

From (1.16) and (3.3), we obtain

$$\begin{aligned} (1 - |z|^{2a}) \left| \frac{zh''(z)}{h'(z)} \right| &\leq (1 - |z|^{2a}) \sum_{i=1}^n |\alpha_i| \left| \frac{zp'_i(z)}{p_i(z)} \right| \\ &\leq (1 - |z|^{2a}) \left( \frac{2|z|}{1 - |z|^2} \right) \sum_{i=1}^n |\alpha_i|. \end{aligned}$$

Since  $\frac{2|z|}{1 - |z|^2} \leq \frac{2}{1 - |z|}$  for  $z \in \mathcal{U}$ , we have

$$(1 - |z|^{2a}) \left| \frac{zh''(z)}{h'(z)} \right| \leq \left( \frac{2(1 - |z|^{2a})}{1 - |z|} \right) \sum_{i=1}^n |\alpha_i|.$$

Define the function  $\Psi : (0, 1) \rightarrow \mathbb{R}$  by

$$\Psi(x) = \frac{2(1 - x^{2a})}{1 - x}, \quad (a > 0, x = |z|).$$

It is easy to show that

$$(3.4) \quad \Psi(x) \leq \begin{cases} 2 & \text{if } 0 < a \leq 1/2 \\ 4a & \text{if } a \geq 1/2. \end{cases}$$

We thus find from (3.4) and the hypothesis (3.1) that

$$(1 - |z|^{2a}) \left| \frac{zh''(z)}{h'(z)} \right| \leq a, \quad (z \in \mathcal{U}).$$

Applying Lemma 1.3 for the function  $h(z)$  with  $\operatorname{Re}(\beta) = a$ , we prove that  $I_\beta(p_1, \dots, p_n; \alpha_1, \dots, \alpha_n)(z) \in \mathcal{S}$ . This evidently completes the proof of Theorem 3.1.  $\square$

Letting  $n = 1$ ,  $\alpha_1 = \alpha$ ,  $p_1 = p$  in Theorem 3.1, we have

**Corollary 3.2.** *Let  $\alpha \in \mathbb{C}$  and  $\beta \in \mathbb{C}$  with  $\operatorname{Re}(\beta) = a$ . If*

$$(3.5) \quad |\alpha| \leq \begin{cases} a/2 & \text{if } 0 < a \leq 1/2 \\ 1/4 & \text{if } a \geq 1/2. \end{cases}$$

*Then the integral operator  $I_\beta(p; \alpha)(z) = \left\{ \int_0^z \beta t^{\beta-1} (p(t))^\alpha dt \right\}^{\frac{1}{\beta}}$  is analytic and univalent in  $\mathcal{U}$ , where  $p(z)$  is analytic in  $\mathcal{U}$  and satisfy  $p(0) = 1$ ,  $\operatorname{Re}\{p(z)\} > 0$ .*

Letting  $\beta = 1$  in Corollary 3.2, we have

**Corollary 3.3.** *Let  $p(z)$  be analytic in  $\mathcal{U}$  and satisfy  $p(0) = 1$ ,  $\operatorname{Re}\{p(z)\} > 0$ . Then the integral operator  $\int_0^z (p(t))^\alpha dt$  is analytic and univalent in  $\mathcal{U}$ , where  $|\alpha| \leq 1/4$ ;  $\alpha \in \mathbb{C}$ .*

Letting  $p(z) = f(z)/z$  in Corollary 3.3, we get the following result obtained by Kim and Merkes [10].

**Corollary 3.4.** *Let  $f \in \mathcal{S}$ , and  $\alpha \in \mathbb{C}$ ;  $|\alpha| \leq 1/4$ . Then the integral operator  $\int_0^z (f(t)/t)^\alpha dt$  is analytic and univalent in  $\mathcal{U}$ .*

Letting  $p(z) = f'(z)$  in Corollary 3.3, we get the following result obtained by Pfaltzgraß [16].

**Corollary 3.5.** *Let  $f \in \mathcal{S}$ , and  $\alpha \in \mathbb{C}$ ;  $|\alpha| \leq 1/4$ . Then the integral operator  $\int_0^z (f'(t))^\alpha dt$  is analytic and univalent in  $\mathcal{U}$ .*

**Remark 3.6.** Taking different choices of  $p_i(z)$ , where  $p_i(z)$  as stated in Section 1, Theorem 3.1 leads to new sufficient conditions for the integral operators defined in Section 1 to be analytic and univalent in  $\mathcal{U}$ .

## References

- [1] W. Alexander, *Functions which map the interior of the unit circle upon simple regions*, Anna. Math., **17**(1915), 12-22.
- [2] F. M. Al-Oboudi, *On univalent functions defined by a generalized Sălăgean operator*, Int. J. Math. Math. Sci., **27**(2004), 1429-1436.
- [3] D. Breaz and N. Breaz, *Two integral operator*, Studia Universitatis Babes-Bolyai, Mathematica, Cluj-Napoca, **3**(2002), 13-21.
- [4] D. Breaz, H. Güney and G. Sălăgean, *A new integral operator*, Tamsui Oxford Journal of Mathematical Sciences, **25**(4)(2009), 407-414.
- [5] D. Breaz, S. Owa and N. Breaz, *A new integral univalent operator*, Acta Univ. Apul., **16**(2008), 11-16.
- [6] S. Bulut, *Some properties for an integral operator defined by Al-Oboudi differential operator*, JIPAM, **9**(2008), Issue 4, Atr. 115, 5 pp.
- [7] S. Bulut, *Univalence preserving integral operators defined by generalized Al-Oboudi differential operators*, An. St. Univ. Ovidius Constata, **17**(2009), 37-50.
- [8] B. C. Carlson, D. B. Shaffer, *Starlike and prestarlike hypergeometric functions*, SIAM J. Math. Anal., **15**(4)(1984), 737-745.



- [9] B. A. Frasin, *General integral operator defined by Hadamard product*, Math Vesnik, **62**, **2**(2010), 127-136.
- [10] I. J. Kim and E. P. Merkes, *On an integral of powers of a spirallike function*, Kyungpook Math. J., **12**(2)(1972), 249-253.
- [11] G. I. Oros, G. Oros and D. Breaz, *Sufficient conditions for univalence of an integral operator*, J. Ineq. Appl., Vol. 2008, Article ID 127645, 7 pages.
- [12] T. MacGregor, *The radius of univalence of certain analytic functions*, Proc. Amer. Math. Soc., **14**(1963), 514-520.
- [13] S. S. Miller, P. T. Mocanu, and M. O. Reade, *Starlike integral operators*, Pacific J. Math., **79**(1978), 157-168.
- [14] N. Pascu, *An improvement of Backer's univalence criterion*, Proceedings of the Commemorative Session Simion Stoilow, Brasov, (1987), 43-48.
- [15] N. Pascu and V. Pescar, *On the integral operators of Kim-Merkis and Pfaltzgraff*, Mathematica, **32**(55)(1990), 185-192.
- [16] J. A. Pfaltzgraff, *Univalence of the integral of  $f'(z)^\lambda$* , Bull. London Math. Soc., **7**(3)(1975), 254-256.
- [17] St. Ruscheweyh, *New criteria for univalent functions*, Proc. Amer. Math. Soc., **49**(1975), 109-115.
- [18] G. Sălăgean, *Subclasses of univalent functions*, Lecture Notes in Math., (Springer-Verlag), **1013**(1983), 362-372.
- [19] C. Selvaraj and K. R. Karthikeyan, *Sufficient conditions for univalence of a general integral operator*, Bull. Korean Math. Soc., **46**(2)(2009), 367-372.