

# Pre-operative Evaluation of Consistency in Intra-axial Brain Tumor with Diffusion-weighted Images (DWI) and Conventional MR Images

Moon Sik Oh, Kook Jin Ahn, Hyun Seok Choi, So Lyung Jung, Yoon Joo Lee, Bum Soo Kim

**Purpose :** To retrospectively evaluate the usefulness of diffusion-weighted images, ADC maps and conventional MR images for determination of brain tumor consistency.

**Materials and Methods :** Twenty-three patients with brain tumor underwent MR examinations with T1, T2 and diffusion-weighted images. Regions of interest (ROIs) were drawn in the tumors, and the measured signal intensities (SI) were normalized with the contralateral side. We evaluated the correlation between SI ratios from various images and tumor consistency assessed at surgery. In three patients with both cystic and solid components, each component was evaluated independently. Qualitatively observed SIs were also correlated with tumor consistency.

**Results :** Statistical analysis revealed significant correlation between tumor consistency and ADC ratio ( $r = -0.586$ ,  $p = 0.002$ ), SI ratios on T2-weighted images ( $r = -0.497$ ,  $p = 0.010$ ), and observed SIs on T2-weighted images ( $r = -0.461$ ,  $p = 0.018$ ). The relative ratio of ADC value correlated with tumor consistency most strongly.

**Conclusion :** The measured ratio of ADC, SI ratio and observed SI grade on T2-weighted images can provide valuable information about the consistency of brain tumor.

**Index words :** Consistency  
Brain tumor  
Diffusion weighted image

## Introduction

The surgical management of brain tumor is the first step of treatment. Preoperative planning for excision of intra-axial brain tumor requires knowledge about

tumor consistency as well as location, size and invasiveness. Operating time is also affected by tumor consistency; whereas soft tumors can easily removed with suction, careful and diligent dissection is required for harder tumors. Thus information of the consistency of intra-axial brain tumor may help the clinician plan

---

### JKSMRM 15:102-109(2011)

Department of Radiology, Collage of Medicine, The Catholic University of Korea

Received; July 27, 2011, revised; August 29, 2011, accepted; August 30, 2011

Corresponding author : Koon Jin Ahn, M.D., Department of Radiology, Seoul St. Mary' Hospital, College of Medicine, The Catholic University of Korea, 505 Banpo-dong, Seocho-gu, Seoul 137-701, Korea.

Tel. 82-2-2258-1441 Fax. 82-2-599-6771 E-mail: ahn-kj@catholic.ac.kr

the proper surgical technique before the procedure.

Several earlier studies assessed the ability of conventional MR imaging for determining the consistency of pituitary adenomas (1–6) or meningiomas (7–11). Only a few studies evaluated the consistency of pituitary macroadenomas using apparent diffusion coefficient (ADC) with diffusion weighted imaging (DWI) (12–14). To the best of the authors' knowledge, a detailed analysis of DWI and conventional MR imaging for prediction of consistency of the intra-axial brain tumor has not yet been reported. Therefore, the aim of this study was to evaluate the usefulness of DWI, ADC maps and conventional MR images for determination of intra-axial brain tumor consistency.

### Materials and Methods

#### Subjects

We evaluated the MR images of 23 patients (10 men, 13 women; mean age, 52 years ± 18.20) with intra-axial brain tumors, who had undergone MR examinations between 2004 and 2008. Histologic types were glioblastoma multiforme (n = 12), metastasis (n = 4), anaplastic oligodendroglioma (n = 2), pilocystic astrocytoma (n = 2), anaplastic oligoastrocytoma (n = 1), anaplastic astrocytoma (n = 1), atypical teratoid/rhabdoid tumor (n = 1).

#### MR Sequence and Image Analysis

MR images were acquired using 1.5T MR scanner (Signa Excite, GE Medical Systems, Milwaukee, WI, U.S.A. or Achieva, Philips Medical Systems, Best, Netherlands) with a 8-channel head coil. The conventional MR images and DW images were

obtained separately. The interval between two studies was shorter than 1 week. The axial spin echo T1-weighted image (T1WI) (repetition time (TR), 500–650 ms; echo time (TE), 12–13 ms; section thickness, 5 mm; field of view (FOV), 220 × 220 mm; matrix size, 256–320 × 224; number of excitation (NEX), 1) and the axial turbo spin echo T2-weighted image (T2WI) (TR, 3600–4500 ms; TE, 80–122 ms; Echo train length (ETL), 12–16; section thickness, 5 mm; FOV, 220 × 220 mm; matrix size, 256–320 × 220–256; NEX, 1–2) were used for the signal intensity (SI) measurement of the tumors.

DWI was performed using a single-shot spin-echo echo-planar imaging sequence (TR, 2330–2640 ms; TE, 65–89 ms; flip angle, 90°; section thickness, 5 mm; FOV, 230 × 230 mm; matrix size, 128 × 128; b-values of 0 and 1000 s/mm<sup>2</sup>).

Measurement of SI of the tumors on T1WI, T2WI, DWI and ADC maps was performed by placing regions of interest (ROIs) drawn in the tumors. The measured SIs of the tumors were normalized by dividing the tumor SI values by the SIs of the contralateral normal regions. In three patients with both cystic and solid components, each component was evaluated independently.

Qualitative observed SIs by two experienced neuroradiologists were graded into 7 steps (grade 3 to grade -3) on various images : grade 3, marked hyperintense; grade 2, moderate hyperintense; grade 1, mild hyperintense; grade 0, isointense relative to the normal brain parenchyma; grade -1, mild hypointense; grade -2, moderate hypointense; grade -3, black signal intensity.

Table 1. Relative SI Ratios and Qualitative SI Grade of MR Images

Consistency	Number	Relative SI Ratio				Qualitative SI Grade			
		T1WI	T2WI	ADC	DWI	T1WI	T2WI	ADC	DWI
Cystic	3	0.67 ± 0.50	2.73 ± 0.94	3.46 ± 0.55	0.38 ± 0.04	-2±1	2.67 ± 0.58	2.67 ± 0.58	-2.67 ± 0.58
Gelatinous	2	0.78 ± 0.10	1.96 ± 0.39	2.75 ± 0.29	0.78 ± 0.12	-2±1.4	2.50 ± 0.71	3.00 ± 0.00	-2.00 ± 0.00
Friable	6	0.79 ± 0.11	1.68 ± 0.20	1.84 ± 0.78	1.18 ± 0.26	-1.17 ± 0.41	1.50 ± 0.55	0.67 ± 1.37	0.17 ± 1.94
Soft	9	0.84 ± 0.12	1.78 ± 0.38	1.87 ± 0.62	1.49 ± 1.83	-1.11 ± 0.60	1.22 ± 0.67	1.33 ± 0.87	-0.67 ± 1.41
Firm	4	0.76 ± 0.08	1.81 ± 0.37	1.82 ± 0.67	0.77 ± 0.41	-1.25 ± 0.50	1.75 ± 0.50	1.50 ± 0.58	0.75 ± 1.71
Hard	2	0.83 ± 0.03	1.44 ± 0.52	1.63 ± 0.10	0.69 ± 0.36	-1.00 ± 0.00	1.00 ± 0.00	1.00 ± 0.00	-1.50 ± 0.71
p value		0.260*	0.037*	0.012*	0.728*	0.434 <sup>§</sup>	0.035 <sup>§</sup>	0.031 <sup>§</sup>	0.039 <sup>§</sup>

Note.— Values are expressed as mean ± standard deviation.

\* = calculated with one-way ANOVA.

<sup>§</sup> = calculated with Kruskal-Wallis test.

**Surgery**

All patients with intra-axial brain tumor underwent surgical resection performed by one neurosurgeon. 16 tumors were totally resected and 7 tumors were partially removed. At surgery, consistency of the tumors was classified as cystic, gelatinous, friable, soft, firm or hard.

**Statistical Analysis**

Statistical analysis was performed by using the Statistical Package for the Social Sciences. We used analysis of variance to assess the difference among the relative SI ratios on various MR images across tumor consistency group, followed by a post hoc comparison (Tukey’s honest significance test). The Kruskal-Wallis test was used to evaluate differences in qualitative observed SI grade on various MR images among all consistency groups. We evaluated the correlation between relative SI ratios and qualitatively observed SI grade from various MR images and tumor consistency assessed at surgery by using Pearson’s and Spearman’s correlation coefficient. A p value of less than 0.05 was considered a statistically significant difference.

**Results**

The table 1 and 2 summarize the relationship between MR imaging findings and consistency of intra-axial brain tumors. No correlation was found between tumor consistency and relative SI ratio on T1WI. Statistical analysis also revealed no significant correlation between tumor consistency and qualitative SI grade on T1WI.

**Relative SI Ratio of ADC and Tumor Consistency**

Representative conventional MR images and ADC maps are shown in Figs. 1–4. The mean ratio of ADC in the cystic consistency group was  $3.46 \pm 0.55$ ,  $2.75 \pm 0.29$  in the gelatinous group,  $1.84 \pm 0.78$  in the friable group,  $1.87 \pm 0.62$  in the soft group,  $1.82 \pm 0.67$  in the firm group, and  $1.63 \pm 0.10$  in the hard group. The difference in the mean ratio of ADC values among various consistency groups was statistically significant ( $p = 0.012$ ). Pearson’s correlation analysis also demonstrated that relative SI ratios of ADC was a significant predictor of tumor consistency ( $r = -0.586$ ,  $p$

Table 2. Correlation Coefficient between Tumor Consistency and Various MR Imaging

	Relative SI Ratio*				Qualitative SI Grade <sup>§</sup>			
	T1WI	T2WI	ADC	DWI	T1WI	T2WI	ADC	DWI
Correlation coefficient	0.329	-0.497	-0.586	0.099	0.312	-0.461	-0.376	0.377
p value	0.100	0.010	0.002	0.629	0.121	0.018	0.059	0.058

\* = calculated with Pearson’s correlation

§ = calculated with Spearman’s correlation




Fig. 1. Metastatic tumor with hard consistency in 44-year-old woman. (a) Axial T2WI shows a homogenous hyperintense (grade 1) mass. (b) Axial DWI shows a hypointense lesion (grade -2) with respect to normal white matter. (c) ADC maps shows a mass with increased diffusion coefficient (grade 1).




Fig. 2. Metastatic mucoepidermoid carcinoma with gelatinous consistency in 67-year-old man. (a) Axial T2WI shows homogenous hyperintense (grade 3) masses. (b) Axial DWI shows hypointense masses (grade -2) with respect to normal white matter. (c) ADC maps shows masses with increased diffusion coefficient (grade 3).




Fig. 3. Glioblastoma with friable consistency in 69-year-old man. (a) Axial T2WI shows a homogenous hyperintense (grade 2) mass. (b) Axial DWI shows a hypointense lesion (grade -1) with respect to normal white matter. (c) ADC maps shows a mass with slightly increased diffusion coefficient (grade 1).




Fig. 4. Glioblastoma with friable consistency in 42-year-old man. (a) Axial T2WI shows a homogenous hyperintense (grade 1) mass. (b) Axial DWI shows a hyperintense lesion (grade 2) with respect to normal white matter. (c) ADC maps shows a mass with slightly decreased diffusion coefficient (grade -2).

= 0.002) (Fig. 5). Cystic consistency tumor was associated with a significantly higher relative SI ratio than friable, soft, firm and hard groups (Tukey HSD  $p < 0.05$ ).

**Relative SI Ratio on Conventional MR Images and Tumor Consistency**

The mean SI ratio on T2WI in the cystic consistency group was  $2.73 \pm 0.94$ ,  $1.96 \pm 0.39$  in the gelatinous group,  $1.68 \pm 0.20$  in the friable group,  $1.78 \pm 0.38$  in the soft group,  $1.81 \pm 0.37$  in the firm group, and  $1.44 \pm 0.52$  in the hard group with a statistically significant difference ( $p = 0.037$ ). Significant correlation was found between relative SI ratio on T2WI and tumor consistency ( $r = -0.497$ ,  $p = 0.01$ ) (Fig. 6). Cystic tumor was associated with significantly higher SI ratio than hard tumor (Tukey HSD  $p < 0.05$ ). The SI ratio on T1WI showed no significant difference among various consistency groups.

**Qualitatively Observed SI Grade on Conventional MR Images, ADC and DWI and Tumor Consistency**

The qualitatively observed SI on T2WI in the cystic consistency group was  $2.67 \pm 0.58$ ,  $2.50 \pm 0.71$  in the gelatinous group,  $1.50 \pm 0.55$  in the friable group,  $1.22 \pm 0.67$  in the soft group,  $1.75 \pm 0.50$  in the firm group, and  $1.0 \pm 0.0$  in the hard group with a




Fig. 5. Box plot shows relative SI ratios on ADC in tumor consistency groups. Relative SI ratios on ADC are significantly correlated with intra-axial brain tumor consistency. \*, §, ¶ = significant different from cystic consistency group \*  $p = 0.02$ , §  $p = 0.014$ , ¶  $p = 0.032$

statistically significant difference ( $p = 0.035$ ) (Fig. 7). Spearman’s correlation analysis showed significant correlation between qualitatively observed SI grade and tumor consistency ( $r = -0.461$ ,  $p = 0.018$ ). Kruskal-Wallis test indicated that there was a significant difference between observed SI grade on ADC and DWI for consistency groups ( $p = 0.031$ ,  $0.039$ , respectively). But there was no significant correlation between tumor consistency and SI grade on ADC and




Fig. 6. Box plot shows relative SI ratios on T2WI in tumor consistency groups. Relative SI ratios on T2WI are significantly correlated with intra-axial brain tumor consistency. \*, §, ¶ = significant different from cystic consistency group \*  $p = 0.033$ , §  $p = 0.044$ , ¶  $p = 0.045$




Fig. 7. Box plot shows qualitative SI grade on T2WI in tumor consistency groups. SI grade on T2WI are significantly correlated with intra-axial brain tumor consistency.

DWI ( $p = 0.059, 0.058$ , respectively).

## Discussion

Several reports assessed the ability of conventional MR images and DWI to predict consistency of pituitary adenomas or meningiomas and their results varied (12–14). Our study is the first research that evaluated the consistency of intra-axial brain tumors by various MR images. We found significant correlation between relative SI ratios of tumors on T2WI and qualitatively observed SI grade on T2WI and tumor consistency. In the present study, softer tumors showed higher T2 SI and grade than harder tumors did. Pierallini et al. (12) described inverse relation between SI on T2WI and tumor consistency of pituitary macroadenoma. Other researchers reported no correlation between tumor consistency and SI on conventional MR images in pituitary adenomas and meningiomas (4–6, 11). Such divergent findings may result from different tissue component of tumors. Collagen content, cellularity, extracellular space and free water contents are strongly correlated with tumor consistency. Harder tumor has more collagen content, higher cellularity, lower extracellular space and free water content than softer tumor does. Whereas extracellular space and free water contents produce an increase in SI on T2WI, fibrous tissue and high cellularity cause a decrease in SI. Therefore harder tumor demonstrates lower SI on T2WI than softer tumor does.

We found no statistically significant correlation between tumor consistency and relative SI ratio on T1WI and qualitatively observed SI grade on T1WI. Our results agreed with those in previous studies that showed no significant difference in SI on T1WI among tumor consistency groups (12, 13). Fibrous tissue which is abundant in harder tumor decrease SI on T1WI and extracellular space and free water contents which are rich in softer tumor also produce a decrease in SI on T1WI. Therefore, T1WI is of limited value in the evaluation of tumor consistency.

In this study, relative ratios of ADC of harder tumors were significantly lower than those of softer tumors. Pearson's correlation analysis also demonstrated that relative ratios of ADC were significant predictors of tumor consistency. More collagen content, higher cellularity and lower extracellular space may inhibit

diffusion, resulting in a lower ADC (15, 16). In pituitary macroadenomas, a few reports evaluated the usefulness of DWI and ADC maps for determination of tumor consistency and their conclusions did not agree (12–14). Although line-scan diffusion-weighted imaging (LSDWI) (17) and periodically rotated overlapping parallel lines with enhanced reconstruction (PROPELLER) DWI (18) were less susceptible to artifact than echo-planar DWI, susceptibility artifact was inevitable particularly in pituitary region. Tumor consistency may also affected by necrosis, cystic degeneration and hemorrhage within tumors which may make tumor softer. In earlier studies, SI measurements were performed by placing ROIs only in solid portions of the tumors. However, we measured SI of the whole tumor. For ADC values, earlier researchers used absolute value of ADC for analysis. ADC values are affected by magnetic field inhomogeneity and other factors. So in this study, ADC values were normalized for interimager and interimage variability. They may make different result.

There were some limitations to this study. First, the sample size of 26 lesions is too small to confirm an exact relationship between tumor consistency and various MR images. Further studies with larger sample sizes are required to confirm these findings. Second, histologic analysis such as collagen content and quantitative estimation of fibrous tissue was not performed in this study. Third, some tumors were not totally removed. If totally removed, their consistencies may be rated differently. Lastly, intraoperative assessment of tumor consistency was qualitative. More precise quantification of tumor consistency may help in preoperative planning and we recommend that this should be performed in future studies.

In conclusion, the measured ratio of ADC, SI ratio and observed SI grade on T2WI can provide valuable information about the consistency of intra-axial brain tumor.

## References

1. Iuchi T, Saeki N, Tanaka M, Sunami K, Yamaura A. MRI prediction of fibrous pituitary adenomas. *Acta Neurochir (Wien)* 1998;140:779-786
2. Snow RB, Johnson CE, Morgello S, Lavyne MH, Patterson RH, Jr. Is magnetic resonance imaging useful in guiding the operative approach to large pituitary tumors? *Neurosurgery*

- 1990;26:801-803
3. Naganuma H, Satoh E, Nukui H. Technical considerations of transsphenoidal removal of fibrous pituitary adenomas and evaluation of collagen content and subtype in the adenomas. *Neurol Med Chir (Tokyo)* 2002;42:202-212; discussion 213
  4. Bahuleyan B, Raghuram L, Rajshekhar V, Chacko AG. To assess the ability of MRI to predict consistency of pituitary macroadenomas. *Br J Neurosurg* 2006;20:324-326
  5. Chakraborty S, Oi S, Yamaguchi M, Tamaki N, Matsumoto S. Growth hormone-producing pituitary adenomas: MR characteristics and pre- and postoperative evaluation. *Neurol Med Chir (Tokyo)* 1993;33:81-85
  6. Hagiwara A, Inoue Y, Wakasa K, Haba T, Tashiro T, Miyamoto T. Comparison of growth hormone-producing and non-growth hormone-producing pituitary adenomas: imaging characteristics and pathologic correlation. *Radiology* 2003;228:533-538
  7. Yamaguchi N, Kawase T, Sagoh M, Ohira T, Shiga H, Toya S. Prediction of consistency of meningiomas with preoperative magnetic resonance imaging. *Surg Neurol* 1997;48:579-583
  8. Suzuki Y, Sugimoto T, Shibuya M, Sugita K, Patel SJ. Meningiomas: correlation between MRI characteristics and operative findings including consistency. *Acta Neurochir (Wien)* 1994;129:39-46
  9. Maiuri F, Iaconetta G, de Divitiis O, Cirillo S, Di Salle F, De Caro ML. Intracranial meningiomas: correlations between MR imaging and histology. *Eur J Radiol* 1999;31:69-75
  10. Yrjana SK, Tuominen H, Karttunen A, Lahdesluoma N, Heikkinen E, Koivukangas J. Low-field MR imaging of meningiomas including dynamic contrast enhancement study: evaluation of surgical and histopathologic characteristics. *AJNR Am J Neuroradiol* 2006;27:2128-2134
  11. Kashimura H, Inoue T, Ogasawara K, et al. Prediction of meningioma consistency using fractional anisotropy value measured by magnetic resonance imaging. *J Neurosurg* 2007;107:784-787
  12. Pierallini A, Caramia F, Falcone C, et al. Pituitary macroadenomas: preoperative evaluation of consistency with diffusion-weighted MR imaging--initial experience. *Radiology* 2006;239:223-231
  13. Suzuki C, Maeda M, Hori K, et al. Apparent diffusion coefficient of pituitary macroadenoma evaluated with line-scan diffusion-weighted imaging. *J Neuroradiol* 2007;34:228-235
  14. Mahmoud OM, Tominaga A, Amatya VJ, et al. Role of PROPELLER diffusion-weighted imaging and apparent diffusion coefficient in the evaluation of pituitary adenomas. *Eur J Radiol* 2010.(Epub ahead of print)
  15. Kono K, Inoue Y, Nakayama K, et al. The role of diffusion-weighted imaging in patients with brain tumors. *AJNR Am J Neuroradiol* 2001;22:1081-1088
  16. Guo AC, Cummings TJ, Dash RC, Provenzale JM. Lymphomas and high-grade astrocytomas: comparison of water diffusibility and histologic characteristics. *Radiology* 2002;224:177-183
  17. Gudbjartsson H, Maier SE, Mulkern RV, Morocz IA, Patz S, Jolesz FA. Line scan diffusion imaging. *Magn Reson Med* 1996;36:509-519
  18. Pipe JG, Farthing VG, Forbes KP. Multishot diffusion-weighted FSE using PROPELLER MRI. *Magn Reson Med* 2002;47:42-52

## 확산강조영상과 고식적 자기공명영상을 이용한 수술 전 축내 뇌종양의 정도 평가

가톨릭대학교 의과대학 영상의학과

오문식 · 안국진 · 최현석 · 정소령 · 이운주 · 김범수

**목적:** 수술 전 축내 뇌종양의 정도를 평가하는데 확산강조영상 및 현성확산계수영상과 고식적 MR 영상의 유용성을 알아보고자 하였다.

**대상 및 방법:** 축내 뇌종양으로 수술을 받은 23명의 환자를 대상으로 T1, T2, 확산강조영상 소견을 후향적으로 분석하였다. 다양한 MR영상에서의 신호강도와 수술에서 평가한 종양의 정도를 정량적, 정성적으로 비교 분석하였다. 수술소견에서 종양의 정도는 낭성, 젤리같은, 쉽게 부서지는, 부드러운, 단단한, 딱딱한 정도로 나눴다. 세 명의 환자에서는 낭성 부분과 고형성 부분이 함께 있어서 각각에 대해서 평가하였다.

**결과:** 종양이 단단할수록 현성확산계수와 T2강조영상에서의 신호강도의 비는 더 낮았다 ( $p = 0.002$ ,  $p = 0.01$ ). 종양의 정도와 현성확산계수가 가장 강한 선형상관관계를 보였다 ( $r = -0.586$ ,  $p = 0.002$ ). 정성적 분석에서는 단단할수록 T2강조영상에서 정성적 신호강도 등급이 낮았다 ( $p = 0.018$ ). 그 외 다른 MR소견은 통계분석에서 종양의 정도와 유의한 상관관계를 보이지 않았다.

**결론:** 축내 뇌종양의 현성확산계수, T2강조영상에서 신호강도의 비와 정성적 신호강도 등급은 종양의 정도와 상관관계가 있었고 이는 수술 전 전략 수립에 많은 도움이 될 것으로 생각된다.

통신저자 : 안국진, (137-701) 서울특별시 서초구 반포동 505번지, 서울성모병원 영상의학과  
Tel. (02) 2258-1441 Fax. (02) 599-6771 E-mail: ahn-kj@catholic.ac.kr