

Clinical Article

Aneurysmal Subarachnoid Hemorrhage in Third and Fourth Decades of Life

Su-Yong Kim, M.D., Chul-Hee Lee, M.D., In Sung Park, M.D., Jae Ha Hwang, M.D., Soo Hyun Hwang, M.D., Jong Woo Han, M.D.
Department of Neurosurgery, Gyeongsang National University School of Medicine & Gyeongsang Institute of Health Sciences, Jinju, Korea

Objective : The aim of this study was to compare clinical characteristics of ruptured aneurysms in young adults, of the third and fourth decades of life, and to compare several clinical characteristics affecting the outcome of patients.

Methods : We retrospectively investigated 1459 patients who underwent surgery and endovascular treatment for ruptured cerebral aneurysms from June 1992 to December 2010 and compared clinical characteristics. We also reviewed pre-existing medical conditions and perioperative complications.

Results : Among 1459 patients, there were 21 patients (1.44%) in the third decade and 104 patients (7.13%) in the fourth decade of life. Within two age groups, 88 (70.4%) were male and 37 (29.6%) were female, a ratio of 2.37 : 1. In both groups, we observed the anterior cerebral artery (ACA) aneurysm with the most frequency ($p=0.028$). In general, favorable outcome was achieved in both age groups (90.5% and 81.7%, respectively). An initial univariate analysis showed Hunt-Hess grade, Fisher grade, location of aneurysm, and rebleeding significantly associated with outcome after aneurysm rupture. Further, multivariate analysis demonstrated that only Hunt-Hess grade (grade 4-5) was a risk factor for the outcome (odds ratio=9.730, 95% confidence interval 2.069-45.756, $p=0.004$).

Conclusion : The incidence of subarachnoid hemorrhage (SAH) was higher in the male population of the third and fourth decades of life. Aneurysms on the ACA were most frequently occurred in both age groups and the outcome of aneurysmal SAH among the third and fourth decades was favorable. Multivariate analysis revealed that high Hunt-Hess grade was a risk factor for patient's outcome.

Key Words : Aneurysm · Subarachnoid hemorrhage · Young adult.

INTRODUCTION

Subarachnoid hemorrhage (SAH) accounts for 10% of stroke and 20% of hemorrhage stroke, with an incidence of 10 in 100000 each year¹⁵. Aneurysmal SAH in patients younger than 40 years of age is uncommon^{11,33}. However, its incidence in young adults has recently increased³¹. Although the etiologic and prognostic features that characterize SAH in the middle aged and elderly may not apply to young adults. We undertook a study of a series of young adult patients who were surgically and endovascularly treated for ruptured cerebral aneurysms. We compared the clinical features between patients in the third and fourth decades of life with published reports and compared several clinical characteristics affecting the outcome in patients. In addition, risk factors for unfavorable outcome were also assessed.

MATERIALS AND METHODS

Patients and methods

We retrospectively reviewed the medical records of 1652 patients with SAH who were treated from June 1992 to December 2010. Among those 1652 patients, 1459 patients were treated for ruptured aneurysms. Patients with unruptured, traumatic, and mycotic aneurysms were excluded from this study. The analysis focused on young adult patients aged from 20 to 39 years of age. We compared those in the third decade of life with those in the fourth decade, while comparing several clinical characteristics, including sex, Hunt-Hess grade, Fisher grade, size, location of aneurysm, multiplicity and outcome. We also reviewed pre-existing medical conditions (previous history of hypertension) and perioperative complications (rebleeding, vasospasm). The locations of aneurysm were classified into four

• Received : March 15, 2012 • Revised : June 23, 2012 • Accepted : September 17, 2012

• Address for reprints : Chul-Hee Lee, M.D.

Department of Neurosurgery, Gyeongsang National University School of Medicine, 79 Gangnam-ro, Jinju 660-702, Korea
Tel : +82-55-750-8112, Fax : +82-55-759-0817, E-mail : chl68@gnu.ac.kr

• This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

groups : internal carotid artery (ICA), anterior cerebral artery (ACA), middle cerebral artery (MCA) and vertebrobasilar artery (VBA). Aneurysms were also divided into three groups based on their diameters : small (<10 mm), large (10-25 mm), and giant (>25 mm). The patient's outcome at discharge was




Fig. 1. Bar graph showing the age and sex distributions of 1459 patients who were surgically and endovascularly treated for aneurysmal subarachnoid hemorrhage.

categorized according to the Glasgow Outcome Scale (GOS), as a favorable outcome (good recovery and moderate disability) or an unfavorable outcome (severe disability, vegetative state and death). We compared several clinical characteristics affecting the outcome of the enrolled patients, including sex, age, previous history of hypertension, Hunt-Hess grade, Fisher grade, size, location and multiplicity of aneurysm, rebleeding and lastly, vasospasm.

Statistical analysis

All data are presented as a mean value±standard deviation. Differences between patient's groups were tested via univariate analysis using t-test, chi-square test, or Mann-Whitney U test, as appropriate. Logistic regression was used to derive odds ratios (OR) for multilevel categorical variables. Multivariate models were adjusted for age, gender, hypertension or not, Hunt-Hess grade, Fisher grade, size and location of aneurysm, multiplicity, rebleeding, and vasospasm. All analyses were performed using Predictive Analytics SoftWare Statistics 18.0. A 2-tailed probability value of <0.05 was considered to be statistically significant.

Table 1. Characteristics of patients and aneurysms in young adults in the third and fourth decades of life who were surgically and endovascularly treated for aneurysmal subarachnoid hemorrhage

| Clinical characteristics | No. of patients (%) | | p-value |
|----------------------------------|---------------------|---------------|---------|
| | Third decade | Fourth decade | |
| Male | 14 (66.7) | 74 (71.2) | 0.681* |
| Previous history of hypertension | 2 (9.5) | 22 (21.2) | 0.217* |
| Hunt-Hess grade | | | 0.491* |
| 1 | 4 (19.0) | 26 (25.0) | |
| 2 | 11 (52.4) | 35 (33.7) | |
| 3 | 1 (4.8) | 14 (13.5) | |
| 4 | 3 (14.3) | 21 (20.2) | |
| 5 | 2 (9.5) | 8 (7.7) | |
| Fisher grade | | | 0.504* |
| 1 | 3 (14.3) | 7 (6.7) | |
| 2 | 6 (28.6) | 25 (24.0) | |
| 3 | 4 (19.0) | 33 (31.7) | |
| 4 | 8 (38.1) | 39 (37.5) | |
| Size of aneurysm | | | 0.834* |
| Small | 17 (81.0) | 87 (83.7) | |
| Large | 4 (19.0) | 16 (15.4) | |
| Giant | 0 (0) | 1 (1.0) | |
| Location of aneurysm | | | 0.028† |
| ICA | 6 (28.6) | 17 (16.3) | |
| ACA | 12 (57.1) | 48 (46.2) | |
| MCA | 3 (14.3) | 30 (28.8) | |
| VBA | 0 (0) | 9 (8.7) | |
| Multiplicity | 2 (9.5) | 14 (13.5) | 0.622* |
| Rebleeding | 5 (23.8) | 12 (11.5) | 0.135* |
| Vasospasm | 3 (14.3) | 20 (19.2) | 0.594* |
| Favorable outcome | 19 (90.5) | 85 (81.7) | 0.328* |

*Chi-square test, †Mann-Whitney U-test. ICA : internal carotid artery, ACA : anterior cerebral artery, MCA : middle cerebral artery, VBA : vertebrobasilar artery

RESULTS

From June 1992 to December 2010, 1459 patients were surgically and endovascularly treated for aneurysmal SAH. From which, 973 were female and 486 were male. The age and sex distributions of the patients with aneurysmal SAH are shown in Fig. 1. The mean age of the patients was 54.9±11.6 years (ranged from 14 to 86). The mean age of the female patients (58.0±10.7) was higher than that of the male patients (48.9±11.1). Among the 1459 patients, there were 21 patients (1.44%) in the third decade of life [20-29 years (26.3±2.7)] and 104 patients (7.13%) in the fourth decade of life [30-39 years (35.5±2.9)]. Within the third and fourth decade age group, 88 (70.4%) were male and 37 (29.6%) were female, a ratio of 2.37 : 1 (p=0.001). Mean follow up rate in two age groups was 41 months (ranged from 0 to 206) and 30-day mortality after ictus was 12.8%.

The age, Hunt-Hess grade, Fisher grade, size of aneurysm, rebleeding, vasospasm and outcomes did not differ significantly between the two age groups. And the distribution of ruptured aneurysms did not also significantly different between the two age groups. Aneurysms on the ACA and ICA were prevalent in

the third decade and aneurysms on the ACA and MCA were prevalent in the fourth decade of life. In particular, ACA aneurysms occurred with the greatest frequency in both age groups (57.1% in third decade and 45.8% in fourth decade) ($p=0.028$) (Table 1).

Favorable outcome rate for third and fourth decade were 90.5% and 81.7%, respectively (Table 2). An initial univariate analysis showed Hunt-Hess grade, Fisher grade, location of aneurysm, and rebleeding which significantly associated with outcome after aneurysm rupture (Table 2). A multiple logistic regression analysis demonstrated that only high Hunt-Hess grade (grade 4-5) was a risk factor for outcome, with an adjusted OR of 9.730 and confidence interval between 2.069 and 45.756 ($p=0.004$) (Table 3).

DISCUSSION

Aneurysmal SAH among young adults is rare^{1,30}. Sixty percent of aneurysms are diagnosed in patients between 40 and 60 years of age, with 10 to 20% found in patients younger than 40 years of age¹. Previously, it was reported that the prevalence of aneurysmal SAH in young adults ranged from 5.3% to 18.4%^{4,9,11,17,22}. In our study, the prevalence was 8.57% (125/1459), which was not significantly different from the previous studies.

In general, it has long recognized that cerebral aneurysm occur more frequently in females than in males^{10,16,25}. It has been reported that aneurysms in infants and children have different distributions from those seen in adults^{2,11}. Further, a male predominance in young children with SAH has been reported in previous studies^{3,9,19,20,33}. In our study, male dominance was observed during both the third and fourth decades of life and the incidence of female aneurysmal SAH progressively increases with age. In contrast, some authors have reported that incidence is higher in female young adults^{11,17}. This discrepancy may reflect the fact that intracranial aneurysm formation and growth is associated with multiple factors in young adults²².

Aneurysms on the ACA and ICA were prevalent in the third decade and aneurysms on the ACA and MCA were

prevalent in the fourth decade of life. Especially, ACA aneurysms were most frequent in these age groups in our study. Previous authors^{8,19,20,23} have noted that a vast majority of ruptured aneurysms in children and adolescents occur in the anterior circulation, especially in the ICA. The ICA has a blood flow much greater than the anterior communicating artery and MCA. Such flow may exert intense hemodynamic stress on the

Table 2. Clinical characteristics of patients with aneurysmal subarachnoid hemorrhage by outcome

| Clinical characteristics | No. of patients (%) | | p-value |
|----------------------------------|---------------------|---------------------|---------|
| | Favorable outcome | Unfavorable outcome | |
| Sex | | | 0.524* |
| Male | 72 (69.2) | 16 (76.2) | |
| Female | 32 (30.8) | 5 (23.8) | |
| Decade | | | 0.328* |
| Third | 19 (18.3) | 2 (9.5) | |
| Fourth | 85 (81.7) | 19 (90.5) | |
| Previous history of hypertension | 18 (17.3) | 6 (28.6) | 0.232* |
| Hunt-Hess grade | | | 0.000* |
| 1 | 28 (26.9) | 2 (9.5) | |
| 2 | 46 (44.2) | 0 (0) | |
| 3 | 12 (11.5) | 3 (14.3) | |
| 4 | 14 (13.5) | 10 (47.6) | |
| 5 | 4 (3.8) | 6 (28.6) | |
| Fisher grade | | | 0.001* |
| 1 | 10 (9.6) | 0 (0) | |
| 2 | 29 (27.9) | 2 (9.5) | |
| 3 | 34 (32.7) | 3 (14.3) | |
| 4 | 31 (29.8) | 16 (76.2) | |
| Size of aneurysm | | | 0.073* |
| Small | 88 (84.6) | 16 (76.2) | |
| Large | 16 (15.4) | 4 (19.0) | |
| Giant | 0 (0) | 1 (4.8) | |
| Location of aneurysm | | | 0.031* |
| ICA | 18 (17.3) | 5 (23.8) | |
| ACA | 55 (52.9) | 5 (23.8) | |
| MCA | 26 (25.0) | 7 (33.3) | |
| VBA | 5 (4.8) | 4 (19.0) | |
| Multiplicity | 14 (13.5) | 19 (90.5) | 0.622* |
| Rebleeding | 10 (9.6) | 7 (33.3) | 0.004* |
| Vasospasm | 16 (15.4) | 7 (33.3) | 0.053* |

*Chi-square test. ICA : internal carotid artery, ACA : anterior cerebral artery, MCA : middle cerebral artery, VBA : vertebrobasilar artery

Table 3. Multivariate Logistic Regression Model results

| Clinical characteristics | p-value | Adjusted OR | 95% confidence interval | |
|-----------------------------|---------|-------------|-------------------------|--------|
| | | | Lower | Upper |
| Hunt-Hess grade (grade 4-5) | 0.004 | 9.730 | 2.069 | 45.756 |
| Fisher grade | 0.483 | 1.398 | 0.547 | 3.572 |
| Size | 0.531 | 1.038 | 0.923 | 1.168 |
| Location | 0.580 | 0.821 | 0.407 | 1.653 |
| Rebleeding | 0.529 | 1.546 | 0.399 | 5.996 |

OR : odds ratio

arterial walls and thus, result in the formation of an aneurysm and rupture. Controversy exists concerning whether there is a difference in the location of ruptured aneurysms among different age groups^{14,28}. Although the causes of these differences in the location of ruptured aneurysms between childhood and young adults are uncertain, there is one hypothesis. Padget²¹ proposed that in the anterior circulation, the development of the ICA is followed by that of the ACA and eventually, by that of the MCA. Thus, it is possible that a fragile, congenital portion of the ICA may develop into an aneurysm due to hemodynamic stress. Further, the development of an ACA aneurysm may take a longer time than the time for an ICA lesion to grow. We agree with this hypothesis, but further studies and investigations about the causes of these differences in the location of ruptured aneurysms among different age groups are needed in the future.

The mortality rate for SAH in the 1966 cooperative Study on Intracranial Aneurysms was 50% at 29 days²⁷ and 33% in a recent analysis of in-hospital deaths among SAH patients admitted through an emergency department⁶. In a population-based study by Broderick et al.², the 30-day mortality rate among all patients who suffered SAH was 45%, with the majority of deaths occurring in the first days after SAH. The most important determinants of 30-day survival have been previously elucidated as initial neurological status and age¹⁸. Outcome is reported to be poor in female patients. In addition, old age is also reported to be a strong predictor of poor outcome^{5,7,26}. Outcomes of aneurysmal SAH among young adults are generally favorable^{11,12,33}. Outcome results from previous studies were compatible with our results. Among 125 patients, 104 (83.2%) had a GOS better than 3 (90.5% in third decade and 81.7% in fourth decade). Further, when perioperative complications (rebleeding, vasospasm) did not occur, patient's outcome was more favorable. Chang et al.⁴ reported that young patients usually present with a good neurological status and thus, have lower incidences of intraventricular hemorrhage, hydrocephalus, symptomatic vasospasm, and other underlying medical problems. Therefore, these young patients have a relatively good outcome.

Location of aneurysm is associated with mortality. The mortality at 6 months in conservatively treated patients was 34% to 39% after rupture of an anterior circulation aneurysm. Mortality at 6 months in patients with multiple aneurysm approached 50% and was higher in those with aneurysm arising from the posterior circulation, 61%³². Schievink et al.²⁹ reported that survival was greatly dependent on the location of the aneurysm and the prognosis of ruptured posterior circulation aneurysm is poor. In our study, aneurysms arising from VBA had more unfavorable outcome ($p=0.031$).

In addition to aneurysm location, clinical grade on admission is also associated with mortality. Winn et al.³² reported that clinical grade on admission and mortality during the first 6 months after the initial hemorrhage showed a strong statistical correlation. This association was subsequently confirmed in a study by

Phillips et al.²⁴ and more recently by Lagares et al.¹³. In our study, when Hunt-Hess grade was low, Fisher grade was low, and thereby, patient's outcome was more favorable. In addition, multivariate analysis demonstrated that only high Hunt-Hess grade (grade 4-5) showed a risk factor for outcome [OR, 9.730 (confidence interval, 2.069 to 45.756)]. Thus, level of consciousness following SAH strongly influenced patient's outcome.

The limitation of this study is being a retrospective review and patient's data were collected from only one institution. A large prospective multicenter study about aneurysmal SAH of young adult will be necessary in the future.

CONCLUSION

The incidence of SAH was higher in the male population in the third and fourth decades of life. Aneurysms on the ACA were most frequent in these age groups. Outcome of aneurysmal SAH among the third and fourth decades was favorable. Multivariate analysis revealed that high Hunt-Hess grade was a risk factor for patient's outcome.

References

- Bonita R, Beaglehole R, North JD : Subarachnoid hemorrhage in New Zealand : an epidemiological study. *Stroke* 14 : 342-347, 1983
- Broderick JP, Brott TG, Duldner JE, Tomsick T, Leach A : Initial and recurrent bleeding are the major causes of death following subarachnoid hemorrhage. *Stroke* 25 : 1342-1347, 1994
- Bröcheler J, Thron A : Intracranial arterial aneurysms in children. Clinical, neuroradiological and histological findings. *Neurosurg Rev* 13 : 309-313, 1990
- Chang CH, Kim JM, Ahn JS, Kwon Y, Kwun BD : Spontaneous intracranial aneurysm in young adult patients : retrospective review of 157 patients. *J Korean Neurosurg Soc* 33 : 149-153, 2003
- Chiang VL, Claus EB, Awad IA : Toward more rational prediction of outcome in patients with high-grade subarachnoid hemorrhage. *Neurosurgery* 46 : 28-35; discussion 35-36, 2000
- Cross DT 3rd, Tirschwell DL, Clark MA, Tuden D, Derdeyn CB, Moran CJ, et al. : Mortality rates after subarachnoid hemorrhage : variations according to hospital case volume in 18 states. *J Neurosurg* 99 : 810-817, 2003
- Deruty R, Pelissou-Guyotat I, Mottolese C, Amat D, Bogнар L : Level of consciousness and age as prognostic factors in aneurysmal SAH. *Acta Neurochir (Wien)* 132 : 1-8, 1995
- Heiskanen O, Vilkkii J : Intracranial arterial aneurysms in children and adolescents. *Acta Neurochir (Wien)* 59 : 55-63, 1981
- Horiuchi T, Tanaka Y, Hongo K, Kobayashi S : Aneurysmal subarachnoid hemorrhage in young adults : a comparison between patients in the third and fourth decades of life. *J Neurosurg* 99 : 276-279, 2003
- Juvela S, Porras M, Poussa K : Natural history of unruptured intracranial aneurysms : probability of and risk factors for aneurysm rupture. *J Neurosurg* 93 : 379-387, 2000
- Kamitani H, Masuzawa H, Kanazawa I, Kubo T : Saccular cerebral aneurysms in young adults. *Surg Neurol* 54 : 59-66; discussion 66-67, 2000
- Kongable GL, Lanzino G, Germanson TP, Truskowski LL, Alves WM, Torner JC, et al. : Gender-related differences in aneurysmal subarachnoid hemorrhage. *J Neurosurg* 84 : 43-48, 1996
- Lagares A, Gómez PA, Lobato RD, Alén JF, Alday R, Campollo J : Prog-

- nostic factors on hospital admission after spontaneous subarachnoid haemorrhage. *Acta Neurochir (Wien)* **143** : 665-672, 2001
14. Lanzino G, Kassell NF, Germanson TP, Kongable GL, Truskowski LL, Torner JC, et al. : Age and outcome after aneurysmal subarachnoid hemorrhage : why do older patients fare worse? *J Neurosurg* **85** : 410-418, 1996
 15. Linn FH, Rinkel GJ, Algra A, van Gijn J : Incidence of subarachnoid hemorrhage : role of region, year, and rate of computed tomography : a meta-analysis. *Stroke* **27** : 625-629, 1996
 16. McConmick WF, Nofzinger JD : Saccular intracranial aneurysm : an autopsy study. *J Neurosurg* **22** : 155-159, 1965
 17. Ogungbo B, Gregson B, Blackburn A, Barnes J, Vivar R, Sengupta R, et al. : Aneurysmal subarachnoid hemorrhage in young adults. *J Neurosurg* **98** : 43-49, 2003
 18. Oshiro EM, Walter KA, Piantadosi S, Witham TF, Tamargo RJ : A new subarachnoid hemorrhage grading system based on the Glasgow Coma Scale : a comparison with the Hunt and Hess and World Federation of Neurological Surgeons Scales in a clinical series. *Neurosurgery* **41** : 140-147; discussion 147-148, 1997
 19. Ostergaard JR : A long-term follow-up study of juvenile aneurysm patients. *Acta Neurochir (Wien)* **77** : 103-109, 1985
 20. Ostergaard JR : Aetiology of intracranial saccular aneurysms in childhood. *Br J Neurosurg* **5** : 575-580, 1991
 21. Padget DH : The development of the cranial arteries in the human embryo. *Contrib Embryol* **32** : 207-261, 1948
 22. Park SK, Kim JM, Kim JH, Cheong JH, Bak KH, Kim CH : Aneurysmal subarachnoid hemorrhage in young adults : a gender comparison study. *J Clin Neurosci* **15** : 389-392, 2008
 23. Pasqualin A, Mazza C, Cavazzani P, Scienza R, DaPian R : Intracranial aneurysms and subarachnoid hemorrhage in children and adolescents. *Childs Nerv Syst* **2** : 185-190, 1986
 24. Phillips LH 2nd, Whisnant JP, O'Fallon WM, Sundt TM Jr : The unchanging pattern of subarachnoid hemorrhage in a community. *Neurology* **30** : 1034-1040, 1980
 25. Rinkel GJ, Djibuti M, Algra A, van Gijn J : Prevalence and risk of rupture of intracranial aneurysms : a systematic review. *Stroke* **29** : 251-256, 1998
 26. Rosenørn J, Eskesen V, Schmidt K : Age as a prognostic factor after intracranial aneurysm rupture. *Br J Neurosurg* **1** : 335-341, 1987
 27. Sahs AL, Perret GE, Locksley HB, Nishioka H : **Intracranial Aneurysms and Subarachnoid Hemorrhage : A Cooperative Study**. Philadelphia : Lippincott, 1969, pp276-280
 28. Sakaki S, Ohta S, Ohue S, Kohno K, Matsuoka K : Outcome in elderly patients with ruptured intracranial aneurysm. *Clin Neurol Neurosurg* **91** : 21-27, 1989
 29. Schievink WI, Wijdicks EF, Piepgras DG, Chu CP, O'Fallon WM, Whisnant JP : The poor prognosis of ruptured intracranial aneurysms of the posterior circulation. *J Neurosurg* **82** : 791-795, 1995
 30. van Gijn J, Rinkel GJ : Subarachnoid haemorrhage : diagnosis, causes and management. *Brain* **124** : 249-278, 2001
 31. Weir B : Unruptured intracranial aneurysms : a review. *J Neurosurg* **96** : 3-42, 2002
 32. Winn HR, Richardson AE, Jane JA : The assessment of the natural history of single cerebral aneurysms that have ruptured in Hopkins LN, Long DM (eds) : **Clinical management of intracranial aneurysms**. New York : Raven Press, 1982, pp1-10
 33. Yoshimoto T, Uchida K, Suzuki J : Intracranial saccular aneurysms in the first three decades. *Surg Neurol* **9** : 287-291, 1978