

## The Ascidian *Numb* Gene Involves in the Formation of Neural Tissues

Hong Ryul Ahn<sup>1</sup> and Gil Jung Kim<sup>†</sup>

Dept. of Marine Molecular Biotechnology, Gangneung-Wonju National University, Gangneung 210-702, Korea

**ABSTRACT** : Notch signaling plays fundamental roles in various animal development. It has been suggested that Hr-Notch, a Notch homologue in the ascidian *Halocynthia roretzi*, is involved in the formation of peripheral neurons by suppressing the neural fates and promoting the epidermal differentiation. However, roles of Notch signaling remain controversial in the formation of nervous system in ascidian embryos. To precisely investigate functions of Notch signaling, we have isolated and characterized Hr-Numb, a Numb homologue which is a negative regulator of Notch signaling, in *H. roretzi*. Maternal expression of *Hr-Numb* mRNAs was detected in egg cytoplasm and the transcripts were inherited by the animal blastomeres. Its zygotic expression became evident by the early neurula stage and the transcripts were detected in dorsal neural precursor cells. Suppression of Hr-Numb function by an antisense morpholino oligonucleotide resulted in larvae with defect in brain vesicle and palps formation. Similar results have been obtained by overexpression of the constitutively activated *Hr-Notch* forms. Therefore, these results suggest that Hr-Numb is involved in Notch signaling during ascidian embryogenesis.

**Key words** : Numb, Neural specification, Notch signaling, Ascidian.

### INTRODUCTION

Notch signaling is used in a variety of cellular contexts during animal development to specify distinct cell fate among individual cells. In vertebrate embryos, Notch signaling is important for the formation of central nervous system (CNS), proliferation of the undifferentiated neural progenitor cells, and generating a variety of neural cell types (Latimer & Appel, 2006; Artavanis-Tsakonas & Muskavitch, 2010). In *Drosophila*, the sensory organ precursor cells develop from a group of equipotent neurogenic epithelial cells, which use Notch signaling to select neural progenitor cells (Campos-Ortega, 1995; Lai, 2004). The neural progenitor cells again use Notch signaling to specify different cell fates following asymmetric cell division. The asymmetric cell

division gives rise to a pIIa cell in which the Notch signaling becomes activated (non-neural fate) and a pIIb cell in which this signaling remains inactive (neural fate). The difference in Notch signaling activity is established by Numb (Uemura et al., 1989; Guo et al., 1996). In this process, Numb protein is asymmetrically segregated and localized to the pIIb daughter cell, in which suppresses Notch signaling activity. Numb encodes an adaptor protein that is able to bind the Notch receptor and interacts with endocytic proteins such as alpha-adaptin and EPS15, and then functions to promote the pIIb cell fate (Santolini et al., 2000; Berdnik et al., 2002). Recently, it was reported that Numb inactivates Notch by promoting its endocytosis during asymmetric cell division in *Drosophila* embryos (Couturier et al., 2012). Interaction of Numb with the Notch signaling pathway appears to be conserved in animal development, particularly in specification of the neural precursor cells (Kandachar & Roegiers 2012).

Ascidians belong to the Urochordata. They are one member of the phylum Chordata with vertebrates and cephalochordates. The ascidian tadpole larvae possess a

<sup>†</sup> Corresponding author: Gil Jung Kim, Dept. of Marine Molecular Biotechnology, Gangneung-Wonju National University, 7 Jukheon-gil, Gangneung 210-702, Korea. Phone: +82-33-640-2415, Fax: +82-33-640-2849, E-mail: gjkim@gwnu.ac.kr

<sup>1</sup> Present address: Functional Food Center, Korea Institute of Science and Technology, Gangneung Institute, Gangneung 210-340, Korea

typical chordate body plan with a CNS composed of only ~100 neurons (Meinertzhagen et al., 2004). The simplicity of ascidian CNS makes it an ideal model for understanding the development and function of chordate-specific neuronal networks. Notch signaling is also involved in specification of the neural progenitor cells and neural tube patterning in ascidian embryos. Overexpression of Notch led to defects in neural tube closure and in the formation of brain vesicle, palps, and peripheral neurons (Akanuma et al., 2002). It was also reported that a Delta2/Notch-mediated relay from the posterior motor ganglion specifies the fate of anterior motor ganglion (Stolfi et al., 2011). During ascidian embryogenesis, however, function of Numb in Notch signaling remains unknown.

In this study, we isolated and characterized the ascidian *Numb* gene. The ascidian *Numb* transcript was found to be expressed both maternally and zygotically. Its zygotic expression is observed in dorsal neural precursor cells from the early neural stage embryo. Larvae injected with *Numb* morpholino oligonucleotide showed abnormalities in brain and palps formation, but not pigment cells.

## MATERIALS AND METHODS

### 1. Animals and embryos

Adults of the ascidian *Halocynthia roretzi* were collected or purchased from fishermen in the vicinity of the Marine Biology Center for Research and Education at Gangneung-Wonju National University, Gangneung, Korea. Naturally spawned eggs were fertilized with a suspension of sperm from another individual, and then raised in filtered seawater containing 50 µg/mL streptomycin sulfate and 50 µg/mL kanamycin sulfate at 10-13°C. Embryos were collected at appropriate stages and fixed for whole-mount in situ hybridization.

### 2. Isolation and characterization of cDNA clone for *Halocynthia Numb* gene

The cDNA fragments encoding a part of N-terminus

region of *Halocynthia Numb* were isolated by RT-PCR with degenerate oligonucleotides, 5'-CA(A/G)TGGCA(A/G)(A/C/G)(C/A)IGA(C/T)GA(A/G)G-3' for the upstream, 5'-(G/A)AAIGC(G/A)CAICCIACIGC(G/A)T-3' for the downstream, and 5'-(A/G)GTI(A/T)(G/C)ITI(C/T)TG(C/T)GCICCI GA-3' for the nested downstream primers from gastrula stage poly(A) RNA. Larger fragments of *Hr-Numb* covering the complete ORF were obtained by 5' and 3' rapid amplification of cDNA ends (RACE) using a SMART RACE cDNA amplification kit (Clontech). The primers for RACE were the following: 5'-GTCGTGATGG AACTACGAGACGCTGGATATG-3' for the upstream and 5'-GGCTTAAACGTTACCCGAATCCTTAATGGC-3' for the downstream. Molecular phylogenetic relationships among the Numb products were estimated with MEGA 5.05 program using the neighbor-joining method (Saitou & Nei, 1987). Sequence data used in this study were taken from GenBank databases, with following accession numbers: Homo a, *Homo sapiens* Numb isoform CRA-a (EAW81100.1); Homo d, *Homo sapiens* Numb isoform CRA-d (EAW81110.1); Homo e, *Homo sapiens* Numb isoform CRA-e (EAW81104.1); Homo f, *Homo sapiens* Numb isoform CRA-f (EAW81114.1); Mus 1, *Mus musculus* Numb1 (AAD47835.1); Mus 2, *Mus musculus* Numb2 (AAD47836.1); Gallus, *Gallus gallus* Numb (AAD49434.1); Xenopus, *Xenopus tropicalis* Numb (CAL49325.1); Danio 1, *Danio rerio* Numb (AAT85677.1); Danio 2, *Danio rerio* Numb-like (AAI07954.1); Ciona, *Ciona intestinalis* Numb (BAE06599.1); Drosophila, *Drosophila melanogaster* Numb isoform A (AAF52776.1).

### 3. Whole-mount in situ Hybridization

Whole-mount in situ hybridization was performed using a digoxigenin-labeled *Hr-Numb* antisense probe, as described previously (Miya et al., 1997; Lee et al., 2011). Specimens were hybridized with the probe at 50°C.

### 4. Microinjection of MOs

To suppress functions of Hr-Numb, we injected the

morpholino antisense oligonucleotide (MO, Gene Tools) into eggs as described by Kim et al. (2007). The nucleotide sequence of *Hr-Numb* MO was 5'-GCTTTGTCTTAT TGTCCTTATCATG-3'. The standard control MO provided by Gene Tools was used as a control experiment. MOs were dissolved in sterile distilled water with Fast Green and injected into fertilized eggs. The final concentration of each MO to be injected was approximately 1 mg/mL. The injected eggs were allowed to develop up to hatching stage.

## RESULTS AND DISCUSSION

To clarify the mechanism by which Notch signaling specify neural cells during ascidian embryogenesis, we isolated *Numb* homologue (*Hr-Numb*) in the ascidian *Halocynthia roretzi* by the classical RT-PCR and 5' and 3' RACE. In various *Numb* proteins, the N-terminal region, which contains the phosphotyrosine-binding (PTB) domain interacting with a wide variety of different proteins, is highly conserved among vertebrates and invertebrates (Berdnik et al., 2002; Reugels et al., 2006). Using degenerate oligonucleotide primers (see Materials and Methods), we amplified target fragments from the *Halocynthia* gastrula stage poly(A) RNA by RT-PCR. Then, a fragment of *Halocynthia Numb* covering the complete ORF was obtained by RACE. A cDNA clone for *Hr-Numb* was 1,624 base pairs in full length, and the predicted protein has 456 amino acids (Fig. 1). The predicted amino acid sequence contained a well-conserved N-terminus sequence between various *Numb* homologues. Although the over degree of amino acid identity was not high (about ~ 30%), the amino acid sequence of *Hr-Numb* PTB domain was highly conserved when compared with the PTB domains of other *Numb* proteins (data not shown).

In order to understand the relationships among the *Numb* genes from various animal groups, we assembled a molecular phylogenetic tree (Fig. 2). The phylogenetic tree clearly shows that *Hr-Numb* is an *Halocynthia* orthologue

of vertebrate *Numb*. *Numb-like* gene (e.g. Danio 2 in Fig. 2) is a related member of the *Numb* family of endocytosis promoting genes (Niikura et al., 2006). *Numb* and *Numb-like* share high sequence similarities at amino acid level. However, *Numb-like* has two specific motifs that were not found in *Numb*, in the N-terminal and middle regions (Zhong et al., 1997; Niikura et al., 2006). *Hr-Numb* did not contain these motifs, suggesting to be an orthologue of *Numb* rather than that of *Numb-like*. There is no report that *Ciona intestinalis*, another ascidian species, has a *Numb-like* gene.

Whole-mount in situ hybridization demonstrated that the maternal transcripts of *Hr-Numb* are distributed evenly in the egg cytoplasm (Fig. 3A). Cleavage of *Halocynthia* eggs is bilaterally symmetrical. At the 8-cell stage, the transcripts were distributed in the cytoplasm of the animal pole-side blastomeres, although the signal was weak in the vegetal pole-side blastomeres (Fig. 3B, b). This expression pattern was inherited by the early gastrula stage. The maternal transcripts became concentrated near the cytoplasm of the animal blastomeres (Fig. 3c-f). Zygotic expression of *Hr-Numb* was firstly observed in the dorsal neural precursor cells at the early neurula stage (Fig. 3G, arrows). At the tailbud stage, the expression became strongly in the A- and a-lineage neural precursor cells of the dorsomedial head region (Fig. 3H). These cells give rise to brain, peripheral neurons, and palps. The expression pattern of *Hr-Numb* is similar to that of *Hr-Notch* reported by Hori et al. (1997). *Hr-Notch* was maternally expressed and its zygotic expression was visible in the dorsal part of the A-lineage nerve cord precursor, and in the a-line brain and palps precursor cells at the neurula stage. Thus, it is suggested that *Hr-Numb* is involved in Notch signaling in which specify the anterior region of neural precursor cells during ascidian embryogenesis. The expression pattern of *Hr-Numb* is also similar to that of zebrafish *Numb*. In zebrafish, *Numb* transcripts were detectable in all regions of the embryos at the mid-blastula transition

```

CACTTCTGAATCTTGTGAACTCTTGAATAAACAACAACATGATAAGGACAATAAGACAAAGCTTTAGACGGAAAAAGAAACC 90
      M I R T I R Q S F R R K K K P

TAAAGTTCAGAGAGCACACAGCCACCGGAATGGCAACAAGATGAGACTGCTGTACAAGCCGGATCTTGTAATTTTCATGTGAAGTATCT 180
K V P E S T Q P P E W Q Q D E T A V Q A G S C K F H V K Y L

TGGGTGCGTAGAAGTTCAAGATTCTAGAGGGATGCAGGTCTGTGAACAAGCTGTGAAGCAACTAAAATCTGGGAGGAAAAAGAAGATCAA 270
G C V E V Q D S R G M Q V C E Q A V K Q L K S G R K K K I K

AGCCATATTATATGTGTCTCTGATGCTTTACGGGTCGTGGAAGACTCCAGCAAAGCATTGTTGCTAGATCAAACAATTGAGAAGGTTTC 360
A I L Y V S P D A L R V V E D S S K A L L L D Q T I E K V S

GTTTTGTGCTCCCGATCGCAATTACAACCGTCTTTCTCCTATATATGTGCTGATGGAACACGAGACGCTGGATATGCCATTCATTTTT 450
F C A P D R N Y N R A F S Y I C R D G T T R R W I C H S F F

TGCCATTAAGGATTCCGGTGAACGTTTAAGCCATGCTGTTGGTTGTCCTTCGCTGCTTGCCTTGAAAAAACAGAAAAGAGAACGAGA 540
A I K D S G E R L S H A V G C A F A A C L E K K Q K R E R D

TACAAGCGCGTTACTGCAAGATATGATAGTAAACGTACCATGTTTACAAGAGAAGGATCGTTTAGACCAAAAATGGAAAGTAAATCGA 630
T S G V T A R Y D S K R T M F T R E G S F R P K M E S E I E

GAAACCTCCACAACCTGAAATCACTGAAGTACCTGCAGACGTTTACAGCTCAAGTTCAAGTGCAGTACCACGTCACATGCTCCATTACA 720
K P S T T E I T E V P A D V H S S S S S A V P R P H A P L H

CCTATTGAGACAAGCCTCTTTAAGGAATTCAGTAACAAGAAAATGTTTGTGCTGAAGGTGATATAAACTCTCCTTTTAAAAAGAACTTTC 810
L L R Q A S L R N F S N K K M F A E G D I N S P F K R N L S

TTTGCAGTAAATGAATTACCATCTACCATTACAAGACAACAAGTCATGAATCCAGTCCAGAGGATGAATGGGAAAATGAAATGGACAC 900
L R V N E L P S T I T R Q Q V M N P V P E D E W E N E M D T

ACACTTTTCTGGACAGACTACAAAATCCCTTTTCTATGAATGATGATCCTTTCACCAAGGCTCCTCCAATTTCAAACCAATTTTCAGTGT 990
H F S G Q T T N A F S M N D D P F T K A P P I S K P I S A V

TCGCCCTACACAGCTTAATGGATGTGGTGTGGTGGATTACGTGATCATTGCTATTGGAATATCATCAGACAAGCCTAAACCACT 1080
R P T Q L N G C G D G G L R D H S L S I G I S S D K P K P L

GAACTCCAGCGCTCTACTGCTACATCTCCGGTCCGTGAAACAAATCCATGGGCAAGTATCGCTCAACATCAAACCTGGTCATAAACG 1170
N S S G P T A T S P V R E T N P W A S D I A Q H Q T G H K R

AACTCCGCTGAGAGCAGACCAATGGCTTTTAGGTTTGCAAAGCAAAGTGAAGTCAACAAGTTATGATGCAGCATAACGCATTGCAATC 1260
T P S E A D Q W L L G L Q S K V Q G Q Q V M M Q H N A F E S


AATGTCTATGATGCAACGAAACCTTCATCAGCAGCAAGGTATAATGCCATGTCTATGAACAAGTATCCAATGAATCAGCAACCATATGG 1350
M S M M Q R N L H Q Q Q G I M P M S M N K Y P M N Q Q P Y G

TTTTGCAAGTGTACTCAAATACTGTTTCCAACCTCAGTCTAATGCATTTATGACAGGAATGTGAACCTGAATAATTACTTTGTTT 1440
F A S V T Q N T V F Q P Q S N A F M T G M *

GTGTGCTACTGGTAGTATATACTAACCAGTATCTTTCAGTGATCAACAAAGAATCCTATCTCGATCGGTGGCTTTTCAATATTGTTAT 1530
ATCAGAGTTGGTTTCTAATTTTTCTCTAATTAAGTTGGTGAACCTTTTCATTTCTTACCAGTGAATAAATGAACGTGAAGCAGGA 1620
GTCG 1624

```

**Fig. 1. The nucleotide and deduced amino acid sequences of a cDNA clone for the *Hr-Numb*.** The sequence of the cDNA encompasses 1,624 bp including 5' and 3'untranslational regions. The ATG at the position 47-49 represents the putative start codon of the *Hr-Numb*-encoded protein of 456 amino acids. The termination codon is shown by an asterisk and the PTB domain is underlined. The 25 base pairs for MO design are shown by dotted lines around the putative start codon.




**Fig. 2. Molecular phylogenetic tree of Numb family members.** Phylogenetic relationships for amino acid sequences of the Numb family members including *Halocynthia* inferred from the neighbor-joining analysis with *Drosophila* sequence as outgroup. Results from bootstrap resampling (1,000 replicates) are shown above internal nodes. The scale bar indicates an evolutionary distance of 0.1 amino acid substitutions per sites.

stage (Reugels et al., 2006). Expression of zebrafish *Numb* was found in the midline from the head to the tail regions at the neurula stage, and then the signal was restricted in the brain and eye progenitor cells at the later stage. Therefore, it is likely that roles of Numb protein are conserved in Notch signaling, which specify neural precursor cells in chordate development.

Next, we attempted to inhibit functions of Hr-Numb

by injecting fertilized eggs with antisense morpholino oligonucleotide (MO). Antisense MO prevents translation of the coding region of the targeted mRNA. The timing of initiation of gastrulation was delayed about one hour in the *Hr-Numb* MO injected embryos (85%,  $n=47$ ) compared with normal embryos. Injection of *Hr-Numb* MO resulted in the loss of brain vesicle (Fig. 4B, black arrow) and palps (Fig. 4B, red arrow) in 89% ( $n=47$ )



**Fig. 3. Whole-mount in situ hybridization pattern of *Hr-Numb* expression during *Halocynthia* embryogenesis.** (A) A fertilized egg. (B, b) 8-, (C, c) 16-, (D, d) 32-, (E, e) 64-, (F, f) 110-cell stage embryos. (B-F) Vegetal pole views. (b-f) Animal pole views. (G) An early neurula. Anterior is to the left. Arrows indicate zygotic expression of *Hr-Numb* in the dorsoanterior neural precursor cells. (H) A tailbud stage embryo. *Hr-Numb* is expressed in the dorsal neural cells. Scale bars = 100  $\mu$ m.

of cases. However, the *Hr-Numb* MO-injected larvae showed formation of pigment cells, although the cells located outside of head region (Fig. 4B, black arrow). This might be caused by incomplete neural tube closure. No specific abnormality was observed in the larvae injected with control MO in 91% of cases ( $n=32$ ) (Fig. 4A). These results suggest that *Hr-Numb* is essential for brain vesicle and palps formation, but not for pigment cells formation. Similar results have been reported by Akanuma et al. (2002). Overexpression of the constitutively activated *Hr-Notch* forms resulted in larvae with defects in neural tube closure and brain vesicle formation. The activated *Hr-Notch* also suppressed formation of palps, but not pigment cells. Therefore, it is plausible that



**Fig. 4. Effects of *Hr-Numb* MO injected into eggs.** (A) A larva injected with control MO shows normal morphology. A brain vesicle, pigment cells (black arrow) and palps (red arrow) were formed. Anterior is to the left. (B) A larva injected with *Hr-Numb* MO. Two pigment cells (black arrow) locate laterally in the outer region of head. Red arrow in B indicates no palps formation. Scale bar = 100  $\mu$ m.

*Hr-Numb* is involved in control of Notch signaling during ascidian embryogenesis.

In zebrafish, Numb and Numb-like are involved in differentiation of primitive erythrocytes (Bresciani et al., 2010). The zebrafish Numb and Numb-like knockdown experiments showed severe reduction or absence of embryonic erythrocytes. There is a report that Notch signaling controls the Numb protein level. In the developing chick CNS, a reciprocal negative regulation presents between Notch and Numb proteins, namely, high levels of Notch activation cause a reduction in the Numb and Numb-like protein levels (Chapman et al., 2006). Inhibition of Notch signaling by Numb is essential for various cell fate decisions. Future studies using *Hr-Numb* may

provide new insights for Notch signaling during ascidian embryogenesis.

### ACKNOWLEDGEMENT

This work was supported by a Gangneung-Wonju National University Research Grant (2010-0150) to G.J.K.

### REFERENCES

- Akanuma T, Hori S, Darras S, Nishida H (2002) Notch signaling is involved in nervous system formation in ascidian embryos. *Dev Genes Evol* 212:459-472.
- Artavanis-Tsakonas S, Muskavitch MA (2010) Notch: the past, the present, and the future. *Curr Top Dev Biol* 92:1-29.
- Berdnik D, Török T, González-Gaitán M, Knoblich JA (2002) The endocytic protein alpha-Adaptin is required for numb-mediated asymmetric cell division in *Drosophila*. *Dev Cell* 3:221-231.
- Bresciani E, Confalonieri S, Cermenati S, Cimbri S, Foglia E, Beltrame M, Di Fiore PP, Cotelli F (2010) Zebrafish numb and numblike are involved in primitive erythrocyte differentiation. *PLoS One* 5:e14296.
- Campos-Ortega JA (1995) Genetic mechanisms of early neurogenesis in *Drosophila melanogaster*. *Mol Neurobiol* 10:75-89.
- Chapman G, Liu L, Sahlgren C, Dahlqvist C, Lendahl U (2006) High levels of Notch signaling down-regulate Numb and Numblike. *J Cell Biol* 175:535-540.
- Couturier L, Vodovar N, Schweisguth F (2012) Endocytosis by Numb breaks Notch symmetry at cytokinesis. *Nat Cell Biol* 14:131-139.
- Guo M, Jan LY, Jan YN (1996) Control of daughter cell fates during asymmetric division: interaction of Numb and Notch. *Neuron* 17:27-41.
- Hori S, Saitoh T, Matsumoto M, Makabe KW, Nishida H (1997) *Notch* homologue from *Halocynthia roretzi* is preferentially expressed in the central nervous system during ascidian embryogenesis. *Dev Genes Evol* 207:371-380.
- Kandachar V, Roegiers F (2012) Endocytosis and control of Notch signaling. *Curr Opin Cell Biol* 24:534-540.
- Kim GJ, Kumano G, Nishida H (2007) Cell fate polarization in ascidian mesenchyme/muscle precursors by directed FGF signaling and role for an additional ectodermal FGF antagonizing signal in notochord/nerve cord precursors. *Development* 134:1509-1518.
- Lai EC (2004) Notch signaling: control of cell communication and cell fate. *Development* 131:965-973.
- Latimer AJ, Appel B (2006) Notch signaling regulates midline cell specification and proliferation in zebrafish. *Dev Biol* 298:392-402.
- Lee WY, Ham HS, Kim GJ (2011) Expression of *Weel* gene in the ascidian, *Halocynthia roretzi* embryo. *Dev Reprod* 15:1-7.
- Meinertzhagen IA, Lemaire P, Okamura Y (2004) The neurobiology of the ascidian tadpole larva: recent developments in an ancient chordate. *Annu Rev Neurosci* 27:453-485.
- Miya T, Morita K, Suzuki A, Ueno N, Satoh N (1997) Functional analysis of an ascidian homologue of vertebrate Bmp-2/Bmp-4 suggests its role in the inhibition of neural fate specification. *Development* 124:5149-5159.
- Niikura Y, Tabata Y, Tajima A, Inoue I, Arai K, Watanabe S (2006) Zebrafish Numb homologue: phylogenetic evolution and involvement in regulation of left-right asymmetry. *Mech Dev* 123:407-414.
- Reugels AM, Boggetti B, Scheer N, Campos-Ortega JA (2006) Asymmetric localization of Numb: EGFP in dividing neuroepithelial cells during neurulation in *Danio rerio*. *Dev Dyn* 235:934-948.
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406-425.
- Santolini E, Puri C, Salcini AE, Gagliani MC, Pelicci PG, Tacchetti C, Di Fiore PP (2000) Numb is an

endocytic protein. *J Cell Biol* 151:1345-1352.

Stolfi A, Wagner E, Taliaferro JM, Chou S, Levine M (2011) Neural tube patterning by Ephrin, FGF and Notch signaling relays. *Development* 138:5429-5439.

Uemura T, Shepherd S, Ackerman L, Jan LY, Jan YN (1989) Numb, a gene required in determination of cell fate during sensory organ formation in *Drosophila* embryos. *Cell* 58:349-360.

Zhong W, Jiang MM, Weinmaster G, Jan LY, Jan YN (1997) Differential expression of mammalian Numb, Numbl-like and Notch1 suggests distinct roles during mouse cortical neurogenesis. *Development* 124:1887-1897.

---

(Received 2 October 2012, Received in revised form 19 October 2012, Accepted 17 November 2012)