

Clinical Article

Clinical Outcomes of Spontaneous Spinal Epidural Hematoma : A Comparative Study between Conservative and Surgical Treatment

Tackeun Kim, M.D., Chang-Hyun Lee, M.D., Seung-Jae Hyun, M.D., Ph.D., Sang Hoon Yoon, M.D., Ki-Jeong Kim, M.D., Ph.D., Hyun-Jib Kim, M.D., Ph.D.

Department of Neurosurgery, Spine Center, Seoul National University Bundang Hospital, Seoul National University College of Medicine, Seongnam, Korea

Objective : The incidence of spontaneous spinal epidural hematoma (SSEH) is rare. Patients with SSEH, however, present disabling neurologic deficits. Clinical outcomes are variable among patients. To evaluate the adequate treatment method according to initial patients' neurological status and clinical outcome with comparison of variables affecting the clinical outcome.

Methods : We included 15 patients suffered from SSEH. Patients were divided into two groups by treatment method. Initial neurological status and clinical outcomes were assessed by the American Spinal Injury Association (ASIA) impairment scale. Also sagittal hematoma location and length of involved segment was analyzed with magnetic resonance images. Other factors such as age, sex, pre-morbid medication and duration of hospital stay were reviewed with medical records. Nonparametric statistical analysis and subgroup analysis were performed to overcome small sample size.

Results : Among fifteen patients, ten patients underwent decompressive surgery, and remaining five were treated with conservative therapy. Patients showed no different initial neurologic status between treatment groups. Initial neurologic status was strongly associated with neurological recovery ($p=0.030$). Factors that did not seem to affect clinical outcomes included : age, sex, length of the involved spinal segment, sagittal location of hematoma, pre-morbid medication of antiplatelets or anticoagulants, and treatment methods.

Conclusion : For the management of SSEH, early decompressive surgery is usually recommended. However, conservative management can also be feasible in selective patients who present neurologic status as ASIA scale E or in whom early recovery of function has initiated with ASIA scale C or D.

Key Words : Spinal epidural hematoma · Surgery · Conservative management · Outcome.

INTRODUCTION

The incidence of spinal hematoma is rare, and epidural hematomas are the most common type of this disease¹⁹. Spinal epidural hematomas are divided into two subgroups; spontaneous spinal epidural hematomas (SSEH) and traumatic spinal epidural hematomas. The incidence of SSEH is estimated to be 0.1 per 100000 patients per year¹⁵. The clinical presentation is usually characterized by the acute onset of back or neck pain with rapidly progressive neurological deficits due to the compression of the spinal cord or spinal nerve roots.

Since a severe neurologic deficit is accompanied with SSEH, rapid diagnosis and treatment is important. As reported in the

literatures, the primary treatment option is decompressive surgery but the number of reports regarding spontaneous resolution of SSEH has increased^{8,10,12,26-28}. However, evidence based arguments for choosing the optimal treatment method have not yet been established. We tried to describe the adequate method of treatment according to the analysis of patients' neurological status and assess clinical outcome and prognostic factors by the comparison of variables affecting the clinical outcome.

MATERIALS AND METHODS

Retrospectively, 15 patients with spontaneous spinal epidural hematomas between 2004 and 2012 were included in the study.

• Received : June 4, 2012 • Revised : September 7, 2012 • Accepted : December 18, 2012

• Address for reprints : Seung-Jae Hyun, M.D., Ph.D.

Department of Neurosurgery, Spine Center, Seoul National University Bundang Hospital, Seoul National University College of Medicine, 82 Gumi-ro 173beon-gil, Bundang-gu, Seongnam 463-707, Korea

Tel : +82-31-787-7164, Fax : +82-31-787-4059, E-mail : neurospine@snuh.org

• This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

We reviewed the patient's medical records and image studies under approval of institutional review board.

Initial neurologic status and clinical outcome was assessed by the American Spinal Injury Association (ASIA) impairment scale (Table 1)²⁴. The hematoma location was assessed by magnetic resonance (MR) axial image. According to relative position to the spinal cord, sagittal location was classified as anterior and posterior. The involved segment was defined as present hematoma between center level of upper and lower discs by MR sagittal image. The duration of hospital stay was calculated with admission and discharge date. All of premorbid medications were identified using prescriptions.

Nonparametric statistical techniques were used to overcome the small sample size. And subgroup analysis according to clinical outcomes regardless initial neurological status and method of treatment was also carried. Continuous evaluated parameters were compared using the Mann-Whitney U test, and the categorical data were analyzed with the Fisher's exact test and Linear-by-Linear Association test. *p*-value less than 0.05 was considered statistically significant.

RESULTS

There were four male and eleven female patients and their median age was 63 years. The patients were divided into two groups, one with surgery and the other with conservative treatment. Decompressive laminectomy (n=6), hemilaminectomy (n=3) and cervical laminoplasty (n=1) with hematoma evacuation were performed in surgically treated patients. Conservatively treated patients received 10 mg of dexamethasone over an intravenous line after admission. One patient was excluded since he presented without neurologic deficits other than pain. Until

motor weakness was improved, 4×4 mg of dexamethasone was administered over an average time range of two days. The dose of steroid was then slowly tapered off during five to seven days (Fig. 1).

Two (13.3%) patients had SSEH at the cervical spine; five (33.3%) at the cervicothoracic junction; four (26.7%) at the thoracic spine; and four (26.7%) at the thoracolumbar junction. Twelve (80.0%) patients manifested epidural hematomas located posteriorly to the spinal cord. Only three (20.0%) had anteriorly located SSEH. Table 2 showed the summarized cases.

There were no statistical differences in age, sex and the number of the involved vertebral segments determined by MR imaging between the two groups. The surgically treated group showed tendency of poorer initial neurologic status as determined by the ASIA scale without a statistically significant difference (*p*=0.067). And the period of hospitalization was longer in the surgically treated group (*p*=0.040). Three patients (30.0%) in the surgical management group and three (60.0%) in conservative treated patients took antiplatelet drugs or anticoagulant therapy, without statistically significant differences. All conservatively treated patients and seven (70.0%) out of ten patients in the surgical group presented posteriorly located hematomas. Table 3 shows these results from the comparison of variables in relation to treatment methods.

Subgroup analysis according to clinical outcomes

Regardless of treatment methods, normal neurological status (ASIA scale E) after treatment showed a strong relationship with initial ASIA scale (*p*=0.030). However, there is no significant difference between treatment methods. Patients characteristics such as age, sex, premorbid medication of antiplatelets or anticoagulants did not seem to affect complete recovery. Also, sagittal loca-

Table 1. ASIA impairment scales

ASIA scale	Description
A	Complete : no motor or sensory function preserved
B	Incomplete : sensory but no motor function preserved below the neurologic level
C	Incomplete motor function (>50%) of the key muscles below the neurologic level, motor grade <III
D	Incomplete motor function (>50%) of the key muscles below the neurologic level, motor grade ≥III
E	Normal

ASIA : American Spinal Injury Association




Fig. 1. T2 oblique sagittal (A) and axial (B) MRI of a patient treated with conservative management. T2 oblique sagittal (C) and axial (D) magnetic resonance images after a 1-month follow-up demonstrate the disappearance of the epidural hematoma.

Table 2. Summary of included cases

Surgical management group										
ID	Sex	Age	Location			Medication	Hospital days	Initial ASIA scale	Follow-up ASIA scale	Surgery
			Level	Sagittal	Segments					
1	Male	68	C	Posterior	2	Aspirin, cilostazol	19	C	D	Total laminectomy
2	Female	55	CT	Anterior	8		60	D	E	Total laminectomy
3	Female	59	CT	Posterior	4		15	D	E	Hemilaminectomy
4	Female	93	TL	Posterior	12		9	D	D	Hemilaminectomy
5	Female	78	TL	Posterior	6	Warfarin	38	B	C	Total laminectomy
6	Female	76	CT	Posterior	7		10	B	D	Laminoplasty
7	Male	68	CT	Posterior	4		26	B	E	Hemilaminectomy
8	Female	53	T	Anterior	3		18	A	D	Total laminectomy
9	Female	35	CT	Anterior	6		110	C	C	Total laminectomy
10	Female	64	TL	Posterior	6	Aspirin	31	A	B	Total laminectomy

Conservative management group										
ID	Sex	Age	Location			Medication	Hospital days	Initial ASIA scale	Follow-up ASIA scale	Treatment
			Level	Sagittal	Segments					
11	Female	70	TL	Posterior	8	Aspirin, clopidogrel	2	E	E	Observation
12	Male	60	T	Posterior	7	Aspirin	10	D	E	Steroid
13	Female	60	T	Posterior	3		34	C	D	Steroid
14	Female	63	C	Posterior	2	Aspirin, clopidogrel	7	D	E	Steroid
15	Male	3	T	Posterior	3		7	C	E	Steroid

C : cervical, CT : cervicothoracic, T : thoracic, TL : thoracolumbar, ASIA : American Spinal Injury Association

tion and length of involved segments of hematomas did not correlated with neurological recovery. These analyses were summarized in the Table 4.

DISCUSSION

SSEH usually presents with acute symptoms such as neck or back pain, radiating pain, progressive weakness, and cauda equina syndrome due to compression of the spinal cord or nerve roots²³. Because these symptoms can mimic a disc herniation, epidural tumor or an infection, differential diagnosis is important. MRI has been currently recognized as the most accurate diagnostic method for establishing a differential diagnosis^{10,19,27}. The MR imaging appearance of SSEH is heterogeneously hyperintense on the T2-weighted MR images and homogeneously isointense on the T1-weighted images. But signals of MR imaging can be varied with the lapse of time. In some cases, T2-weighted images appear as heterogeneously hyperintense and T1-weighted images can range from homogeneously isointense

Table 3. Summary of variables related to treatment methods

	Surgical management	Conservative management	p-value
Age [†]	66.0 (54.5-76.5)	60.0 (31.5-66.5)	0.513
Hospital days [†]	22.5 (13.8-43.5)	7.0 (4.5-22.0)	0.040*
Involved segments [†]	6.0 (3.8-7.3)	3.0 (2.5-7.5)	0.513
Sex [†]			
	Male	2	0.560
	Female	8	
Antiplatelet drugs or anticoagulants [†]			
	Yes	3	0.329
	No	7	
Sagittal location [†]			
	Anterior	3	0.505
	Posterior	7	
Initial ASIA scale [§]			
	A	2	0.067
	B	3	
	C	2	
	D	3	
	E	0	

Continuous variables were presented as median (interquartile range). *Statistical significance, [†]Analyzed with Mann-Whitney U test, [‡]Analyzed with Fisher's exact test, [§]Analyzed with Linear-by-Linear Association. ASIA : American Spinal Injury Association

to hyperintense. Most of SSEH do not display enhancement²⁰, however, in the hyperacute stage, the lesion can be enhanced²⁵.

The traditional treatment of choice for SSEH has been surgical treatment such as decompressive laminectomy and hemo-

Table 4. Overall neurologic outcome by other variables

		Neurologic deficit after treatment		
		Present	Absent	p-value
Initial ASIA scale [†]	A	2	-	0.030*
	B	2	1	
	C	3	1	
	D	1	4	
	E	-	1	
Treatment [‡]	Surgery	7	3	0.119
	Conservative	1	4	
Age [§]		66.0 (54.8-77.5)	60.0 (55.0-68.0)	0.336
Sex [‡]	Male	1	3	0.282
	Female	7	4	
Medication [‡]	Yes	3	3	1.000
	No	5	4	
Sagittal location [‡]	Anterior	2	1	1.000
	Posterior	6	6	
Involved segments [§]		6.0 (3.0-6.8)	4.0 (3.0-8.0)	0.955

Continuous variables were presented as median (interquartile range). *Statistical significance, [†]Analyzed with Linear-by-Linear Association, [‡]Analyzed with Fisher's exact test, [§]Analyzed with Mann-Whitney U test. ASIA : American Spinal Injury Association

ma evacuation^{10,19-21}. Many authors emphasize the role of early surgery, especially within 48 hours in cases with incomplete spinal cord dysfunction, or 36 hours for complete spinal cord dysfunction^{1,11,17,22}. Patients with severe neurological deficits preoperatively had poor clinical outcomes^{2,11,16}. Similarly, in the analysis of clinical outcomes of this study, better initial neurological status correlated with an improved clinical outcome.

On the other hand, there are several reports regarding the spontaneous resolution of SSEH without surgery. The authors chose non-surgical treatment in the case of rapid improvement of neurological deficits, inappropriate medical conditions for operation such as coagulopathy, and the refusal of surgery^{8,10,12,26-28}. A literature review revealed that only 10 out of 64 patients treated with conservative management experienced incomplete recovery because they tended to manifest a milder presentation compared to patients with surgical management. However, a literature review revealed no factors that advocate conservative treatment in SSEH¹⁰.

A high dose of methylprednisolone was the generally accepted treatment method for acute spinal cord injury³. But, another double blind randomized clinical trial showed no significant difference between the high dose (1000 mg daily) and the low dose (100 mg daily) treatment groups⁴. In this study, considering the side effects of treatment with a high dose of steroids and the patients' neurologic status, conservative group patients were treated with the low-dose protocol as the equivalent dose of dexamethasone.

Previous studies have reported that the segment involved most frequently in SSEH was the cervicothoracic junction and the sagittal location of the hematoma in the spinal canal was predominantly the dorsal side, which correlates with our results.

The posterior internal vertebral venous plexus is bigger and more convoluted than anterior one in the cervicothoracic junction. Furthermore, posterior plexus is uncovered by ligamentous structures. Therefore, it seems to play an important role in the etiology of SSEH⁶. Others, however, believe arterial rupture as the origin of the hematoma, as the intrathecal pressure is greater than the venous epidural pressure⁷. Additional risk factors were discussed in several studies such as anticoagulation therapy, antiplatelet drugs, coronary thrombolysis, and hypertension^{5,13,14,18,29,30}. These factors can play an important role in the progression of the hematoma¹⁹.

Several reports have stressed the issue that immediate replacement therapy in patients with impaired coagulation prevents progression of the hematoma with improvement of neurological deficits,

excluding the need for an operative intervention⁹. In the present investigation, two patients had been taking aspirin alone, two were treated with aspirin with clopidogrel, one was treated with aspirin with cilostazol and a remaining patient was treated with warfarin. Three of five patients (60.0%) managed conservatively had been taking antiplatelet drugs, but only 3/10 patients (30.0%) in the surgical group took medication. According to Groen et al., in the context of impaired coagulation, a hematoma can remain liquid for a longer period of time when compared to normal clotting, enabling spread of the hematoma into the spinal epidural space. In our study, patients who took antiplatelet drugs or anticoagulants did not tend to present themselves with mild neurological symptoms or exhibit improved clinical outcomes.

Due to the relatively small number of patients, especially in the conservative management group, it is difficult to draw a firm conclusion from this study. To overcome this weak point, nonparametric statistical methods and subgroup analysis were used, but statistical significance may have been overestimated or underestimated due to potential bias with low statistical power. A prospective randomized controlled trial should be performed to identify optimal treatments and therapies. However, there are ethical problems in forcing patients to submit to conservative management when they present progressive neurologic impairment. Furthermore, the rarity of SSEH is a serious obstacle to the design of a new, randomized controlled trial.

CONCLUSION

In the management of patients with SSEH, early decompressive surgical management is usually recommended because the

neurologic deficit is mainly caused by compression of the cord and nerve roots by the hematoma. However, conservative management can be considered in patients who present neurologic status as ASIA scale E or in whom early recovery of function has initiated with ASIA scale C or D.

References

- Baek BS, Hur JW, Kwon KY, Lee HK : Spontaneous spinal epidural hematoma. *J Korean Neurosurg Soc* 44 : 40-42, 2008
- Börm W, Mohr K, Hassepas U, Richter HP, Kast E : Spinal hematoma unrelated to previous surgery : analysis of 15 consecutive cases treated in a single institution within a 10-year period. *Spine (Phila Pa 1976)* 29 : E555-E561, 2004
- Bracken MB, Shepard MJ, Collins WF, Holford TR, Young W, Baskin DS, et al. : A randomized, controlled trial of methylprednisolone or naloxone in the treatment of acute spinal-cord injury. Results of the Second National Acute Spinal Cord Injury Study. *N Engl J Med* 322 : 1405-1411, 1990
- Bracken MB, Shepard MJ, Hellenbrand KG, Collins WF, Leo LS, Freeman DF, et al. : Methylprednisolone and neurological function 1 year after spinal cord injury. Results of the National Acute Spinal Cord Injury Study. *J Neurosurg* 63 : 704-713, 1985
- Clark MA, Paradis NA : Spinal epidural hematoma complicating thrombolytic therapy with tissue plasminogen activator--a case report. *J Emerg Med* 23 : 247-251, 2002
- Clemens HJ : [A contribution to the histology of the internal spinal venous plexus]. *Z Mikrosk Anat Forsch* 67 : 183-189, 1961
- Dinsmore AJ, Leonard RB, Manthey D : Spontaneous spinal epidural hematoma : a case report. *J Emerg Med* 28 : 423-426, 2005
- Duffill J, Sparrow OC, Millar J, Barker CS : Can spontaneous spinal epidural haematoma be managed safely without operation? A report of four cases. *J Neurol Neurosurg Psychiatry* 69 : 816-819, 2000
- García López A, Pérez Lara JM, Herrainz Hidalgo R, Puente Gonzalo E : Spinal epidural hematoma following thrombolytic therapy for acute myocardial infarction. *Orthopedics* 22 : 987-988, 1999
- Groen RJ : Non-operative treatment of spontaneous spinal epidural hematomas : a review of the literature and a comparison with operative cases. *Acta Neurochir (Wien)* 146 : 103-110, 2004
- Groen RJ, van Alphen HA : Operative treatment of spontaneous spinal epidural hematomas : a study of the factors determining postoperative outcome. *Neurosurgery* 39 : 494-508; discussion 508-509, 1996
- Hentschel SJ, Woolfenden AR, Fairholm DJ : Resolution of spontaneous spinal epidural hematoma without surgery : report of two cases. *Spine (Phila Pa 1976)* 26 : E525-E527, 2001
- Heppner PA, Monteith SJ, Law AJ : Spontaneous spinal hematomas and low-molecular-weight heparin. Report of four cases and review of the literature. *J Neurosurg Spine* 1 : 232-236, 2004
- Heye N : Is there a link between acute spinal epidural hematoma and aspirin? *Spine (Phila Pa 1976)* 20 : 1931-1932, 1995
- Holtás S, Heiling M, Lönnroft M : Spontaneous spinal epidural hematoma : findings at MR imaging and clinical correlation. *Radiology* 199 : 409-413, 1996
- Hsieh CF, Lin HJ, Chen KT, Foo NP, Te AL : Acute spontaneous cervical spinal epidural hematoma with hemiparesis as the initial presentation. *Eur J Emerg Med* 13 : 36-38, 2006
- Hussenbocus SM, Wilby MJ, Cain C, Hall D : Spontaneous spinal epidural hematoma : a case report and literature review. *J Emerg Med* 42 : e31-e34, 2012
- Kirazli Y, Akkoc Y, Kanyilmaz S : Spinal epidural hematoma associated with oral anticoagulation therapy. *Am J Phys Med Rehabil* 83 : 220-223, 2004
- Kreppel D, Antoniadis G, Seeling W : Spinal hematoma : a literature survey with meta-analysis of 613 patients. *Neurosurg Rev* 26 : 1-49, 2003
- Liao CC, Hsieh PC, Lin TK, Lin CL, Lo YL, Lee SC : Surgical treatment of spontaneous spinal epidural hematoma : a 5-year experience. *J Neurosurg Spine* 11 : 480-486, 2009
- Liao CC, Lee ST, Hsu WC, Chen LR, Lui TN, Lee SC : Experience in the surgical management of spontaneous spinal epidural hematoma. *J Neurosurg* 100 : 38-45, 2004
- Liu Z, Jiao Q, Xu J, Wang X, Li S, You C : Spontaneous spinal epidural hematoma : analysis of 23 cases. *Surg Neurol* 69 : 253-260; discussion 260, 2008
- Matsumura A, Namikawa T, Hashimoto R, Okamoto T, Yanagida I, Hoshi M, et al. : Clinical management for spontaneous spinal epidural hematoma : diagnosis and treatment. *Spine J* 8 : 534-537, 2008
- Maynard FM Jr, Bracken MB, Creasey G, Ditunno JF Jr, Donovan WH, Ducker TB, et al. : International Standards for Neurological and Functional Classification of Spinal Cord Injury. American Spinal Injury Association. *Spinal Cord* 35 : 266-274, 1997
- Nawashiro H, Higo R : Contrast enhancement of a hyperacute spontaneous spinal epidural hematoma. *AJNR Am J Neuroradiol* 22 : 1445, 2001
- Pahapill PA, Lownie SP : Conservative treatment of acute spontaneous spinal epidural hematoma. *Can J Neurol Sci* 25 : 159-163, 1998
- Schröder J, Palkovic S, Wassmann H : Spontaneous spinal epidural haematoma : a therapeutical challenge? Report of an unusual case. *Emerg Med J* 22 : 387-388, 2005
- Silber SH : Complete nonsurgical resolution of a spontaneous spinal epidural hematoma. *Am J Emerg Med* 14 : 391-393, 1996
- Spengos K, Tsvigoulis G, Zakopoulos N : Could high blood pressure be the cause of acute spontaneous spinal epidural hematoma? *Eur J Emerg Med* 14 : 59, 2007
- Sung JH, Hong JT, Son BC, Lee SW : Clopidogrel-induced spontaneous spinal epidural hematoma. *J Korean Med Sci* 22 : 577-579, 2007