

RESEARCH COMMUNICATION

Increased Trend of Breast Cancer Mortality in Iran

Afsoon Taghavi¹, Zeinab Fazeli², Mohsen Vahedi³, Ahmad Reza Baghestani⁴, Asma Pourhoseingholi², Farnoosh Barzegar², Mohamad Amin Pourhoseingholi^{2*}

Abstract

Background: Breast cancer is the most commonly diagnosed cancer in women worldwide. In Iran, it ranks first among cancers diagnosed in women and is the fifth most common cause of death. The aim of this study was to present the mortality trends from breast cancer for Iranian women during a period of almost a decade, in order to provide update information regarding the likely future. **Methods:** We analyzed National death Statistics reported by the Iranian Ministry of Health and Medical Education from 1995 to 2004 to generate annual mortality rates/100,000, overall, by age group (<15, 15-49 and ≥50 years of age) and age standardized rate (ASR). **Results:** The age standardized mortality rate of breast cancer increased dramatically during these years from 1.40 to 3.52 per 100,000 and its mortality was increasing 151.4% for Iranian women, although it seemed that the rate leveled off from 2002 to 2004. Moreover the increasing rate was higher for those aged between 15-49 compared to age >50 years old. **Conclusion:** There is an increasing trend for breast cancer mortality in Iran. Thus, health education programs to rectify the lack of women awareness about breast cancer signs and effective screening are urgently needed.

Keywords: Breast cancer - mortality - burden - Iranian females

Asian Pacific J Cancer Prev, 13, 367-370

Introduction

Breast cancer (BC) is the most commonly diagnosed cancer in women worldwide (Parkin et al., 2005). Global statistics show the annual incidence of breast cancer is increasing and this is occurring more rapidly in countries with a low incidence rate of breast cancer (Parkin et al., 200; Wilson et al., 2004). It has been reported that each year over 502,000 women die from the disease (World Health Organization., 2009) and it causes more than \$7 billion direct economic loss (medical costs) per year worldwide (Forbes., 1997). BC is the most commonly diagnosed cancer and the second-leading cause of cancer-related deaths among women in the U.S (U.S. Cancer Statistics Working Group., 2010). It is also the most common tumor in European women and is the first cause of death by cancer in females (Izquierdo et al., 2008).

Breast cancer occurs more frequently in wealthy countries (Parkin et al., 1997) due to a higher prevalence of BC risk factors, such as older age at first pregnancy, low parity, high-calorie intake, sedentary occupation and use of hormonal replacement therapy (Kogevinas et al., 1997; Marks and Shinberg., 1998; Zheng et al., 1993). On the other hand, BC survival is lower in less affluent countries and in women with low income or educational level (Kogevinas et al., 1997).

Cancer is a major public health problem in Iran. Based on recent reports from the Ministry of Health and Medical Education (MOHME); it is the third cause of death in Iran after coronary heart disease and accidents (Naghavi et al., 2009). In Iran, breast cancer ranks first among cancers diagnosed in women (Sadjadi et al., 2005) comprising 24.4% of all malignancies with a crude incidence rate and ASR of 17.4 and 23.1 per 100,000, respectively (Mousavi et al., 2009). BC is the most frequent cancer in women population in Tehran too (Mohagheghi et al., 2009) and the fifth most common cause of death for Iranian women (Akbari., 2008). BC has increased in Iran since 1999 according to Iranian cancer registry data (Shamsa and Mohagheghi., 2002).

Hence, it is necessary and important to get accurate projections of age-specific breast cancer mortality rates. These projections are of major public health interest. The aim of this study was to present the mortality trends from BC for Iranian women during a period of almost a decade, in order to provide update information regarding time trends for this cancer.

Materials and Methods

National death Statistics Reported by the MOH&ME from 1995 to 2000 (registered death statistics for Iranian

¹Department of Pathology, ²Research Center for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, ³Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, ⁴ Department of Mathematics, Islamic Azad University - South Tehran Branch, Tehran, Iran *For correspondence: aminphg@gmail.com

population at the Information Technology and Statistic Management Center, MOH&ME) and from 2001 to 2004 (published by MOH&ME) (Naghavi., 2002; Naghavi., 2003; Naghavi., 2004) stratified by age group and cause of death (coded according to the 9th revision of the International Classification of Diseases [ICD-10] are included in this analysis. Breast cancer [ICD-10: C50] were expressed as the annual mortality rates/100,000, overall, by age group (<15, 15-49 and \geq 50 years of age) and age standardized rate (ASR) (Ahmad et al., 2001). The populations of Iran in 1995-2004 were estimated, using the census from 1996 conducted by Statistics Centre of Iran and its estimation according to population growth rate for years before and after national census.

Results

All death records for women due to BC from 1995 to 2004 are included in this study. The age standardized mortality rate of BC increased dramatically during these years from 1.40 to 3.52 per 100,000 (Figure 1 and Table 1) and BC mortality was increasing 151.43% for Iranian women, however it seems that the rate would be leveled off from 2002 to 2004. Moreover BC mortality was higher for older age (Table 1 and Figure 2). But the increasing rate was higher for age between 15-49 compare to age >50 years old. In younger age, the mortality from 0.74 per 100,000 in 1995 was increased to 2.08 per 100,000 in 2004 (181.08% increasing) whereas, in older age, there




Figure 1. Trends of Breast Cancer Mortality During the Period 1995-2004 (Crude Rate and Age Standardized Rate per 100,000)




Figure 2. Age Specific Rate (per 100,000) for Breast Cancer Mortality During the Period Rate Per 100,000)

Table 1. Age Specific Rate (per 100,000) and Age Standardized Rate (per 100,000) for Breast Cancer Mortality

Year	<15	15-49	\geq 50	Crude Rate	ASR
1995	0	0.74	4.63	0.96	1.4
1996	0.03	1.08	6.24	1.36	1.93
1997	0.02	1.14	7.47	1.54	2.23
1998	0.02	1.2	6.24	1.41	1.99
1999	0.05	1.49	8.93	1.91	2.74
2000	0.06	1.65	10.58	2.12	3.18
2001	0.01	1.82	9.88	2.38	3.11
2002	0	2.14	11.8	2.75	3.69
2003	0	2.1	11.9	2.72	3.69
2004	0	2.08	11.16	2.58	3.52

was 141.04% increasing from 1995 to 2004.

Discussion

There is an increasing trend for BC mortality in Iran and it is possible that, breast cancer will become a leading cancer in our country, although its mortality is still relatively low compared with Western industrialized countries.

Recently, breast cancer mortality figures in Europe varied between 12 deaths per 100,000 in Spain and 19.3 deaths per 100,000 women in Hungary (World Health Organization: Cancer., 2008). Since the 1990s, age-adjusted mortality rates for BC have declined in most of the developed world, particularly in the young and middle-aged groups (35–64 years). In the United Kingdom and Switzerland, BC mortality decreased about 30% between 1990 and 2006 (Bulliard et al., 2006), whereas in most Southern, Northern, and Western European countries the decline was between 15% and 25%, and in the Eastern Europe, breast cancer mortality only decreased moderately or remained stable during the same period (Hery et al., 2009; Autier et al., 2011). In contrast, Russia was experiencing an increase trend (Hirte et al., 2007), where the ASR increased until 2004 and decreased thereafter.

According to WHO mortality database, between 1990 and 2006, the ASRs for BC in women in three Asian countries/regions (Japan, Hong Kong and Korea) were lower than the ASRs in Western countries. Korea had the lowest ASRs among the three Asian countries regions, and the UK had the highest ASRs among Western countries. In Asia, an increasing trend was observed for the ASR in Japan and Korea, whereas the ASR in Hong Kong appeared to be slowly decreasing (Katanoda and Yako-Suketomo., 2010).

Iranian data suggested a similar decreasing trend, in compare to other Asian countries. The data of Korea (in the period 1984–2003) revealed a similar results to Iranian data according to increasing trend and also amount of mortality pre 100,000 (Choi et al., 2006). A study in China indicated that, between 2002 and 2008, the mortality from BC increased by 201% among urban women, while among rural women the rate did not show the significant increase. This study concluded that the reversed urban–rural is primarily caused by the increases among elderly urban women (He et al., 2011) and a Japanese study showed,

there has been a marked increase in years of potential life lost (YPLL) associated with breast cancer among women (from 0.06 years in 1950 to 0.30 years in 2000) (Kono et al., 2005).

Iran is located in the western part of Asia which in this region, breast cancer in women is number one, generally followed by gastric, oesophageal or cervical lesions (Moore., 2010). The incidence of the BC is rising in our country, patients present with advanced stage of disease and they are relatively younger (about 10 years) than their western counterparts (Harirchi et al., 2000; Mousavi et al., 2007). It seems that breast cancer affects Iranian women at least one decade younger than women in developed countries, with the mean age ranging from 47.1 to 48.8 years (Harirchi et al., 2004). And more than 36% of the tumors occur in women under 40 years old (Mousavi et al., 2006).

Early detection of breast cancer plays the leading role in reducing mortality rates and improving the patients' prognosis among women (Elmore et al., 2005; Hoerger et al., 2011). In US, BC mortality rates are now decreasing because of the widely generalized use of mammography screening and the improvement in treatment (IARC., 2002). These recent falls of BC mortality in westernized countries have been larger in younger than older women. The relative contribution of screening and better systemic treatment to this downward trend is complex to disentangle (Levi et al., 2005; Botha et al., 2003). A relevant impact of screening on breast cancer mortality has been reported (Threlfall et al., 2003; Fracheboud et al., 2004), and, mammography use was inversely related to trends in breast cancer mortality in various US states over the period 1985–2000 (Das et al., 2005).

In Korea, the National Cancer Screening Program against breast cancer in 2002 started and the five-year survival rates for BC have improved significantly from 78.0% in early 1993-1995 to 90.0% in 2004-2008 which indicated that improvement of the survival rate may be partially due to the early diagnosis of BC (Park et al., 2011).

Our study provides comprehensive projection for Burden of death due to BC, indicating that the trend of BC mortality in Iranian women was dramatically increased in recent decade. In Iran, there is not a national screening program for controlling and early diagnosis of BC. Early detection of breast cancer is the key for decreasing its burden. So, specific screening measures should be implemented for Iranian women and special programs should be considered for younger women (Mousavi et al., 2006; Harirchi et al., 2011). On the other hand Iranian studies indicated that the women's awareness of BC warning signs and effective screening were very inadequate (Yavari and Pourhoseingholi., 2007; Montazeri et al., 2008). Thus, health education program to rectify the lack of women awareness is urgently needed. Indeed the focus of primary health care providers should be to raise awareness about breast care among women.

References

Ahmad O, Boschi Pinto C, Lopez AD, et al (2001). Age

- standardization of rates: a new WHO standard. Geneva, World Health Organization.
- Akbari ME (2008). Iran cancer report. Cancer Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Qom: Darolfekr.
- Autier P, Boniol M, Gavin A, et al (2011). Breast cancer mortality in neighbouring European countries with different levels of screening but similar access to treatment: trend analysis of WHO mortality database. *BMJ*, **28**, 343-?.
- Botha JL, Bray F, Sankila R, et al (2003). Breast cancer incidence and mortality trends in 16 European countries. *Eur J Cancer*, **39**, 1718-29.
- Bulliard JL, La Vecchia C, Levi F (2006). Diverging trends in breast cancer mortality within Switzerland. *Ann Oncol*, **7**, 57-9.
- Choi Y, Kim Y, Park SK, et al (2006). Age-Period-Cohort Analysis of Female Breast Cancer Mortality in Korea. *Breast Cancer*, **13**, 266-71.
- Das B, Feuer EJ, Mariotto A (2005). Geographic association between mammography use and mortality reduction in the US. *Cancer Causes Control*, **16**, 691-9.
- Elmore JG, Armstrong K, Lehman CD, et al (2005). Screening for breast cancer. *JAMA*, **293**, 1245-56.
- Forbes JF (1997). The incidence of breast cancer: the global burden, public health considerations. *Semin Oncol*, **24**, 20-35.
- Fracheboud J, Otto SJ, Van Dijck JA, et al (2004). Decreased rates of advanced breast cancer due to mammography screening in The Netherlands. *Br J Cancer*, **91**, 861-7.
- Harirchi I, Ebrahimi M, Zamani N, et al (2000). Breast cancer in Iran: a review of 903 case records. *Public Health*, **114**, 143-5.
- Harirchi I, Karbakhsh M, Kashefi A, et al (2004). Breast cancer in Iran: results of a multi-center study. *Asian Pac J Cancer Prev*, **5**, 24-7.
- Harirchi I, Kolahdoozan S, Karbakhsh M, et al (2011). Twenty years of breast cancer in Iran: downstaging without a formal screening program. *Ann Oncol*, **22**, 93-7.
- He M, Guo Q, Hu G (2011). Reversed urban–rural differences in breast cancer mortality (China, 2002–2008). *Breast Cancer Res Treat*, **126**, 231-4.
- Hery C, Ferlay JM, Boniol M, et al (2009). Quantification of changes in breast cancer incidence and mortality since 1990 in 35 countries with Caucasian-majority populations. *Ann Oncol*, **19**, 1187-94.
- Hirte L, Nolte E, Bain C, et al (2007). Breast cancer mortality in Russia and Ukraine 1963–2002: an age-period-cohort analysis. *Int J Epidemiol*, **36**, 900-6.
- Hoerger TJ, Ekwueme DU, Miller JW, et al (2011). Estimated Effects of the National Breast and Cervical Cancer Early Detection Program on Breast Cancer Mortality. *Am J Prev Med*, **40**, 397-404.
- IARC (2002). Globocan 2002 Database. Mortality. Available at <http://wwwwdep.iarc.fr/>.
- Izquierdo A, Gispert R, Saladie F, et al (2008). Analysis of cancer incidence, survival and mortality according to the main tumoral localizations, 1985-2019: Breast cancer. *Med Clin (Barc)*, **131**, 50-2.
- Katanoda K, Yako Suketomo H (2010). Comparison of Time Trends in Breast Cancer Mortality (1990–2006) in the World, from the WHO Mortality Database. *Jpn J Clin Oncol*, **40**, 182-?.
- Kogevinas M, Pearce N, Susser M, et al (1997). Social inequalities and cancer. Lyon. IARC Scientific Publications no. 138.
- Kono A, Misumi J, Misumi J (2005). The time trend of breast cancer mortality in Japan. *Arch Gynecol Obstet*, **272**, 187-90.
- Levi F, Bosetti C, Lucchini F, et al (2005). Monitoring the

- decrease in breast cancer mortality in Europe. *Eur J Cancer Prev*, **14**, 497-502.
- Marks NF, Shinberg DS (1998). Socioeconomic status differences in hormone therapy. *Am J Epidemiol*, **148**, 581-93.
- Mohagheghi MA, Mosavi-Jarrahi A, Malekzadeh R, et al (2009). Cancer Incidence in Tehran Metropolis: The First Report from the Tehran Population-Based Cancer Registry, 1998 – 2001. *Arch Iran Med*, **12**, 15-23.
- Montazeri A, Vahdaninia M, Harirchi I, et al (2008). Breast cancer in Iran: need for greater women awareness of warning signs and effective screening methods. *Asia Pac Fam Med*, **7**, 6-?.
- Moore MA, Eser S, Iginov N, et al (2010). Cancer epidemiology and control in North-Western and Central Asia - past, present and future. *Asian Pac J Cancer Prev*, **11**, 17-32.
- Mousavi SM, Gouya MM, Ramazani R, et al (2009). Cancer incidence and mortality in Iran. *Ann Oncol*, **20**, 556-63.
- Mousavi SM, Mohagheghi MA, Mousavi-Jarrahi A, et al (2006). Burden of breast cancer in Iran: a study of the Tehran population based cancer registry. *Asian Pac J Cancer Prev*, **7**, 571-4.
- Mousavi SM, Montazeri A, Mohagheghi MA, et al (2007). Breast cancer in Iran: an epidemiological review. *Breast J*, **13**, 383-91.
- Naghavi M (2002). Death report from 18 provinces in Iran. 1st edition. Ministry of Health and Medical Education, Tehran, Iran.
- Naghavi M (2003). Death report from 18 provinces in Iran. 1st edition. Ministry of Health and Medical Education, Tehran, Iran.
- Naghavi M (2004). Death report from 18 provinces in Iran. 1st edition. Ministry of Health and Medical Education, Tehran, Iran.
- Naghavi M, Abolhassani F, Pourmalek F, et al (2003). The burden of disease and injury in Iran 2003. *Popul Health Metr*, **15**, 7-9.
- Park SK, Kim Y, Kang D, et al (2011). Risk factors and control strategies for the rapidly rising rate of breast cancer in Korea. *J Breast Cancer*, **14**, 79-87.
- Parkin DM, Bray F, Ferlay J, et al (2005). Global cancer statistics, 2002. *CA Cancer J Clin*, **55**, 74-108.
- Parkin DM, Whelan SL, Ferlay J, et al (1997). Cancer incidence in five continents, vol. VII. Lyon. *IARC Scientific Publications* no. 143.
- Sadjadi A, Nouraie M, Mohagheghi MA, et al (2005). Cancer occurrence in Iran in 2002, an international perspective. *Asian Pac J Cancer Prev*, **6**, 359-63.
- Shamsa AZ, Mohagheghi MA (2002). Final report of the National project for cancer registry. Islamic Republic of Iran.
- Threlfall AG, Collins S, Woodman CB (2003). Impact of NHS breast screening on advanced disease and mortality from breast cancer in the North West of England. *Br J Cancer*, **89**, 77-80.
- U.S. Cancer Statistics Working Group (2010). U.S. Cancer Statistics: 2005 Incidence and Mortality. Atlanta GA: USDHHS, CDC and National Cancer Institute, www.cdc.gov/uscs.
- Wilson CM, Tobin S, Young RC (2004). The exploding worldwide cancer burden: the impact of cancer on women. *Int J Gynecol Cancer*, **14**, 1-11.
- World Health Organization (2009). Mortality database. WHO, , 2-6.
- World Health Organization: Cancer (2008). [http://www.who.int/mediacentre/factsheets/fs297/en]. accessed 30.07.
- Yavari P, Pourhoseingholi MA (2007). Socioeconomic factors association with knowledge and practice of breast self-examination among Iranian women. *Asian Pac J Cancer Prev*, **8**, 618-22.
- Zheng W, Shu XO, McLaughlin JK, et al (1993). Occupational physical activity and the incidence of cancer of the breast, corpus uteri, and ovary in Shanghai. *Cancer*, **71**, 3620-4.