

## The effect of a nutritional education program on the nutritional status of elderly patients in a long-term care hospital in Jeollanamdo province: health behavior, dietary behavior, nutrition risk level and nutrient intake

Bok Hee Kim<sup>1</sup>, Mi-Ju Kim<sup>2</sup> and Yoonna Lee<sup>3§</sup>

<sup>1</sup>Department of Food and Nutrition, Chosun University, Gwangju 501-759, Korea

<sup>2</sup>Graduate School of Education, Chosun University, Gwangju 501-759, Korea

<sup>3</sup>Nutrition Policy & Promotion Team, Korea Health Industry Development Institute, Osong Health Technology Administration Complex, 187 Osongsaeangmyeong2-ro, Gangoe-myeon, Cheongwon-gun, Chungcheongbuk 363-951, Korea

### Abstract

This study was conducted to assess improvements in nutritional status following the application of nutrition education to elderly patients in a long-term care hospital. The study was carried out from January to May 2009, during which a preliminary survey, a pretest, the application of nutrition education, and a post-test were applied in stages. The number of subjects at pretest was 81, and the number of participants included in the final analysis was 61 (18 men, 43 women), all of whom participated in both the nutrition education program and the post-test. The survey consisted of general demographic items, health behaviors, dietary behaviors, the Nutrition Screening Initiative checklist, and nutrient intake assessment (24 hour recall method). The nutrition education program lasted for four weeks. It included a basic education program, provided once a week, and mini-education program, which was offered daily during lunch times. The survey was conducted before and after the education program using the same assessment method, although some items were included only at pretest. When analyzing the changes in elderly patients after the nutritional education program, we found that, among subjective dietary behaviors, self-rated perceptions of health ( $P < 0.001$ ) and of depression ( $P < 0.001$ ) improved significantly and that dietary behavior scores also improved significantly ( $P < 0.001$ ), while nutritional risk levels decreased. In terms of nutrient intake, subjects' intake of energy, protein, fat, carbohydrate, calcium, phosphorus, iron, vitamin A, thiamin, riboflavin, niacin, and vitamin C all increased significantly ( $P < 0.001$ ). These results indicated that nutritional education is effective in improving the nutritional status of elderly patients. We hope that the results of this study can be used as preliminary data for establishing guidelines for nutrition management tailored to elderly patients in long-term care hospitals.

**Key Words:** Long-term care hospital for the elderly, nutritional education program, health behavior, dietary behavior, nutrient intake

### Introduction

With a declining birthrate and increasing life expectancy, Korea is rapidly aging in its demographic structure. While the population aged 65 or over was 3.8% in 1980, this grew to 11.0% in 2010 and is expected to reach 38.2% in 2050. It is also estimated that the old age dependency ratio will increase from 6.1% to 72.0% within this timeframe. An increase in old age dependency has already led to an increase in medical expenses; the percentage of the total medical cost that was expended on elderly care increased from 17.0% in 1999 to 30.5% in 2009, according to National Health Insurance statistics [1].

The aged population is, in general, an undernourished group in Korea. Although the nutrient intake of the total population has increased, the nutritional status of Korean elderly people has not much improved over the past ten years. While the rate of

under-nutrition based on the Korea National Health & Nutrition Survey was 14.6% in those aged 1 or over, it was 22.7% in those aged 65 or over (17.8% in men, 26.1% in women). The percentage of people with excess energy/fat consumption was 1.3% among the elderly, while it was 4.5% among the entire population [2]. These results imply that problems related to inadequate nutrition may be more prevalent among elderly people, and more serious in elderly women than in elderly men. Considering the sociodemographic factors that influence nutritional problems, these problems were also more serious for women than their male counterparts and for those who live alone, who live in rural areas, who have lower economic and educational levels or who are of an older age [3]. The problem of under-nutrition in the elderly results in increased health problems, leading to a rise in medical costs for the elderly. These problems cannot be solved solely by individuals, and the government is thus

§ Corresponding Author: Yoonna Lee, Tel. 82-43-713-8616, Fax. 82-43-713-8907, Email. ynlee@khidi.or.kr

Received: July 25, 2011, Revised: January 19, 2012, Accepted: January 19, 2012

©2012 The Korean Nutrition Society and the Korean Society of Community Nutrition

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

making various efforts to address the issue. Such efforts include the provision of meals to older people who are more likely to skip meals, hub-health promotion programs, health check-ups, long-term care insurance systems, and home-visit health management programs. However, nutritional programs for the elderly need to be further expanded.

As the population ages, the number of nuclear families grows, more women enter the workforce, and more households comprise aged single persons, it has become more difficult for family structures to provide care for the elderly. Considering this, the government introduced a long-term care insurance system in July 2008, which defines care for the elderly as being one of its social roles. Under this system, elderly people with a senility-related disorder who are considered to be unable to manage daily life by themselves for six months or more can receive long-term care [4]. Currently in Korea, long-term care hospitals and long-term care facilities provide services to senior citizens, with the two being run without a clear divide in terms of function or conditions [5]. With an increase in the aged population, the number of people eligible for the system is increasing, and this in turn, may be increasing insurance costs. Therefore, the main issue at stake is how to prevent currently healthy older people from becoming eligible for long-term care insurance, alongside the issue of how to reduce the cost of medical treatment in these facilities. In this context, the importance of nutrition management is emphasized as one approach. Nutrition management designed to attenuate nutritional problems can include nutrition education, nutrition counseling, tube feeding and other supplemental nutrition programs, in which proper methodology should be applied considering the characteristics of each individual.

When addressing the nutritional problems of the elderly, it is important to consider their characteristics and potential limitations; due to a weakening of physical and mental functions, the elderly may not only have less motivation to acquire new knowledge and lower studying abilities but may also be somewhat rigid in their habits, all of which factors could make nutrition interventions less effective. Previous studies, however, have indicated that there is a positive effect of nutrition education or counseling for the elderly [6-10].

This study was designed to measure the effect of nutritional education that is targeted to elderly patients in long-term care hospitals. Most of these patients are beneficiaries of the long-term care insurance system. In the hospital setting, dietitians manage patients' diet, but the cost of this is not covered by the insurance system, with patients having to pay for it [11]. This makes it difficult for patients or caretakers to engage with diverse nutrition management programs including education, counseling, tube feeding, and supplemental nutrition programs. This study therefore attempts to measure the extent to which long-stay patients' nutritional status can be improved when nutrition education is added to their basic meal management service, which is the minimum level nutrition service provided currently by hospitals. We hope that the results of this study can be used as preliminary

data for establishing guidelines for nutritional management programs targeted at elderly patients in long-term care hospitals.

## Subjects and Methods

### *Subjects*

Subjects comprised elderly, long-stay patients in a long-term care hospital in Naju in the Cheollanamdo province of South Korea, from January to May 2009. To evaluate improvements in their nutritional status following the application of nutrition education, we conducted a comparative analysis of changes before and after the program. The preliminary survey was conducted in January 2009, and based on the results of this a final, revised version of the survey questionnaire and methods were prepared. The pretest was conducted in February targeting 81 people who agreed to take part in the study, and based on its results, a nutritional education program was provided to these patients and their caretakers for four weeks. After the education was completed, we conducted a post-test, applying the same assessment method as with the pre-test, although some items were included only in the pretest. Considering the patients' age, a face-to-face interview was conducted by well-trained hospital dietitians and college students who were food and nutrition majors. The number of participants included in the final analysis was 61 (18 men, 43 women), excluding those who did not finish the nutritional education program or who did not completed the post-test.

### *General factors and health behaviors*

The survey consisted of five factors: general factors, health behaviors, dietary behaviors, a Nutrition Screening Initiative checklist, and nutrient intake assessment. General factors included household structure, educational level, and economic status. Health behaviors included self-rated perceptions about health and depression, smoking and drinking status, exercise, and previous participation in nutrition education.

### *Anthropometry and biochemical indices*

For data on subjects' height and weight, blood glucose, albumin, total cholesterol, HDL-cholesterol, triglycerides, and hemoglobin, we referred to the records of the hospital lab.

### *Dietary behaviors and nutrition screening initiative checklist*

Dietary behaviors were examined based on the 'Dietary Guidelines for the Elderly (Ministry of Health & Welfare)' [12]. The survey consisted of 12 questions which asked participants whether they ate regularly, had a balanced diet, ate various types of foods, and so on. To measure the level of compliance with dietary guidelines, a score of 3 was given for

yes, 2 for moderate, and 1 for no for each question, and these scores were used to calculate a total dietary behavior score. To assess participants' nutritional risk level, we applied the Nutrition Screening Initiative checklist (NSI) [13], in which the higher the total score received from 10 questions, the higher the risk level. Scores below 2 are regarded as indicating low risk, scores between 3 and 5 are considered moderate risk, while scores above 6 indicate high risk.

### Nutrient intake

The amount of foods consumed was examined over a five day period. Dieticians working in the long-term care hospital and college students studying food and nutrition measured both the amount of food provided and the leftovers of each meal for each individual to calculate consumption. The intake of foods brought from outside the hospital was also reflected in nutrient intake based on interviews with caretakers. To assess nutrient intake, CAN-PRO 3.0 (Computer Aided Nutritional Analysis Program, The Korea Nutrition Society (KNS)), a nutrition-evaluating program, was used to calculate the intake of each nutrient, and the average intake over the five days was used to measure nutrient intake data for each individual. We used the Dietary Reference Intakes for Koreans [14] to evaluate the adequacy of nutrient intakes.

### Nutritional education

Nutritional education was provided to the elderly patients and their caretakers in a long-term care hospital to improve their nutritional status and thus to help with treatment of their diseases. Implementing the education program was the responsibility of dieticians working within the hospital. The program was provided once a week for four weeks. Each session lasted for 60 minutes during the first three weeks, broken down into 20 minutes of information being given and 40 minutes of practical training and games. In the fourth week, the session lasted for 120 minutes, consisting of meal planning and nutrition counseling for each individual. Along with the four week basic education program, we also held mini-education sessions at lunch times for patients and caretakers. Educational materials consisted of PowerPoint resources made by our research team, and pamphlets and leaflets

issued by medical institutions or public health centers. For practical training, we utilized the food guide tower, a food model and a salt amount model, and used multiple visual materials with pictures to help the elderly, illiterate patients understand the contents.

Table 1 shows the detailed content of the nutritional program provided. In the first week, the education was focused on the physiological changes that occur with aging, the importance of nutrition management, and the need for a balanced diet. After this lecture, participants played a game consisting of putting food models into a food guide tower board. In the second week, participants learned about essential nutrients, their functions in the body, minimum requirements, and the major food sources of each nutrient. They also played the food traffic light game (KFDA), which was designed to enhance and reinforce their previous learning. In the third week, the education focused on dietary guidelines for the elderly (Ministry of Health & Welfare), and considering the characteristics of those patients who had one or more chronic diseases, a model based on salt intake was used to help them reduce salt intake. After this lecture, participants were given a chance to talk about their own eating habits and were encouraged to change their dietary behaviors. In the fourth week, participants practiced menu planning and received individualized nutrition counseling. Participants were encouraged to plan their own menu, and one-on-one tailored counseling was provided. To educate about adequate food composition, foods were prepared that were within the categories of rice, soup, side dishes, and snacks in the hospital cafeteria, and each food was provided in one serving size unit based on the food exchange system. Through this, participants could practice making better choices so as to develop a balanced diet. This approach was applied to the caretakers in the same way. The mini-nutritional education that accompanied the main program focused on helping participants to understand the characteristics of daily lunch menus and on how to apply basic food group knowledge in their diet. To reduce the amount of leftovers, a survey was conducted measuring how satisfied patients were with their meals. To measure the effect of the nutrition education program, a one group pretest-posttest design was applied, and changes in their dietary behaviors, health behaviors, Nutrition Screening Initiative checklist scores, and nutrient intake were assessed before and after program participation.

**Table 1.** The contents and tools of the nutritional education program

Week	Subject	Program	Tools
1st	Understanding basic food groups	<ul style="list-style-type: none"> <li>- The importance of nutrition management for the elderly</li> <li>- What is the food guide tower?</li> <li>- 'Make the food guide tower' game</li> </ul>	PowerPoint materials, handouts, food models, tools for games
2nd	Nutrients: functions and main food sources	<ul style="list-style-type: none"> <li>- The essential nutrients for our body</li> <li>- Functions and main food sources of each nutrient</li> <li>- Food &amp; nutrition traffic lights game</li> </ul>	PowerPoint materials, handouts, food models, tools for games
3rd	What is a healthy diet?	<ul style="list-style-type: none"> <li>- Introduce dietary guidelines for the elderly</li> <li>- Salt: how much do you eat?</li> <li>- Present individual dietary habits</li> </ul>	PowerPoint materials, handouts, salt amount model, presentation materials
4th	Menu-planning and nutrition counseling	<ul style="list-style-type: none"> <li>- Decide nutritional status of each individual</li> <li>- Individualized menu planning</li> <li>- Individualized nutrition counseling</li> </ul>	Evaluation sheet, foods for individualized menu-planning

### Statistical analysis

Statistical analysis was conducted using SPSS 17.0. Basic statistical analysis, including mean, standard deviation and frequency data for each survey item was conducted. To analyze differences by sex, a *t*-test was carried out for continuous variables, while an  $\chi^2$ -test was applied to the categorical variables. To assess the differences between pre-test and post-test results in all categories, paired *t*-tests were applied to measure statistical significance.

## Results

### General characteristics and health behaviors

Elderly patients' general characteristics and health behaviors are indicated in Table 2. The average age of the elderly men was 76.7 and that of the elderly women was 78.8. In terms of household structure, the percentage of subjects who indicated that they lived alone was the highest in both men and women (44.4% vs. 46.5%). However, of those not living alone, elderly men were more likely to live with a spouse, while elderly women tended to live with their children. Therefore, household structure differed

by sex ( $P < 0.01$ ). In terms of self-rated perceptions of health, 27.9% indicated being in good health, 59.0% normal, and 45.9% poor. The percentage of subjects who saw themselves as being in poor health was higher in elderly women (51.1%) than in elderly men (33.3%), which corresponds to results from national level statistics (27% in men vs. 45% in women) [4]. In self-rated perception of depression level, 50.8% indicated a good mental state, 29.5% normal and 19.7% poor. For other health related behaviors, 11.1% of elderly men reported that they smoked, and 5.6% of elderly men and 4.7% of elderly women indicated drinking alcohol. 72.2% of elderly men and 53.5% of elderly women reported exercising regularly. On the other hand, only 9.8% of participants had received nutrition education in the past.

### Anthropometry and biochemical indices

The anthropometry and biochemical indices of subjects are indicated in Table 3. The average height of elderly men was 161.6 cm and that of elderly women was 150.9 cm, while the average weight of elderly men was 52.3 kg and that of elderly women was 50.2 kg. The average BMI was 20.0 kg/m<sup>2</sup> in the case of men and 22.0 kg/m<sup>2</sup> in the case of women, which is somewhat low compared to the average BMI recorded in older people aged 70 or over (22.6 kg/m<sup>2</sup> in elderly men and 23.9

**Table 2.** General characteristics and health behaviors of participants

Variables		Total (n = 61)	Male (n = 18)	Female (n = 43)	$\chi^2$
Age (yrs)		78.2 ± 7.6 <sup>1)</sup>	76.7 ± 9.6	78.8 ± 6.7	2.306 <sup>2)</sup>
Household structure	Alone	28 (45.9) <sup>3)</sup>	8 (44.4)	20 (46.5)	12.011**
	With spouse	12 (19.7)	8 (44.4)	4 (9.3)	
	With children	10 (16.4)	1 (5.6)	9 (20.9)	
	With spouse & children	11 (18.0)	1 (5.6)	10 (23.3)	
Education	Illiterate	38 (62.3)	8 (44.4)	30 (69.8)	5.598*
	Beyond elementary school	23 (37.7)	10 (55.6)	13 (30.2)	
Economic status	Good	9 (14.8)	2 (11.1)	7 (16.3)	0.272
	Normal	36 (59.0)	11 (61.1)	25 (58.1)	
	Poor	16 (26.2)	5 (27.8)	11 (25.6)	
Self-rated perception of health	Good	17 (27.9)	5 (27.8)	12 (27.9)	2.439
	Normal	16 (26.2)	7 (38.9)	9 (20.9)	
	Poor	28 (45.9)	6 (33.3)	22 (51.1)	
Self-rated perception of depression	Good	31 (50.8)	9 (50.0)	22 (51.1)	0.114
	Normal	18 (29.5)	5 (27.8)	13 (30.2)	
	Poor	12 (19.7)	4 (22.2)	8 (18.7)	
Smoking	Yes	2 (3.3)	2 (11.1)	-	4.940*
	No	59 (96.7)	16 (88.9)	43 (100.0)	
Drinking	Yes	3 (4.9)	1 (5.6)	2 (4.7)	0.022
	No	58 (95.1)	17 (94.4)	41 (95.3)	
Regular exercise	Yes	36 (59.0)	13 (72.2)	23 (53.5)	1.841
	No	25 (41.0)	5 (27.8)	20 (46.5)	
Previous nutritional education	Yes	6 (9.8)	2 (11.1)	4 (9.3)	0.047
	No	55 (90.2)	16 (88.9)	39 (90.7)	

<sup>1)</sup> Mean ± SD

<sup>2)</sup> t-score

<sup>3)</sup> N (%)

\*  $P < 0,05$ , \*\*  $P < 0,01$

**Table 3.** The anthropometric measurements and biochemical data of participants

Variables	Total (n = 61)	Male (n = 18)	Female (n = 43)	t
Anthropometric measurements				
Height (cm)	154.0 ± 8.4 <sup>1)</sup>	161.6 ± 5.4	150.9 ± 7.3	1.142
Weight (kg)	50.8 ± 8.9	52.3 ± 6.5	50.2 ± 9.7	3.985*
BMI <sup>2)</sup> (kg/m <sup>2</sup> )	21.5 ± 3.7	20.0 ± 2.3	22.0 ± 4.1	0.771
Biochemical data				
Blood glucose (mg/dl)	104.7 ± 31.5	98.9 ± 24.5	106.3 ± 33.6	0.898
Albumin (g/dl)	4.2 ± 0.3	4.1 ± 0.3	4.3 ± 0.3	1.353
Total cholesterol (mg/dl)	191.1 ± 40.2	182.8 ± 35.7	194.8 ± 43.5	2.116
HDL-cholesterol (mg/dl)	43.3 ± 12.7	32.3 ± 5.1	48.2 ± 12.0	0.888
Triglycerides (mg/dl)	185.2 ± 108.2	150.7 ± 40.2	195.6 ± 121.8	3.699*
Hemoglobin (g/dl)	11.5 ± 1.3	12.1 ± 1.2	11.4 ± 1.2	0.038

<sup>1)</sup> Mean ± SD<sup>2)</sup> BMI (Body Mass Index) = Weight (kg) / Height (m)<sup>2</sup>\* *P* < 0,05

kg/m<sup>2</sup> in elderly women) from the Korea National Health & Nutrition Survey [2]. Fasting blood glucose was 98.9 mg/dl in men and 106.3 mg/dl in women. Other biochemical indices measured through blood testing included albumin, which was at 4.1 mg/dl in men vs. 4.3 mg/dl in women; total cholesterol at 182.8 mg/dl in men vs. 194.8 mg/dl in women; HDL-cholesterol at 32.3 mg/dl in men vs. 48.2 mg/dl in women; triglycerides at 150.7 mg/dl in men vs. 195.6 mg/dl in women; and hemoglobin, which was at 12.1 g/dl in men vs. 11.4 g/dl in women. The overall distribution of biochemical indices was found to be within the reasonable range.

#### Dietary behaviors and Nutrition Screening Initiative checklist

The results of the assessment of dietary behaviors, which was developed based on the Dietary Guidelines for the Elderly (Ministry of Health & Welfare), are described in Table 4. The

**Table 4.** Dietary behavior scores of participants

Variables	Total (n = 61)	Male (n = 18)	Female (n = 43)	t
Do you eat three meals a day regularly?	2.9 ± 0.1 <sup>1)</sup>	2.9 ± 0.2	2.9 ± 0.1	1.622
Do you eat foods slowly?	2.5 ± 0.7	2.7 ± 0.4	2.3 ± 0.8	13.027***
Do you try not to overeat?	2.4 ± 0.8	2.6 ± 0.6	2.4 ± 0.8	5.880**
Do you try to eat a balanced diet?	2.3 ± 0.7	2.5 ± 0.6	2.2 ± 0.8	3.567*
Do you try not to eat salty foods?	2.2 ± 0.7	2.2 ± 0.7	2.2 ± 0.7	0.056
Do you eat rice/breads/noodles at every meal?	2.8 ± 0.4	3.0 ± 0.0	2.7 ± 0.5	11.795***
Do you eat meats/fish/eggs/bean products daily?	2.3 ± 0.7	2.4 ± 0.6	2.3 ± 0.7	0.004
Do you eat foods cooked with oil at every meal time?	2.1 ± 0.6	2.3 ± 0.6	2.0 ± 0.5	3.582*
Do you eat milk & milk products daily?	1.8 ± 0.7	2.1 ± 0.7	1.6 ± 0.7	0.845
Do you eat vegetables daily?	2.7 ± 0.5	2.7 ± 0.5	2.6 ± 0.5	0.126
Do you eat fruits daily?	1.8 ± 0.6	1.8 ± 0.7	1.8 ± 0.6	0.143
Do you enjoy your meals?	2.3 ± 0.7	2.3 ± 0.6	2.3 ± 0.7	0.436
Total dietary behavior score	28.1 ± 3.2	29.5 ± 3.6	27.3 ± 2.8	1.053

<sup>1)</sup> Mean ± SD, Higher score indicates more desirable dietary behavior, with a possible total score from 12 to 36.\* *P* < 0,05, \*\* *P* < 0,01, \*\*\* *P* < 0,001

total score for the 12 questions measuring dietary behaviors was 29.5 in men and 27.3 in women, indicating that the dietary behaviors of elderly men were slightly better than those of elderly women. The results also showed a significant difference between elderly men and elderly women from the questions asking whether they ate food slowly, whether rice/bread/noodles formed a part of every meal (*P* < 0.001), whether they overate (*P* < 0.01), whether they had a balanced diet, and whether foods cooked with oils formed a part of every meal (*P* < 0.05).

Subjects' nutritional risk levels, as measured by the Nutrition Screening Initiative checklist (NSI), are indicated in Table 5. The

**Table 5.** Nutrition screening initiative checklist results

NSI Checklist	Total (n = 61)	Male (n = 18)	Female (n = 43)	t
I have an illness or condition that makes me change the kind and/or amount of food I eat	0.3 ± 0.7 <sup>1)</sup>	0.2 ± 0.6	0.4 ± 0.7	2.280
I eat fewer than 2 meals per day	0.4 ± 1.4	0.3 ± 0.9	0.5 ± 1.1	0.804
I eat few fruits or vegetable, or milk products	1.4 ± 0.9	1.8 ± 0.6	1.3 ± 0.9	11.435**
I have three or more drinks of beer, liquor or wine almost everyday	0.1 ± 0.4	0.1 ± 0.2	0.1 ± 0.1	0.009
I have tooth or mouth problems that make it hard for me to eat	1.4 ± 0.9	1.0 ± 1.3	1.5 ± 0.8	5.453*
I don't always have enough money to buy the food I need	1.3 ± 1.8	0.9 ± 1.7	1.4 ± 1.9	4.767*
I eat alone most of the time	0.8 ± 0.3	0.9 ± 0.2	0.8 ± 0.4	2.138
I take 3 or more different prescribed or over-the-counter drugs a day	0.3 ± 0.4	0.3 ± 0.4	0.3 ± 0.4	1.313
Without wanting to, I have lost or gained 10 pounds in the last 6 months	0.4 ± 0.8	0.2 ± 0.6	0.5 ± 0.8	5.816*
I am not always physically able to shop, cook and/or feed myself	0.9 ± 0.9	0.3 ± 0.7	1.1 ± 1.0	31.090***
Total NSI score	7.3 ± 3.5	6.0 ± 3.3	7.9 ± 3.5	5.323*
Nutritional status assessed by NSI score				
N (%)				
≤ 2 (Good)	6 (9.8)	2 (11.1)	4 (9.3)	
3-5 (Moderately risk)	16 (26.2)	7 (38.9)	9 (20.9)	4.280*
> 6 (High risk)	39 (63.9)	9 (50.0)	30 (69.8)	

<sup>1)</sup> Mean ± SD, Higher score indicates higher nutritional risk, with a possible total score from 0 to 21.\* *P* < 0,05, \*\* *P* < 0,01, \*\*\* *P* < 0,001

**Table 6.** Nutrient intakes of participants

	Total (n = 61)		Male (n = 18)		Female (n = 43)		t
	Intakes	%DRI	Intakes	%DRI	Intakes	%DRI	
Energy (kcal)	1,248.9 ± 212.4 <sup>1)</sup>	72.8 <sup>2)</sup>	1,333.9 ± 162.3	65.7	1,213.3 ± 222.2	75.8	0.404
Protein (g)	55.9 ± 11.4	120.5	59.0 ± 11.1	118.1	54.7 ± 11.3	121.6	0.503
Fat (g)	30.5 ± 7.5	-	31.7 ± 8.0	-	30.0 ± 7.4	-	0.810
Cholesterol (mg)	159.9 ± 38.6	-	165.6 ± 46.2	-	157.7 ± 35.4	-	0.077
Carbohydrate (g)	191.9 ± 32.7	-	207.1 ± 20.1	-	185.5 ± 35.0	-	3.956*
CPF ratio <sup>3)</sup>	60.7:17.7:21.6	-	61.7:17.4:20.9	-	60.3:17.8:21.9	-	0.288
Calcium (mg)	395.9 ± 97.2	51.6	405.1 ± 100.9	57.8	392.1 ± 96.6	49.0	0.756
Phosphorus (mg)	711.6 ± 143.7	101.6	744.8 ± 142.3	106.4	697.7 ± 143.7	99.7	0.741
Iron (mg)	11.4 ± 2.4	123.7	12.0 ± 2.3	120.5	11.3 ± 2.5	125.1	0.252
Sodium (mg)	3,099.9 ± 1,003.5	281.8	3,262.7 ± 1,033.0	296.6	3,031.8 ± 995.2	275.6	0.946
Potassium (mg)	2,549.9 ± 582.9	54.2	2,667.0 ± 609.4	56.7	2,499.7 ± 571.5	53.1	0.707
Vitamin A (µg)	500.4 ± 139.9	79.8	508.9 ± 138.2	72.7	496.5 ± 142.1	82.7	0.005
Thiamin (mg)	0.76 ± 0.15	67.4	0.80 ± 0.15	66.7	0.75 ± 0.15	67.8	1.083
Riboflavin (mg)	0.80 ± 0.18	62.4	0.82 ± 0.17	54.8	0.78 ± 0.18	65.5	0.206
Niacin (mg)	12.9 ± 2.7	88.2	13.5 ± 2.6	84.6	12.6 ± 2.7	89.7	0.296
Vitamin C (mg)	72.4 ± 18.1	72.4	74.2 ± 17.2	74.2	71.6 ± 18.7	71.6	0.000

<sup>1)</sup> Mean ± SD

<sup>2)</sup> % Estimated Energy Requirements (EER) for energy, % Recommended Intake (RI) for protein, calcium, phosphorus, iron, vitamin A, thiamin, riboflavin, vitamin C and % Adequate Intake (AI) for sodium, potassium

<sup>3)</sup> CPF ratio indicates percentage of energy intake from carbohydrate, protein and fat

\*  $P < 0,05$

total NSI score was 7.3, on average. Elderly men averaged 6.0 whereas elderly women averaged 7.9. Thus, the average score from both sexes were within the high-risk group, and the nutritional status of women was worse than that of men ( $P < 0.05$ ). By classifying findings using the criteria described earlier, we found that the rate of a low-risk group was 9.8% and that of a moderate-risk group was 26.2% while the rate of a high-risk group was 63.9%. It was also found that the percentage of participants belonging to the high-risk group was higher in elderly women than in elderly men ( $P < 0.05$ ).

### Nutrient intake

Nutrient intake findings are shown in Table 6. This indicates that the intake of most nutrients, including that of energy, were considerably lower than recommended intake levels. The daily energy intake of elderly men was 1,333.9 kcal (65.7% of the

Estimated Energy Requirement, EER) and that of elderly women was 1,213.3 kcal (75.8% EER). In both men and women, the intake of calcium (57.8% Recommended Intake (RI) in men vs. 49.0% RI in women), phosphorus (56.7% RI in men vs. 53.1% RI in women), thiamin (66.7% RI in men vs. 67.8% RI in women), riboflavin (54.8% RI in men vs. 65.5% RI in women), and vitamin C (74.2% RI in men vs. 71.6% RI in women) were below recommended levels.

### Changes to health behavior after nutritional education

Changes to health behaviors after nutritional education are shown in Table 7. In terms of self-rated perception of health status, those who reported being in 'poor' health decreased from 45.9% to 21.3% after the education program, while those who saw their health as 'normal' increased from 26.2% to 49.2%. This result shows that there was an improvement in participants' perceptions of their health, with negative perceptions significantly reduced ( $P < 0.001$ ). Considering self-rated perception of depression, the percentage of participants who said they were in a 'poor' mental state fell from 19.7% to 3.3%, while those responding as being at 'normal' increased from 29.5% to 36.1%, and 'good' responses increased from 50.8% to 60.7%. These results indicate that self-rated perceptions of depression also improved after the nutrition education program had taken place ( $P < 0.001$ ).

### Changes to dietary behaviors after nutrition education

Participants' changes to their dietary behaviors after nutritional education are shown in Table 8. The total dietary behavior score

**Table 7.** Changes in self-rated perceptions of health and depression in participants after nutritional education

	Before education (n = 61)	After education (n = 61)	$\chi^2$
Self-rated perception of health			
Good	17 (27.9)	18 (21.3)	4.271***
Normal	16 (26.2)	30 (49.2)	
Poor	28 (45.9)	13 (21.3)	
Self-rated perception of depression			
Good	31 (50.8)	37 (60.7)	4.619***
Normal	18 (29.5)	22 (36.1)	
Poor	12 (19.7)	2 (3.3)	

\*\*\*  $P < 0,001$

**Table 8.** Changes to dietary behavior scores in participants after nutritional education

Variables	Before education (n = 61)	After education (n = 61)	t
Do you eat three meals a day regularly?	2.9 ± 0.1 <sup>1)</sup>	3.0 ± 0.0	1.426
Do you eat foods slowly?	2.5 ± 0.7	2.7 ± 0.5	4.423*
Do you try not to overeat?	2.4 ± 0.8	2.7 ± 0.4	4.423*
Do you try to eat a balanced diet?	2.3 ± 0.6	2.6 ± 0.5	5.012*
Do you try not to eat salty foods?	2.2 ± 0.3	2.4 ± 0.5	4.619*
Do you eat rice/breads/noodles at every meal?	2.8 ± 0.4	2.9 ± 0.1	2.870
Do you eat meats/fish/eggs/bean products daily?	2.3 ± 0.7	2.6 ± 0.4	5.012**
Do you eat foods cooked with oil at every meal?	2.1 ± 0.6	2.3 ± 0.5	3.081*
Do you eat milk & milk products daily?	1.8 ± 0.7	2.2 ± 0.6	6.485**
Do you eat vegetables daily?	2.7 ± 0.5	2.8 ± 0.3	3.223*
Do you eat fruits daily?	1.8 ± 0.6	2.3 ± 0.5	5.850**
Do you enjoy your meals?	2.3 ± 0.7	2.7 ± 0.4	5.012**
Total dietary behavior score <sup>2)</sup>	28.1 ± 3.2	31.2 ± 2.0	10.838***

<sup>1)</sup> Mean ± SD, A higher score indicates more desirable dietary behavior, with a possible total score from 12 to 36.

\*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$

increased from 28.1 to 31.2 after the education program ( $P < 0.001$ ), and of the 12 questions, scores for 10 increased with statistical significance ( $P < 0.05$ ). Scores were most drastically increased in the following questions: ‘do you eat meats/fish/eggs/bean products daily?’, ‘do you eat fruits daily?’, ‘do you enjoy your meals?’ ( $P < 0.01$ ), ‘do you eat slowly?’, ‘do you try not to overeat?’, ‘do you try to have a balanced diet?’, ‘do you try not to eat salty foods?’, ‘do you eat foods cooked with oil in every meal?’, and ‘do you eat vegetables daily?’ ( $P < 0.05$ ).

**Table 9.** Changes to nutrition screening initiative checklist scores after education

NSI Checklist	Before education (n = 61)	After education (n = 61)	t
I have an illness or condition that makes me change the kind and/or amount of food I eat	0.3 ± 0.7 <sup>1)</sup>	0.3 ± 0.5	-
I eat fewer than 2 meals per day	0.4 ± 1.4	0.2 ± 0.5	2.215*
I eat few fruits, vegetables, or milk products	1.4 ± 0.9	1.2 ± 0.8	2.627*
I have three or more drinks of beer, liquor or wine almost everyday	0.1 ± 0.4	0.1 ± 0.3	-
I have tooth or mouth problems that make it hard for me to eat	1.4 ± 0.9	1.3 ± 0.9	-
I don't always have enough money to buy the foods I need	1.3 ± 1.8	1.3 ± 1.8	-
I eat alone most of the time	0.8 ± 0.4	0.5 ± 0.5	5.410***
I take 3 or more different prescribed or over-the-counter drugs a day	0.3 ± 0.4	0.3 ± 0.4	-
Without wanting to, I have lost or gained 10 pounds in the last 6 months	0.4 ± 0.8	0.4 ± 0.8	-
I am not always physically able to shop, cook and/or feed myself	0.9 ± 0.9	0.8 ± 0.7	-
Total NSI score	7.3 ± 3.5	6.4 ± 3.5	5.387***
Nutritional status assessed by NSI score	N (%)		
≤ 2 (Good)	6 (9.8)	8 (13.1)	
3-5 (Moderately risk)	16 (26.2)	16 (26.2)	1.780
> 6 (High risk)	39 (63.9)	37 (60.7)	

<sup>1)</sup> Mean ± SD, Higher score indicates higher nutritional risk, with a possible total score from 0 to 21.

\*  $P < 0.05$ , \*\*\*  $P < 0.05$

### Changes to nutrition screening initiative checklist score after nutritional education

Table 9 shows changes to the Nutrition Screening Initiative checklist scores after the nutrition education program. The total score on the Nutrition Screening Initiative checklist fell from 7.3 to 6.4, confirming that there were improvements in terms of nutritional risk level ( $P < 0.001$ ). Scores dropped significantly for three questions, namely ‘I eat alone most of the time’ ( $P < 0.001$ ), ‘I eat fewer than two meals per day’, and ‘I eat fruits, vegetables, or milk products’ ( $P < 0.05$ ). When scores were classified by risk level, the percentage of subjects in the high-risk group fell from 63.9% to 60.7% after the education program, while the number within the the low-risk group rose from 9.8% to 13.1%. These results confirm that there were improvements in *Nutrition Screening Initiative* checklist scores after the nutritional education program.

### Changes in nutrient intake after nutritional education

Changes in nutrient intakes after nutritional education are shown in Table 10. This indicates that intake of most nutrients increased significantly after the educational program. Energy intake increased from 72.8% of EER to 82.5% ( $P < 0.001$ ), and the intake of protein (120.5 → 136.6% RI), fat, carbohydrate, calcium (51.6 → 60.7% RI), phosphorus (101.6 → 118.5% RI), iron (123.7 → 143.5% RI), vitamin A (79.8 → 120.6% RI), thiamin (67.4 → 78.9% RI), riboflavin (62.4 → 73.0% RI), niacin (88.2 → 103.4% RI), and vitamin C (72.4 → 85.0% RI) also increased after the educational program ( $P < 0.001$ ). However, intakes of calcium, potassium, and riboflavin remained insufficient after the program, confirming the need for a proper nutritional assistance system that encourages the elderly to increase intake of these nutrients, which are more likely to be deficient in their diet.

**Table 10.** Changes to nutrient intakes in participants after nutritional education

Nutrients	Before education (n = 61)		After education (n = 61)		t
	Intakes	%DRI	Intakes	%DRI	
Energy (kcal)	1,248.9 ± 212.4 <sup>1)</sup>	72.8 <sup>2)</sup>	1,415.7 ± 232.3	82.5	9.921***
Protein (g)	55.9 ± 11.4	120.5	63.4 ± 11.6	136.6	9.990***
Fat (g)	30.5 ± 7.5	-	38.2 ± 7.2	-	9.683***
Cholesterol (mg)	159.9 ± 38.6	-	183.9 ± 39.3	-	8.996***
Carbohydrate (g)	191.9 ± 32.7	-	204.7 ± 35.4	-	8.927***
Calcium (mg)	395.9 ± 97.2	51.6	464.7 ± 99.7	60.7	9.797***
Phosphorus (mg)	711.6 ± 143.7	101.6	829.2 ± 148.6	118.5	9.875***
Iron (mg)	11.4 ± 2.4	123.7	13.3 ± 2.7	143.5	9.922***
Sodium (mg)	3,099.9 ± 1,003.5	281.8	3,130.7 ± 1,017.3	284.6	1.671
Potassium (mg)	2,549.9 ± 582.9	54.2	2,383.6 ± 449.3	50.7	3.479
Vitamin A (µg)	500.4 ± 139.9	79.8	757.6 ± 166.8	120.6	19.50***
Thiamin (mg)	0.76 ± 0.15	67.4	0.89 ± 0.16	78.9	10.05***
Riboflavin (mg)	0.80 ± 0.18	62.4	0.93 ± 0.18	73.0	9.352***
Niacin (mg)	12.9 ± 2.7	88.2	15.1 ± 2.7	103.4	9.760***
Vitamin C (mg)	72.4 ± 18.1	72.4	85.0 ± 18.8	85.0	8.346***

<sup>1)</sup> Mean ± SD

<sup>2)</sup> % Estimated Energy Requirements (EER) for energy, % Recommended Intake (RI) for protein, calcium, phosphorus, iron, vitamin A, thiamin, riboflavin, vitamin C and % Adequate Intake (AI) for sodium, potassium

\*  $P < 0.05$ , \*\*\*  $P < 0.005$

## Discussion

This study was designed to measure improvements to nutritional status after the application of a nutritional education program designed for elderly patients in a long-term care hospital. To analyze whether there were improvements in nutritional status, we compared changes in health behaviors, dietary behaviors, Nutrition Screening Initiative scores, and nutrient intakes.

In general, the purpose of nutrition education programs is to provide proper knowledge and information about nutrition-related issues, to draw attention to current dietary issues, to encourage modification of dietary behaviors and to increase participants' ability to practice healthy eating behaviors through a process involving 'knowledge, attitude, and practice' [6].

Although there have been conflicting results regarding the effect of nutritional education, recent studies conducted in Korea have indicated that nutritional education has a positive effect in improving dietary behaviors and nutritional status [6-9,15]. However, there are specific difficulties related to implementing nutritional education for the elderly. It is relatively hard to motivate this group to actively participate in educational programs. Elderly people may find it relatively difficult to understand the contents of the education and may tend to more easily forget what they have learned compared to other age groups [16]. Considering this, the nutritional education program in this study, from planning to implementation, was led by trained dietitians, whose role was to manage elderly patients' diet and who therefore understood their characteristics very well.

The nutritional status of the Korean elderly was reported to be lower than that of other age groups [2]. Among elderly people aged 65 or above, the percentage whose nutrient intake is

insufficient is relatively high (14.6% for one year or older vs. 22.7% for 65 or older), while the rate of elderly people whose energy/fat intake exceeds the recommended level is rather low (4.5% for one year or older vs. 1.3% for 65 or older). In terms of sociodemographic factors, nutritional status is reported to be lower among elderly women than elderly men, and lower among the recipients of government assistance [17], among those who live alone [18-20], among people with a low educational level or low income [21-23], and among those living in a rural area [24]. The result of the Mini Nutritional Assessment (MNA), which was applied to recipients of the long-term care insurance system, including both the elderly people in long-term care facilities and those living at home, showed that those in care facilities had a better nutritional status than those living at home [11]. The studies that compared the nutritional status of low-income, elderly people living at home with that of elderly people in care facilities also reported that those staying in care facilities were less likely to skip meals and ate more diverse foods [25]. The reason for this difference is that those in care facilities are under nutritional care led by dietitians whereas those living at home, especially given the increasing number of nuclear families and single households, are less likely to have access to adequate diet management. However, care facilities may also have difficulties with providing appropriate nutrition management. The cost for nutrition management programs for patients in care facilities is not covered by insurance but by the patients themselves. Therefore, it is hard for them to receive specialized services such as nutritional assessment, nutritional education, individualized counseling or tube feeding. This study therefore tried to measure improvements to nutritional status when nutritional education was added to the existing meal management service for elderly



patients in a long-term care hospital.

The results indicate that the application of a nutritional education program yielded positive changes in elderly patients in terms of their health behaviors, dietary behaviors, nutrient intake, and nutritional risk levels. After the program was completed, the elderly patients' negative perceptions about their health and depression level decreased while positive perceptions increased ( $P < 0.001$ ). Previous studies showed that nutritional intervention may be more effective when participants have high interest in and awareness of their health status [10] and that elderly people who were highly depressed or under severe stress were more likely to be exposed to nutritional risk [26,27]. It is therefore expected that the positive changes in self-rated perceptions of health and depression following nutritional education will also have a positive impact on patients' nutritional status. This study further confirms that, after nutrition education, elderly patients' dietary behavior scores increased significantly ( $P < 0.05$ ). Previous studies that applied nutritional education to healthy old men or elderly people with chronic diseases also reported that, after education, nutritional knowledge scores increased [6-8] and positive changes occurred in terms of dietary behaviors [7,8].

One of the core purposes of nutritional education is to encourage participants to change their behaviors by acquiring proper knowledge about what constitutes a healthy diet and thus to resolve their nutritional problems. The NSI checklist is a tool designed to check nutritional problems over relatively a long period of time, and it generally has limitations in measuring the effect of short-term nutrition education. In this study, improvement was confirmed in three assessment items out of the 10, suggesting changes in behavior could be observed even in a short period of time (I eat fewer than 2 meals per day/ I eat few fruits or vegetable, or milk products/ I eat alone most of the time) ( $P < 0.05$ ). Due to the fact that the subjects were elderly, long-stay patients with chronic diseases, their nutrient intake was much lower than that of healthy counterparts. After nutrition education, however, their overall nutrient intake increased ( $P < 0.001$ ). Some previous studies that applied nutritional education or intervention to elderly people with poor nutritional status also confirmed a positive effect, indicating that after nutritional education or intervention, intake of nutrients or foods was increased [6,8,9], while intake of sodium, which is harmful in excess, was decreased [7]. Even after nutritional education, however, elderly patients' intake of calcium, potassium, and riboflavin remained low in this study. This finding was also reported in other studies on the elderly, which showed that there were little improvements in terms of intake of essential nutrients such as iron, vitamin A, thiamin, and riboflavin [8,9]. These results suggest that in order to address the poor nutritional status of the elderly, a nutritional assistant system is needed, alongside a nutritional education program, so as to increase the intake of nutrients that more likely to be deficient in the diet of elderly populations.

In summary, after a nutritional education program was applied to the elderly, who comprised long-stay patients in a long-term

care hospital, self-rated perceptions on health and on depression improved, positive changes occurred in dietary behaviors and nutritional risk levels, and there was an increase in nutrient intake. These results suggest that efforts designed to improve the nutritional status of the elderly could be more effective when a nutritional assistant program is combined with nutritional education. We hope that the results of this study can be used as preliminary data when establishing guidelines for nutritional management in elderly care facilities, the demand for which is expected to rise in the future.

## References

1. Statistics Korea. Social Indicators of Korea 2010 (the press release). Daejeon: Statistics Korea; 2011. p.5-7.
2. Ministry of Health and Welfare, Korea Centers for Disease Control and Prevention. 2009 National Health Statistics, Korea National Health & Nutrition Survey, the 4th stage, 3rd year. 2009. p.155-8, 448.
3. Korea Institute for Health and Social Affairs. National Nutrition Policy: Its Development Strategies for an Aging Korea). 2007. p.83-91.
4. Duk SW. A study on research trends and development process of health policy for the elderly. *J Korean Gerontol Soc* 2008; 28:773-84.
5. Lee YS, Moon SK. A study on the affecting factors to utilization of long term care hospitals according to the elderly long term care insurance system in Korea. *Korean J Hosp Manage* 2010; 15:49-69.
6. Yoon HJ, Lee SK. Effect of home-visit nutrition education for the elderly with high fasting blood glucose levels. *Korean J Community Nutr* 2006;11:346-60.
7. Yim KS. The effects of a nutrition education program for hypertensive female elderly at the public health center. *Korean J Community Nutr* 2008;13:640-52.
8. Kang HJ, Shin EM, Kim KW. Evaluation of nutrition education for diabetes mellitus management of older adults. *Korean J Community Nutr* 2009;14:734-45.
9. Park PS, Chun BY, Jeong GB, Huh CH, Joo SJ, Park MY. The effect of follow-up nutrition intervention programs applied aged group of high risk undernutrition in rural area (I). *Korean J Food Cult* 2007;22:127-39.
10. Choi Y, Kim C, Park YS. The effect of nutrition education program in physical health, nutritional status and health-related quality of life of the elderly in Seoul. *Korean J Nutr* 2007;40: 270-80.
11. Ministry of Health and Welfare. Ways to Enhance Preventive Functions of Long-Term Care Insurance System for the Elderly-Centered on Nutrition Management. 2011. p.102-9.
12. Ministry of Health and Welfare. Revision of Dietary Guidelines for Koreans (2002: with Action Guides for Adults and the Elderly). 2002. p.191-8.
13. Posner BM, Jette AM, Smith KW, Miller DR. Nutrition and health risks in the elderly: the nutrition screening initiative. *Am J Public Health* 1993;83:972-8.
14. The Korean Nutrition Society. Dietary Reference Intakes for Koreans. 2005.

15. Lee YA, Kim KN, Chang N. The effect of nutrition education on weight control and diet quality in middle-aged women. *Korean J Nutr* 2008;41:54-64.
16. Kim KW. Issues and directions in developing nutrition education for older adults in Korea. *J Community Nutr* 2000;2:71-84.
17. Yim KS, Lee TY. Sociodemographic factors associated with nutrients intake of elderly in Korea. *Korean J Nutr* 2004;37:210-22.
18. Park JK, Son SM. The dietary behaviors, depression rates and nutrient intakes of the elderly females living alone. *Korean J Community Nutr* 2003;8:716-25.
19. Kim HY, Kim MH, Hong SG, Hwang SJ, Park MH. A study on the nutrient intake, health risk factors, blood health status in elderly Korean women living alone. *Korean J Community Nutr* 2005;10:216-23.
20. Kim EM, Song YJ, Kim HS. Evaluation of eating habit and dietary intake by family type of seniors utilizing social welfare center in Seoul. *Korean J Nutr* 2007;40:753-61.
21. Yang EJ, Bang HM. Nutritional status and health risks of low income elderly women in Gwangju area. *Korean J Nutr* 2008;41:65-76.
22. Ro HK, Oh KA. Gender and age differences in the nutritional status of the low income elderly living in Gwangju. *Korean J Community Nutr* 2003;8:302-10.
23. Kwon SO, Oh SY. Associations of household food insecurity with socioeconomic measures, health status and nutrient intake in low income elderly. *Korean J Nutr* 2007;40:762-8.
24. Yoon H, Kwoun J, Lee S. Nutritional status and energy expenditure in the elderly in a rural community. *Korean J Community Nutr* 2002;7:336-44.
25. Kwak KS, Bae YJ, Kim MH. Nutritional status and dietary quality in the low-income elderly residing at home or in health care facilities. *J Korean Diet Assoc* 2008;14:337-50.
26. Son SM, Park JK, Lee HS. Depression and dietary factors related to hyperlipidemia in urban living elderly female from low income group. *Korean J Community Nutr* 2003;8:938-50.
27. Yim KS. Health-related behavioral factors associated with nutritional risks in Korean aged 50 years and over. *Korean J Community Nutr* 2007;12:592-605.