

PROPERTIES OF INDUCED INVERSE POLYNOMIAL MODULES OVER A SUBMONOID

EUNHA CHO AND JINSUN JEONG*

ABSTRACT. Let M be a left R -module and R be a ring with unity, and $S = \{0, 2, 3, 4, \dots\}$ be a submonoid. Then $M[x^{-s}] = \{a_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n} \mid a_i \in M\}$ is an $R[x^s]$ -module. In this paper we show some properties of $M[x^{-s}]$ as an $R[x^s]$ -module.

Let $f : M \rightarrow N$ be an R -linear map and $\overline{M}[x^{-s}] = \{a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n} \mid a_i \in M\}$ and define $N + \overline{M}[x^{-s}] = \{b_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n} \mid b_0 \in N, a_i \in M\}$. Then $N + \overline{M}[x^{-s}]$ is an $R[x^s]$ -module.

We show that given a short exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ of R -modules, $0 \rightarrow L \rightarrow M[x^{-s}] \rightarrow N + \overline{M}[x^{-s}] \rightarrow 0$ is a short exact sequence of $R[x^s]$ -module. Then we show $E_1 + \overline{E_0}[x^{-s}]$ is not an injective left $R[x^s]$ -module, in general.

1. Introduction

Let M be a left R -module and R be a ring with unity and $S = \{0, 2, 3, 4, \dots\}$ be a submonoid. Then $M[x^s] = \{a_0 + a_2x^2 + a_3x^3 + \dots + a_nx^n \mid a_i \in M\}$ is an $R[x^s]$ -module defined by

$$x^k(a_0 + a_2x^2 + a_3x^3 + \dots + a_nx^n) = a_0x^k + a_2x^{2+k} + a_3x^{3+k} + \dots + a_nx^{n+k}$$

for $x^k \in R[x^s]$.

Also $M[x^{-s}] = \{a_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n} \mid a_i \in M\}$ is an $R[x^s]$ -module defined by

$$x^k(a_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}) = a_k + a_{k+2}x^{-2} + a_{k+3}x^{-3} + \dots + a_nx^{-n+k}$$

Received July 2, 2012. Revised August 30, 2012. Accepted September 5, 2012.

2010 Mathematics Subject Classification: 16E30, 13C11, 16D80.

Key words and phrases: injective module, inverse polynomial modules, induced module.

*Corresponding author.

for $x^k \in R[x^s]$. For example,

$$x^3(a_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}) = a_3 + a_5x^{-2} + a_6x^{-3} + \dots + a_nx^{-n+3}.$$

Induced inverse polynomial module was introduced in ([8]). Let M and N be left R -modules and $f : M \rightarrow N$ be an R -linear map. Then an induced polynomial $N + x^{-1}M[x^{-1}]$ is a left $R[x]$ -module defined by

$$x(b_0 + a_1x^{-1} + \dots + a_nx^{-n}) = b_1 + a_2x^{-1} + \dots + a_nx^{-n+1},$$

where $f(a_1) = b_1, b_0 \in N$, and $a_i \in M$. In this paper we generalized induced polynomial modules over a submonoid.

Let $f : M \rightarrow N$ be an R -linear map and let $\overline{M}[x^{-s}] = \{a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n} \mid a_i \in M\}$ and define $N + \overline{M}[x^{-s}] = \{b_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n} \mid b_0 \in N, a_i \in M\}$. Then $N + \overline{M}[x^{-s}]$ is an $R[x^s]$ -module defined by

$$\begin{aligned} x^k(b_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}) \\ = f(a_k) + a_{k+2}x^{-2} + a_{k+3}x^{-3} + \dots + a_nx^{-n+k}. \end{aligned}$$

For example,

$$x^2(b_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}) = f(a_2) + a_4x^{-2} + a_5x^{-3} + \dots + a_nx^{-n+2}.$$

A left R -module E is said to be injective if given any injective linear map $\sigma : M' \rightarrow M$ and any linear map $h : M' \rightarrow E$, there is a linear map $g : M \rightarrow E$ such that $g \circ \sigma = h$. That is,

$$\begin{array}{ccc} 0 & \longrightarrow & M' \xrightarrow{\sigma} M \\ & & \downarrow h \quad \swarrow g \\ & & E \end{array}$$

can always be completed to a commutative diagram([9]).

The map $f : x^2E[x^s] \rightarrow E[x^s]$ defined by $f(ex^2) = e$ is an $R[x^s]$ -linear map. So consider the following diagram

$$\begin{array}{ccc} 0 & \longrightarrow & x^2E[x^s] \longrightarrow E[x^s] \\ & & \downarrow f \\ & & E[x^s]. \end{array}$$

Then we easily see that the above diagram can not be completed for any $R[x^s]$ -linear map. Thus $E[x^s]$ is not an injective left $R[x^s]$ -module with $E \neq 0$.

Northcott([3]) defined inverse polynomial modules and used inverse polynomial modules to study the properties of injective modules and he studied $K[x^{-1}]$ as $K[x]$ -module on field K . And McKerraw([2]) showed that if R is a left noetherian ring and E is an injective left R -module, then $E[x^{-1}]$ is an injective envelope of $M[x^{-1}]$ as $R[x]$ -module. Inverse polynomial modules were studied in ([4]),([5]) and recently in ([1]), ([5]), ([6]), ([7]).

2. inverse polynomial modules over a submonoid

Through this paper we let $S = \{0, 2, 3, 4, \dots\}$ a submonoid.

DEFINITION 2.1. Let M be a left R -module and R be a ring with unity and $S = \{0, 2, 3, 4, \dots\}$ be a submonoid. Then $M[x^{-s}] = \{a_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n} \mid a_i \in M\}$ is an $R[x^s]$ -module defined by

$$r(a_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}) = ra_0 + ra_2x^{-2} + ra_3x^{-3} + \dots + ra_nx^{-n}$$

and

$$x^k(a_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}) = a_k + a_{k+2}x^{-2} + a_{k+3}x^{-3} + \dots + a_nx^{-n+k}$$

where $r \in R$ and $x^k \in R[x^s]$.

PROPOSITION 2.2. Let $\phi : M[x^{-s}] \rightarrow M[x^{-s}]$ be an $R[x^s]$ -linear map. Then $\phi(M) \subset (M + Mx^{-2}) \cap (M + Mx^{-3}) = M$.

Proof. Suppose $m \in M$ and $\phi(m) = a_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}$. Then for $x^2 \in R[x^s]$, $\phi(x^2m) = \phi(0) = 0$, and $x^2\phi(m) = a_2 + a_4x^{-2} + \dots + a_nx^{-n+2} = 0$ implies $a_2 = a_4 = \dots = a_n = 0$. Therefore, $\phi(M) \subset (M + Mx^{-2}) \cap (M + Mx^{-3}) = M$. □

If N is a submodules of M , then we easily see

$$\frac{M[x^{-s}]}{N[x^{-s}]} \cong \frac{M}{N}[x^{-s}]$$

as $R[x^s]$ -module.

PROPOSITION 2.3. *Let M be a left R -module, then*

$$\frac{M[x^{-s}]}{M + Mx^{-3}} \cong M[x^{-s}].$$

Proof. Define $\phi : M[x^{-s}] \rightarrow M[x^{-s}]$ by

$$\begin{aligned} & \phi(a_0 + a_2x^{-2} + a_3x^{-3} + \cdots + a_nx^{-n}) \\ &= \phi(a_0 + a_2x^{-2} + a_3x^{-3} + \cdots + a_nx^{-n}) \\ &= a_2 + a_4x^{-2} + \cdots + a_nx^{-n+2}. \end{aligned}$$

Then we easily see that ϕ is an $R[x^s]$ -linear map.

Let $a_0 + a_2x^{-2} + a_3x^{-3} + \cdots + a_nx^{-n} \in M[x^{-s}]$, then

$$\phi(a_0x^{-2} + a_2x^{-4} + a_3x^{-5} + \cdots + a_nx^{-n-2}) = a_0 + a_2x^{-2} + a_3x^{-3} + \cdots + a_nx^{-n}.$$

Thus ϕ is surjective.

Let $a_0 + a_3x^{-3} \in M[x^{-s}]$, then $\phi(a_0 + a_3x^{-3}) = 0$, so that $M + Mx^{-3} \subset \ker(\phi)$. Let $a_0 + a_2x^{-2} + a_3x^{-3} + \cdots + a_nx^{-n} \in \ker(\phi)$, then

$$\phi(a_0x^{-2} + a_2x^{-4} + a_3x^{-5} + \cdots + a_nx^{-n-2}) = a_2 + a_4x^{-2} + \cdots + a_nx^{-n+2} = 0,$$

so that $a_2 = a_4 = a_5 = \cdots = a_n = 0$, so that $\ker(\phi) \subset M + Mx^{-3}$. Thus $\ker(\phi) = M + Mx^{-3}$. Therefore,

$$\frac{M[x^{-s}]}{M + Mx^{-3}} \cong M[x^{-s}].$$

□

PROPOSITION 2.4. *Let M be a left R -module, then*

$$\sigma : \frac{M[[x^{-s}]]}{M[x^{-s}]} \rightarrow \frac{M[[x^{-s}]]}{M[x^{-s}]}$$

by $\sigma(f + M[x^{-s}]) = x^2(f + M[x^{-s}])$ is an isomorphism.

Proof. Let $f + M[x^{-s}] \in \ker(\sigma)$ and let $f = a_0 + a_2x^{-2} + a_3x^{-3} + \cdots$. Then

$$\begin{aligned} \sigma(f + M[x^{-s}]) &= x^2(f + M[x^{-s}]) \\ &= a_2 + a_4x^{-2} + a_5x^{-3} + \cdots + M[x^{-s}] \\ &= M[x^{-s}]. \end{aligned}$$

So $a_kx^{-k+2} + a_{k+1}x^{-k+1} + \cdots = 0$, $a_k = a_{k+1} = \cdots = 0$, for some k . Thus $f + M[x^{-s}] = M[x^{-s}]$. Therefore, σ is injective.

Let $f + M[x^{-1}] = (a_0 + a_2x^{-2} + a_3x^{-3} + \dots) + M[x^{-s}] \in \frac{M[[x^{-s}]]}{M[x^{-s}]}$. Then there exists $g + M[x^{-s}] = (a_0x^{-2} + a_2x^{-4} + a_3x^{-5} + \dots + a_kx^{-k-2} + \dots) + M[x^{-s}]$ such that $\sigma(g + M[x^{-s}]) = f + M[x^{-s}]$. Therefore, σ is surjective. Hence, σ is an isomorphism. \square

3. Induced polynomial modules over a submonoid

DEFINITION 3.1. Let $f : M \rightarrow N$ be an R -linear map and $S = \{0, 2, 3, 4, \dots\}$ be a submonoid. Then $N + \overline{M}[x^{-s}]$ is an $R[x^s]$ -module, where $\overline{M}[x^{-s}] = \{a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n} \mid a_i \in M\}$ be defined by $r(b_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}) = rb_0 + ra_2x^{-2} + ra_3x^{-3} + \dots + ra_nx^{-n}$ and

$x^k(b_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}) = f(a_k) + a_{k+2}x^{-2} + a_{k+3}x^{-3} + \dots + a_nx^{-n+k}$ for $x^k \in R[x^s], b_0 \in N$. Similarly, we can define

$N + \overline{M}[x^{-s}] = \{b_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n} \mid b_0 \in N, a_i \in M\}$ as a left $R[x^s]$ -module.

THEOREM 3.2. If $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ is a short exact sequence of R -modules, then

$$0 \rightarrow L \rightarrow M[x^{-s}] \rightarrow N + \overline{M}[x^{-s}] \rightarrow 0$$

is a short exact sequence of $R[x^s]$ -module.

Proof. Let $f[x^{-s}] : L \rightarrow M$ be defined by $f[x^{-s}](l) = f(l)$ for $l \in L$. Then easily $f[x^{-s}]$ is an injective $R[x^s]$ -linear map. Let $g[x^{-s}] : M[x^{-s}] \rightarrow N + \overline{M}[x^{-s}]$ be defined by

$$g[x^{-s}](a_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}) = g(a_0) + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}$$

Then easily $g[x^{-s}]$ is an $R[x^s]$ -linear map.

Let $b_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n} \in N + \overline{M}[x^{-s}]$. Then since g is a surjective R -linear map, there exists $a_0 \in M$ such that $g(a_0) = b_0$. Thus $g[x^{-s}](a_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}) = b_0 + a_2x^{-2} + a_3x^{-3} + \dots + a_nx^{-n}$.

So $g[x^{-s}]$ is a surjective $R[x^s]$ -linear map.

Now $(g[x^{-s}] \circ f[x^{-s}])(l) = g[x^{-s}](f(l)) = g(f(l)) = 0$, so that

$$Im(f[x^{-s}]) \subseteq Ker(g[x^{-s}]).$$

And if $a_0 + a_2x^{-2} + a_3x^{-3} + \cdots + a_nx^{-n} \in \text{Ker}(g[x^{-s}])$, then

$$\begin{aligned} g[x^{-s}](a_0 + a_2x^{-2} + a_3x^{-3} + \cdots + a_nx^{-n}) \\ = g(a_0) + a_2x^{-2} + a_3x^{-3} + \cdots + a_nx^{-n} \\ = 0, \end{aligned}$$

implies $g(a_0) = 0$, and $a_2 = a_3 = \cdots = a_n = 0$. Thus $a_0 \in \text{Ker}(g) = \text{Im}(f)$, so that $\text{Ker}(g[x^{-s}]) \subseteq \text{Im}(f[x^{-s}])$. Therefore, $\text{Im}(f[x^{-s}]) \subseteq \text{Ker}(g[x^{-s}])$. Hence,

$$0 \rightarrow L \rightarrow M[x^{-s}] \rightarrow N + \overline{M}[x^{-s}] \rightarrow 0$$

is a short exact sequence of $R[x^{-s}]$ -modules. \square

THEOREM 3.3. *Let $0 \rightarrow N \xrightarrow{f} E_0 \xrightarrow{g} E_1 \rightarrow 0$ be a short exact sequence of R -modules, with $\text{injdim}_R N = 1$, where E_0, E_1 are injective left R -modules. Then $E_1 + \overline{E_0}[x^{-s}]$ is not an injective left $R[x^{-s}]$ -module.*

Proof. Suppose $E_1 + \overline{E_0}[x^{-s}]$ is an injective left $R[x^{-s}]$ -module. Then we can complete the following diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & E_1 & \xrightarrow{i} & E_1 + x^{-2}E_1 \\ & & \downarrow \text{id} & \swarrow \sigma & \\ & & E_1 + \overline{E_0}[x^{-s}] & & \end{array}$$

as commutative diagram by an $R[x^{-s}]$ -linear map σ . Then there exists an R -linear map $h : E_1 \rightarrow E_0$ such that $g \circ h = \text{id}_{E_1}$. This contradicts the fact that the short exact sequence $0 \rightarrow N \xrightarrow{f} E_0 \xrightarrow{g} E_1 \rightarrow 0$ is not split. Hence, $E_1 + \overline{E_0}[x^{-s}]$ is not an injective left $R[x^{-s}]$ -module. \square

DEFINITION 3.4. *Let M be a left R -module and $S = \{0, 2, 3, 4, \dots\}$ be a submonoid, then $M[x^s, x^{-s}] = \{a_0 + a_2x^2 + a_3x^3 + \cdots + a_ix^i + b_2x^{-2} + b_3x^{-3} + \cdots + b_jx^{-j} \mid a_n, b_m \in M\}$ is a left $R[x^s]$ -module be defined by*

$$\begin{aligned} r(a_0 + a_2x^2 + a_3x^3 + \cdots + a_ix^i + b_2x^{-2} + b_3x^{-3} + \cdots + b_jx^{-j}) \\ = ra_0 + ra_2x^2 + ra_3x^3 + \cdots + ra_ix^i + rb_2x^{-2} + rb_3x^{-3} + \cdots + rb_jx^{-j} \end{aligned}$$

and

$$\begin{aligned} & x^k(a_0 + a_2x^2 + a_3x^3 + \cdots + a_ix^i + b_2x^{-2} + b_3x^{-3} + \cdots + b_jx^{-j}) \\ &= a_0x^k + a_2x^{2+k} + \cdots + ra_ix^{i+k} + b_2x^{-2+k} + \cdots + b_jx^{-j+k} \end{aligned}$$

for $x^k \in R[x^s]$.

Similarly we can define $M[[x^s, x^{-s}]]$ as an $R[x^s]$ -module.

THEOREM 3.5. *For any nonzero left R -module E , $E[x^s, x^{-s}]$ is not an injective left $R[x^s]$ -module.*

Proof. Define $f : (1 + x^2) \longrightarrow E[x^s, x^{-s}]$ by $f(1 + x^2) = e$ for $e \in E$ and consider the following diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & (1 + x^2) & \longrightarrow & R[x^s] \\ & & \downarrow f & & \\ & & E[x^s, x^{-s}] & & \end{array}$$

Then we easily see that the above diagram can not be completed for any $R[x^s]$ -linear map. Thus $E[x^s, x^{-s}]$ is not an injective left $R[x^s]$ -module. \square

References

- [1] Z. Liu. *Injectivity of Modules of Generalized Inverse Polynomials*, Comm. Algebra, **29** (2) (2001), 583–592.
- [2] A.S. McKerrow. *On the Injective Dimension of Modules of Power Series*, Quart J. Math., Oxford, **25** (3) (1974), 359–368.
- [3] D.G. Northcott. *Injective Envelopes and Inverse Polynomials*, London Math. Soc. **3** (2) (1974), 290–296.
- [4] S. Park. *The Macaulay-Northcott Functor*, Arch. Math. **63** (1994), 225–230.
- [5] S. Park, *Gorenstein Rings and Inverse Polynomials*, Comm. Algebra, **28** (2) (2000), 785–789.
- [6] S. Park, *The General Structure of Inverse Polynomial Modules*, Czech. Math. J. **51** (126) (2001), 343–349.
- [7] S. Park and E. Cho. *Injective and Projective Properties of $R[x]$ -modules*, Czech. Math. J. **54** (129) (2004), 573–578.
- [8] S. Park, J. Jeong. *Inverse polynomial modules induced by an R -linear map*, Bull. Korean Math. Soc. **47** (2010), 693–669.
- [9] J. Rotman, *An Introduction to Homological Algebra*, Academic Press Inc., New York (1979).

Department of Mathematics
Dong-A University
Pusan 604-714, Korea
E-mail: choeh@donga.ac.kr

Department of Mathematics
Dong-A University
Pusan 604-714, Korea
E-mail: jsjeong@donga.ac.kr