

## **R**, fuzzy **R**, and Algebraic Kripke-style Semantics\* \*\*

Eunsuk Yang

**【Abstract】** This paper deals with Kripke-style semantics for **FR**, a fuzzy version of **R** of Relevance. For this, first, we introduce **FR**, define the corresponding algebraic structures FR-algebras, and give algebraic completeness results for it. We next introduce an algebraic Kripke-style semantics for **FR**, and connect it with algebraic semantics. We furthermore show that such semantics does not work for **R**.

**【Key Words】** Kripke-style semantics, Algebraic semantics, Many-valued logic, Fuzzy logic, **R**, **FR**.

---

\* 접수일자:2012.04.01 심사 및 수정 완료일:2012.05.20 게재확정일:2012.05.29

\*\* This paper was supported by research funds of Chonbuk National University in 2012. I must thank the anonymous referees for their helpful comments.

## 1. Introduction

It is well known that many relevance logicians have had difficulties in providing binary relational Kripke-style semantics, i.e., semantics with binary accessibility relations, for relevance logics (see e.g. [3, 4]). To the best of my knowledge, any satisfactory such semantics for  $\mathbf{R}$  has not yet been introduced. In this paper we show that such semantics can be provided for a fuzzy version of the system  $\mathbf{R}$  of Relevance, although not  $\mathbf{R}$  itself.

Actually, this is a free continuation of the paper [11]. In it the author provided algebraic Kripke-style semantics for Uninorm logic  $\mathbf{UL}$ . Here we introduce algebraic Kripke-style semantics for  $\mathbf{FR}$ , a fuzzy version of  $\mathbf{R}$ .<sup>1)</sup> For this, first, in Section 2 we introduce  $\mathbf{FR}$ , define the corresponding algebraic structures  $\mathbf{FR}$ -algebras, and give algebraic completeness results for it. In Section 3 we introduce an algebraic Kripke-style semantics for  $\mathbf{FR}$ , and connect them with algebraic semantics. We furthermore show that this semantics does not work for  $\mathbf{R}$  (see Example 3.9).

For convenience, we shall adopt the notation and terminology similar to those in [5, 7, 8, 10], and assume familiarity with them (together with the results found in them).

## 2. The logic $\mathbf{FR}$ and its algebraic semantics

We base  $\mathbf{FR}$  on a countable propositional language with

---

<sup>1)</sup> To see why algebraic Kripke-style semantics are interesting, see [12].

formulas  $FOR$  built inductively as usual from a set of propositional variables  $VAR$ , binary connectives  $\rightarrow$ ,  $\&$ ,  $\wedge$ ,  $\vee$ , and constants  $\mathbf{f}$ ,  $\mathbf{t}$ , with defined connectives:<sup>2)</sup>

$$\text{df1. } \sim\phi := \phi \rightarrow \mathbf{f}$$

$$\text{df2. } \phi \leftrightarrow \psi := (\phi \rightarrow \psi) \wedge (\psi \rightarrow \phi).$$

We moreover define  $\phi_{\mathbf{t}} := \phi \wedge \mathbf{t}$ . For the remainder we shall follow the customary notation and terminology. We use the axiom systems to provide a consequence relation.

We start with the following axiomatization of **FR**.

**Definition 2.1** **FR** consists of the following axiom schemes and rules:<sup>3)</sup>

$$\text{A1. } \phi \rightarrow \phi \quad (\text{self-implication, SI})$$

$$\text{A2. } (\phi \wedge \psi) \rightarrow \phi, (\phi \wedge \psi) \rightarrow \psi \quad (\wedge\text{-elimination, } \wedge\text{-E})$$

$$\text{A3. } ((\phi \rightarrow \psi) \wedge (\phi \rightarrow \chi)) \rightarrow (\phi \rightarrow (\psi \wedge \chi)) \quad (\wedge\text{-introduction, } \wedge\text{-I})$$

$$\text{A4. } \phi \rightarrow (\phi \vee \psi), \psi \rightarrow (\phi \vee \psi) \quad (\vee\text{-introduction, } \vee\text{-I})$$

$$\text{A5. } ((\phi \rightarrow \chi) \wedge (\psi \rightarrow \chi)) \rightarrow ((\phi \vee \psi) \rightarrow \chi) \quad (\vee\text{-elimination, } \vee\text{-E})$$

$$\text{A6. } (\phi \wedge (\psi \vee \chi)) \rightarrow ((\phi \wedge \psi) \vee (\phi \wedge \chi)) \quad (\wedge \vee\text{-distributivity, } \wedge \vee\text{-D})$$

$$\text{A7. } (\phi \& \psi) \rightarrow (\psi \& \phi) \quad (\&\text{-commutativity, } \&\text{-C})$$

$$\text{A8. } (\phi \& \mathbf{t}) \leftrightarrow \phi \quad (\text{push and pop, PP})$$

---

<sup>2)</sup> Note that while  $\wedge$  is the extensional conjunction connective,  $\&$  is the intensional conjunction one.

<sup>3)</sup> A6, indeed, is redundant in **FR**. But we introduce this in order to show that **R** is the **FR** omitting A13. Note that the system omitting both A6 and A13 is not **R** (cf see [1, 2, 4]).

- A9.  $(\phi \rightarrow \psi) \rightarrow ((\psi \rightarrow \chi) \rightarrow (\phi \rightarrow \chi))$  (suffixing, SF)  
 A10.  $(\phi \rightarrow (\psi \rightarrow \chi)) \leftrightarrow ((\phi \& \psi) \rightarrow \chi)$  (residuation, RE)  
 A11.  $\phi \rightarrow (\phi \& \phi)$  (contraction, CR)  
 A12.  $\sim\sim\phi \rightarrow \phi$  (double negation elimination, DNE)  
 A13.  $(\phi \rightarrow \psi)_t \vee (\psi \rightarrow \phi)_t$  (t-prelinearity, PL<sub>t</sub>)  
 $\phi \rightarrow \psi, \phi \vdash \psi$  (modus ponens, mp)  
 $\phi, \psi \vdash \phi \wedge \psi$  (adjunction, adj).

A13 is the axiom scheme for linearity, and logics being complete w.r.t. linearly ordered (corresponding) algebras are said to be fuzzy logics (see e.g. [3]).

Note that the system **R** is the **FR** omitting A13. Note also that in **R** (and so **FR**),  $\phi \rightarrow \psi$  can be defined as  $\sim(\phi \& \sim\psi)$  (df3), and  $\phi \& \psi$  as  $\sim(\phi \rightarrow \sim\psi)$  (df4).

**Proposition 2.2** **FR** proves:

- (1)  $(\phi \& (\psi \& \chi)) \leftrightarrow ((\phi \& \psi) \& \chi)$  (&-associativity, AS)
- (2)  $(\phi \wedge \psi) \rightarrow (\phi \& \psi)$
- (3)  $(\phi \& (\psi \wedge \chi)) \leftrightarrow ((\phi \& \psi) \wedge (\phi \& \chi))$
- (4)  $(\phi \rightarrow (\psi \vee \chi)) \leftrightarrow ((\phi \rightarrow \psi) \vee (\phi \rightarrow \chi))$
- (5)  $((\phi \rightarrow (\psi \vee \chi)) \wedge (\psi \rightarrow \chi)) \rightarrow (\phi \rightarrow \chi)$ .

**Proof:** The proof for (1) to (3) is easy, just noting that in order to prove (3) we need A13 (cf. see [1]). We prove (4) and (5).

For the proof of (4), first note that in **R**, we can easily prove  $(\phi \rightarrow (\psi \vee \chi)) \leftrightarrow (\phi \rightarrow \sim(\sim\psi \wedge \sim\chi))$  and  $(\phi \rightarrow \sim(\sim\psi$

$\wedge \sim \chi)) \leftrightarrow ((\phi \ \& \ (\sim \psi \ \wedge \ \sim \chi)) \rightarrow \mathbf{f})$ . Then, using (3), we can prove  $((\phi \ \& \ (\sim \psi \ \wedge \ \sim \chi)) \rightarrow \mathbf{f}) \leftrightarrow \sim((\phi \ \& \ \sim \psi) \ \wedge \ (\phi \ \& \ \sim \chi))$ , and using de Morgan laws, we get  $\sim((\phi \ \& \ \sim \psi) \ \wedge \ (\phi \ \& \ \sim \chi)) \leftrightarrow \sim(\phi \ \& \ \sim \psi) \ \vee \ \sim(\phi \ \& \ \sim \chi)$ . Hence, by df3, we obtain  $(\phi \rightarrow (\psi \vee \chi)) \leftrightarrow ((\phi \rightarrow \psi) \vee (\phi \rightarrow \chi))$ , as required.

For the proof of (5), first note that in **R**, we can easily prove  $((\phi \rightarrow \psi) \wedge (\psi \rightarrow \chi)) \rightarrow (\phi \rightarrow \chi)$  using (2). Then, since  $((\phi \rightarrow \chi) \wedge (\psi \rightarrow \chi)) \rightarrow (\phi \rightarrow \chi)$ , we can obtain  $((\phi \rightarrow \psi) \wedge (\psi \rightarrow \chi)) \vee ((\phi \rightarrow \chi) \wedge (\psi \rightarrow \chi)) \rightarrow (\phi \rightarrow \chi)$ . Thus, using A6, we get  $((\psi \rightarrow \chi) \wedge ((\phi \rightarrow \psi) \vee (\phi \rightarrow \chi))) \rightarrow (\phi \rightarrow \chi)$ . Hence, using (4), we can obtain that  $((\phi \rightarrow (\psi \vee \chi)) \wedge (\psi \rightarrow \chi)) \rightarrow (\phi \rightarrow \chi)$ , as wished.  $\square$

Note that **R** does not prove (5) in Proposition 2.2 (see [5]).

In **FR**, **f** can be defined as  $\sim \mathbf{t}$ . A *theory* over **FR** is a set **T** of formulas. A *proof* in a theory **T** over **FR** is a sequence of formulas whose each member is either an axiom of **FR** or a member of **T** or follows from some preceding members of the sequence using the two rules in Definition 2.1.  $\mathbf{T} \vdash \phi$ , more exactly  $\mathbf{T} \vdash_{\mathbf{FR}} \phi$ , means that  $\phi$  is *provable* in **T** w.r.t. **FR**, i.e., there is a **FR**-proof of  $\phi$  in **T**. The relevant deduction theorem (RDT<sub>t</sub>) for **FR** is as follows:

**Proposition 2.3** ([7]) Let **T** be a theory, and  $\phi, \psi$  formulas.

(RDT<sub>t</sub>)  $\mathbf{T} \cup \{\phi\} \vdash \psi$  iff  $\mathbf{T} \vdash \phi_t \rightarrow \psi$ .

For convenience, “ $\sim$ ”, “ $\wedge$ ”, “ $\vee$ ”, and “ $\rightarrow$ ” are used ambiguously as propositional connectives and as algebraic operators, but context should make their meaning clear.

The algebraic counterpart of **FR** is the class of *FR-algebras*. Let  $x_t := x \wedge t$ . They are defined as follows.

**Definition 2.4** (i) A *pointed commutative residuated distributive lattice* is a structure  $\mathbf{A} = (A, t, f, \wedge, \vee, *, \rightarrow)$  such that:

(I)  $(A, \wedge, \vee)$  is a distributive lattice.

(II)  $(A, *, t)$  is a commutative monoid.

(III)  $y \leq x \rightarrow z$  iff  $x * y \leq z$ , for all  $x, y, z \in A$  (residuation).

(ii) (Dunn-algebras, [1, 2]) A *Dunn-algebra* is a pointed commutative residuated distributive lattice satisfying:

(IV)  $x \leq x * x$  (contraction).

(V)  $(x \rightarrow f) \rightarrow f \leq x$  (double negation elimination).

(iii) (FR-algebras) A *FR-algebra* is a Dunn-algebra satisfying:

(VI)  $t \leq (x \rightarrow y)_t \vee (y \rightarrow x)_t$  (pl).

Note that the class of Dunn-algebras characterizes the system **R**. Note also that Dunn-algebras are also called De Morgan monoids.

Additional (unary) negation and (binary) equivalence operations are defined as in Section 2.1:  $\sim x := x \rightarrow f$  and  $x \leftrightarrow y := (x \rightarrow y) \wedge (y \rightarrow x)$ .

The class of all FR-algebras is a variety which will be denoted by **FR**.

FR-algebra is said to be *linearly ordered* if the ordering of its algebra is linear, i.e.,  $x \leq y$  or  $y \leq x$  (equivalently,  $x \wedge y = x$  or  $x \wedge y = y$ ) for each pair  $x, y$ .

**Definition 2.5** (Evaluation) Let  $\mathcal{A}$  be an algebra. An  $\mathcal{A}$ -evaluation is a function  $v : \text{FOR} \rightarrow \mathcal{A}$  satisfying:  $v(\phi \rightarrow \psi) = v(\phi) \rightarrow v(\psi)$ ,  $v(\phi \wedge \psi) = v(\phi) \wedge v(\psi)$ ,  $v(\phi \vee \psi) = v(\phi) \vee v(\psi)$ ,  $v(\phi \& \psi) = v(\phi) * v(\psi)$ ,  $v(\mathbf{f}) = \mathbf{f}$ , (and hence  $v(\sim\phi) = \sim v(\phi)$  and  $v(\mathbf{t}) = \mathbf{t}$ ).

**Definition 2.6** Let  $\mathcal{A}$  be a FR-algebra,  $T$  a theory,  $\phi$  a formula, and  $\mathbf{K}$  a class of FR-algebras.

(i) (Tautology)  $\phi$  is a *t-tautology* in  $\mathcal{A}$ , briefly an  $\mathcal{A}$ -tautology (or  $\mathcal{A}$ -valid), if  $v(\phi) \geq \mathbf{t}$  for each  $\mathcal{A}$ -evaluation  $v$ .

(ii) (Model) An  $\mathcal{A}$ -evaluation  $v$  is an  $\mathcal{A}$ -model of  $T$  if  $v(\phi) \geq \mathbf{t}$  for each  $\phi \in T$ . By  $\text{Mod}(T, \mathcal{A})$ , we denote the class of  $\mathcal{A}$ -models of  $T$ .

(iii) (Semantic consequence)  $\phi$  is a *semantic consequence* of  $T$  w.r.t.  $\mathbf{K}$ , denoting by  $T \models_{\mathbf{K}} \phi$ , if  $\text{Mod}(T, \mathcal{A}) = \text{Mod}(T \cup \{\phi\}, \mathcal{A})$  for each  $\mathcal{A} \in \mathbf{K}$ .

**Definition 2.7** (FR-algebra) Let  $\mathcal{A}$ ,  $T$ , and  $\phi$  be as in Definition 2.6.  $\mathcal{A}$  is a *FR-algebra* iff whenever  $\phi$  is FR-provable in  $T$  (i.e.  $T \vdash_{\text{FR}} \phi$ ), it is a semantic consequence of  $T$  w.r.t. the set  $\{\mathcal{A}\}$  (i.e.  $T \models_{\{\mathcal{A}\}} \phi$ ),  $\mathcal{A}$  a FR-algebra. By  $\text{MOD}^{(l)}(\text{FR})$ , we denote the class of (linearly ordered) **FR**-algebras. Finally, we write  $T \models_{\text{FR}}^{(l)} \phi$  in place of  $T \models_{\text{MOD}^{(l)}(\text{FR})} \phi$ .

Note that since each condition for the FR-algebra has a form of equation or can be defined in equation (exercise), it can be ensured that the class of all FR-algebras is a variety.

We first show that classes of provably equivalent formulas form a FR-algebra. Let  $T$  be a fixed theory over  $\mathbf{FR}$ . For each formula  $\phi$ , let  $[\phi]_T$  be the set of all formulas  $\psi$  such that  $T \vdash_{\mathbf{FR}} \phi \leftrightarrow \psi$  (formulas  $T$ -provably equivalent to  $\phi$ ).  $A_T$  is the set of all the classes  $[\phi]_T$ . We define that  $[\phi]_T \rightarrow [\psi]_T = [\phi \rightarrow \psi]_T$ ,  $[\phi]_T * [\psi]_T = [\phi \& \psi]_T$ ,  $[\phi]_T \wedge [\psi]_T = [\phi \wedge \psi]_T$ ,  $[\phi]_T \vee [\psi]_T = [\phi \vee \psi]_T$ ,  $t = [t]_T$ , and  $\perp_t = [f]_T$ . By  $A_T$ , we denote this algebra.

**Proposition 2.8** For  $T$  a theory over  $\mathbf{FR}$ ,  $A_T$  is a  $\mathbf{FR}$ -algebra.

**Proof:** Note that A1 to A6 ensure that  $\wedge$  and  $\vee$  satisfy (I) in Definition 2.4; that A7, A8, and AS ensure that  $\&$  satisfies (II); that A10 ensures that (III) holds; and that A11, A12, and A13 ensure that (IV), (V), and (VI), respectively, hold. It is obvious that  $[\phi]_T \leq [\psi]_T$  iff  $T \vdash_{\mathbf{FR}} \phi \leftrightarrow (\phi \wedge \psi)$  iff  $T \vdash_{\mathbf{FR}} \phi \rightarrow \psi$ . Finally recall that  $A_T$  is a  $\mathbf{FR}$ -algebra iff  $T \vdash_{\mathbf{FR}} \psi$  implies  $T \models_{\mathbf{FR}} \psi$ , and observe that for  $\phi$  in  $T$ , since  $T \vdash_{\mathbf{FR}} t \rightarrow \phi$ , it follows that  $[t]_T \leq [\phi]_T$ . Thus it is a  $\mathbf{FR}$ -algebra.  $\square$

We next note that the nomenclature of the prelinearity condition is explained by the subdirect representation theorem below.

**Proposition 2.9** (Cf. [10]) Each FR-algebra is a subdirect

product of linearly ordered FR-algebras.

**Theorem 2.10** (Strong completeness) Let  $T$  be a theory, and  $\phi$  a formula.  $T \vdash_{FR} \phi$  iff  $T \models_{FR} \phi$  iff  $T \models_{FR}^1 \phi$ .

**Proof:** (i)  $T \vdash_{FR} \phi$  iff  $T \models_{FR} \phi$ . The left-to-right direction follows from definition. The right-to-left direction is as follows: from Proposition 2.8, we obtain  $\mathbf{A}_T \in \text{MOD}(FR)$ , and for  $\mathbf{A}_T$ -evaluation  $v$  defined as  $v(\psi) = [\psi]_T$ , it holds that  $v \in \text{Mod}(T, \mathbf{A}_T)$ . Thus, since from  $T \models_{FR} \phi$  we obtain that  $[\phi]_T = v(\phi) \geq t$ ,  $T \vdash_{FR} t \rightarrow \phi$ . Then, since  $T \vdash_{FR} t$ , by (mp)  $T \vdash_{FR} \phi$ , as required.

(ii)  $T \models_{FR} \phi$  iff  $T \models_{FR}^1 \phi$ . It follows from Proposition 2.9.  $\square$

### 3. Kripke-style semantics for FR

Here we consider algebraic Kripke-style semantics for **FR**.

**Definition 3.1** (Algebraic Kripke frame) An *algebraic Kripke frame* is a structure  $\mathbf{X} = (X, t, f, \leq, *, \rightarrow)$  such that  $(X, t, f, \leq, *, \rightarrow)$  is a linearly ordered residuated pointed commutative monoid. The elements of  $\mathbf{X}$  are called *nodes*.

**Definition 3.2** (FR frame) A *FR frame* is an algebraic Kripke frame, where  $x = (x \rightarrow f) \rightarrow f$ , and  $*$  is contractive, i.e.,  $x \leq x * x$ .

An *evaluation* or *forcing* on an algebraic Kripke frame is a relation  $\Vdash$  between nodes and propositional variables, and arbitrary formulas subject to the conditions below: for every propositional variable  $p$ ,

(AHC) if  $x \Vdash p$  and  $y \leq x$ , then  $y \Vdash p$ ; and

for arbitrary formulas,

(t)  $x \Vdash t$  iff  $x \leq t$ ;

(f)  $x \Vdash f$  iff  $x \leq f$ ;

( $\wedge$ )  $x \Vdash \phi \wedge \psi$  iff  $x \Vdash \phi$  and  $x \Vdash \psi$ ;

( $\vee$ )  $x \Vdash \phi \vee \psi$  iff  $x \Vdash \phi$  or  $x \Vdash \psi$ ;

( $\&$ )  $x \Vdash \phi \& \psi$  iff there are  $y, z \in X$  such that  $y \Vdash \phi$ ,  $z \Vdash \psi$ , and  $x \leq y * z$ ;

( $\rightarrow$ )  $x \Vdash \phi \rightarrow \psi$  iff for all  $y \in X$ , if  $y \Vdash \phi$ , then  $x * y \Vdash \psi$ .

**Definition 3.3** (i) (Algebraic Kripke model) An *algebraic Kripke model* is a pair  $(X, \Vdash)$ , where  $X$  is an algebraic Kripke frame and  $\Vdash$  is a forcing on  $X$ .

(ii) (FR model) A *FR model* is a pair  $(X, \Vdash)$ , where  $X$  is a FR frame and  $\Vdash$  is a forcing on  $X$ .

**Definition 3.4** (Cf. [9]) Given an algebraic Kripke model  $(X, \Vdash)$ , a node  $x$  of  $X$  and a formula  $\phi$ , we say that  $x$  *forces*  $\phi$  to express  $x \Vdash \phi$ . We say that  $\phi$  is *true* in  $(X, \Vdash)$  if  $t \Vdash \phi$ , and

that  $\phi$  is *valid* in the frame  $\mathbf{X}$  (expressed by  $\mathbf{X}$  models  $\phi$ ) if  $\phi$  is true in  $(\mathbf{X}, \Vdash)$  for every forcing  $\Vdash$  on  $\mathbf{X}$ .

For soundness and completeness for **FR**, let  $\vdash_{\text{FR}} \phi$  be the theoremhood of  $\phi$  in **FR**. First we note the following lemma.

**Lemma 3.5** (Hereditary Lemma, HL) Let  $\mathbf{X}$  be an algebraic Kripke frame. For any sentence  $\phi$  and for all nodes  $x, y \in \mathbf{X}$ , if  $x \Vdash \phi$  and  $y \leq x$ , then  $y \Vdash \phi$ .

**Proof:** Easy.  $\square$

**Proposition 3.6** (Soundness) If  $\vdash_{\text{FR}} \phi$ , then  $\phi$  is valid in every FR frame.

**Proof:** We prove the validity of A11 as an example: it suffices to show that if  $x \Vdash \phi$ , then  $x \Vdash \phi \ \& \ \phi$ . Assume  $x \Vdash \phi$ . Then, since  $x \leq x * x$ , using ( $\&$ ), we can obtain  $x \Vdash \phi \ \& \ \phi$ , as required.

The proof for the other cases is left to the interested reader.  $\square$

By a *chain*, we mean a linearly ordered algebra. The next proposition connects algebraic Kripke semantics and algebraic semantics for **FR** (cf. see [9]).

**Proposition 3.7** (i) The  $\{t, f, \leq, *, \rightarrow\}$  reduct of a FR-chain  $\mathbf{A}$  is a FR frame.

(ii) Let  $\mathbf{X} = (X, t, f, \leq, *, \rightarrow)$  be a FR frame. Then the structure  $\mathbf{A} = (X, t, f, \max, \min, *, \rightarrow)$  is a FR-algebra (where *max* and *min* are meant w.r.t.  $\leq$ ).

(iii) Let  $\mathbf{X}$  be the  $\{t, f, \leq, *, \rightarrow\}$  reduct of a FR-chain  $\mathbf{A}$ , and let  $v$  be an evaluation in  $\mathbf{A}$ . Let for every atomic formula  $p$  and for every  $x \in \mathbf{A}$ ,  $x \Vdash p$  iff  $x \leq v(p)$ . Then  $(\mathbf{X}, \Vdash)$  is a FR model, and for every formula  $\phi$  and for every  $x \in \mathbf{A}$ , we obtain that:  $x \Vdash \phi$  iff  $x \leq v(\phi)$ .

**Proof:** The proof for (i) and (ii) is easy. For the proof of (iii), see Proposition 3.8 in [10].  $\square$

**Theorem 3.8** (Strong completeness) **FR** is strongly complete w.r.t. the class of all FR-frames.

**Proof:** It follows from Proposition 3.7 and Theorem 2.10.  $\square$

Let an *R frame*  $\mathbf{X}$  be an FR frame on a partially ordered monoid in place of a linearly ordered monoid, let an evaluation or forcing  $\Vdash$  on an R frame be the same as that on a FR frame, and let  $(\mathbf{X}, \Vdash)$  be an R model. Then, at first glance,  $(\mathbf{X}, \Vdash)$  seems to be a model for **R**. But actually it is not. The following example verifies it.

**Example 3.9** An R model  $(\mathbf{X}, \Vdash)$  validates Proposition 2.2 (5), i.e.,  $t \Vdash ((\phi \rightarrow (\psi \vee \chi)) \wedge (\psi \rightarrow \chi)) \rightarrow (\phi \rightarrow \chi)$ .

**Proof:** By  $(\rightarrow)$  and  $(\wedge)$ , we assume  $x \vdash (\phi \rightarrow (\psi \vee \chi))$  and  $x \vdash \psi \rightarrow \chi$ , and show  $x \vdash \phi \rightarrow \chi$ . For this last, we further assume  $y \vdash \phi$  and show  $x * y \vdash \chi$ . By the suppositions and  $(\rightarrow)$ , we have  $x * y \vdash \psi \vee \chi$ , therefore  $x * y \vdash \psi$  or  $x * y \vdash \chi$  by  $(\vee)$ . Let  $x * y \vdash \psi$ . Then, since  $x \vdash \psi \rightarrow \chi$ , by  $(\rightarrow)$  we obtain  $x * (x * y) \vdash \chi$ , therefore  $(x * x) * y \vdash \chi$  by the associativity of  $*$ . Then, since  $x \leq x * x$ , using Lemma 3.5, we get  $x * y \vdash \chi$ .  $\square$

This sentence is not a theorem of **R** but a theorem of **FR**. Thus this model is not for **R**.

#### 4. Concluding remark

We investigated algebraic Kripke-style semantics for **FR**, a fuzzy version of **R**. We proved soundness and completeness theorems. But we did not provide algebraic Kripke-style semantics for **R**. This is an open problem left in this paper.

## References

- [1] Anderson, A. R., and Belnap, N. D. (1975), *Entailment: The Logic of Relevance and Necessity*, vol. 1, Princeton, Princeton Univ. Press.
- [2] Anderson, A. R., Belnap, N. D., and Dunn, J. M. (1992), *Entailment: The Logic of Relevance and Necessity*, vol. 2, Princeton, Princeton Univ. Press.
- [3] Cintula, P. (2006), “Weakly Implicative (Fuzzy) Logics I: Basic properties”, *Archive for Mathematical Logic*, pp. 673-704.
- [4] Dunn, J. M. (1976), “A Kripke-style semantics for R-Mingle using a binary accessibility relation”, *Studia Logica*, 35, pp. 163-172.
- [5] Dunn, J. M. (1986), “Relevance logic and entailment”, *Handbook of Philosophical Logic*, vol. III, D. Gabbay and F. Guenther (eds.), Dordrecht, D. Reidel Publ. Co., pp. 117-224.
- [6] Dunn, J. M. (2000), “Partiality and its Dual”, *Studia Logica*, 66, pp. 5-40.
- [7] Meyer, R. K., Dunn, J. M., and Leblanc, H. (1976), “Completeness of relevant quantification theories”, *Notre Dame Journal of Formal Logic*, 15, pp. 97-121.
- [8] Montagna, F. and Sacchetti, L. (2003), “Kripke-style semantics for many-valued logics”, *Mathematical Logic Quarterly*, 49, pp. 629-641.
- [9] Montagna, F. and Sacchetti, L. (2004), “Corrigendum to

- “Kripke-style semantics for many-valued logics”, *Mathematical Logic Quarterly*, 50, pp. 104-107.
- [10] Tsinakis, C., and Blount, K. (2003), “The structure of residuated lattices”, *International Journal of Algebra and Computation*, 13, pp. 437-461.
- [11] Yang, E. (2012), “Kripke-style semantics for UL”, *Korean Journal of Logic*, 15/1, pp. 1-15.
- [12] Yang, E., “Algebraic Kripke-style semantics for relevance logics”, manuscript.

전북대학교 철학과

Department of Philosophy, Chonbuk National University  
eunsyang@jbnu.ac.kr

---

## R, fuzzy R, and Algebraic Kripke-style Semantics

양 은 석

---

이 글에서 우리는 연관 논리 **R**을 퍼지화한 체계 **FR**을 위한 크립키형 의미론을 다룬다. 이를 위하여 먼저 **FR** 체계를 소개하고 그에 상응하는 FR-대수를 정의한 후 **FR**이 대수적으로 완전하다는 것을 보인다. 다음으로 **FR**을 위한 대수적 크립키형 의미론을 소개하고 이를 대수적 의미론과 연관 짓는다. 마지막으로 이러한 의미론이 **R**에는 적용될 수 없다는 점을 보인다.

주요어: **R**, **FR**, (대수적) 크립키형 의미론, 대수적 의미론, 다치 논리, 퍼지 논리