

Effects of Rutaecarpine on Hydrogen Peroxide-Induced Apoptosis in Murine Hepa-1c1c7 Cells

Sung-Jin Lee¹, Hyunjin Ahn², Kung-Woo Nam³, Kyeong Ho Kim⁴ and Woongchon Mar^{2,*}

¹Department of Animal Biotechnology, Kangwon National University, Chuncheon 200-701,

²Natural Products Research Institute, College of Pharmacy, Seoul National University, Seoul 151-951,

³SCH-BioPharm Human Resources Development Center, Soonchunhyang University, Asan 336-745,

⁴College of Pharmacy, Kangwon National University, Chuncheon 200-701, Republic of Korea

Abstract

The aim of this study was to investigate the inhibitory effects of rutaecarpine on DNA strand breaks and apoptosis induced by hydrogen peroxide (H₂O₂) in murine Hepa-1c1c7 cells. Oxidative DNA damage was estimated by nuclear condensation assessment, fluorescence-activated cell sorting analysis, and Comet assay. Rutaecarpine inhibited cell death induced by 500 μM H₂O₂, as assessed by 4',6-diamidino-2-phenylindole (DAPI) staining. Treatment with rutaecarpine reduced the number of DNA strand breaks induced by H₂O₂, as assessed by DAPI staining and Comet assay, and increased quinone reductase, phosphatidylinositol 3-kinase, and pAkt protein levels, as assessed by western blotting.

Key Words: Rutaecarpine, Oxidative stress, DNA damage, Quinone reductase, Apoptosis, Hydrogen peroxide

INTRODUCTION

Mammalian cells have evolved numerous complicated mechanisms that protect against toxic insults. Phase II enzymes are the primary lines of defense against reactive chemical species generated by exogenous quinines and related compounds (Talalay *et al.*, 1995). Recent studies have shown that the antioxidant response element (ARE) in the promoters of genes encoding many of these enzymes may allow transcriptional control of various enzymes in response to dietary activation factors (Finley, 2003). In addition to detoxification reactions, phase II proteins may help to maintain the overall antioxidant functions of the cell by maintaining vitamin E and coenzyme Q in their reduced and active state and eliminating reactive oxygen species (ROS) that can potentially cause DNA damage (Loft and Poulsen, 1996).

Oxidative stress is believed to play important roles in many human diseases. Modification of biological molecules, such as nucleic acids, lipids, and proteins, by oxygen free radicals is involved in the pathogenesis of many diseases (Finkle and Holbrook, 2000). Cells are subjected to oxidative stress when levels of ROS exceed the cells' counter-regulatory antioxidant capacity (Martindale and Holbrook, 2002). Such stress can trigger the activation of certain signaling pathways that protect

against cell death (Ahlgren-Bechendorf *et al.*, 1999). Hydrogen peroxide (H₂O₂) treatment induces the phosphorylation of members of the mitogen-activated protein kinase (MAPK) family [p38 MAPK, c-Jun NH₂-terminal kinases (JNK) 1 and 2, and extracellular signal-regulated protein kinases (ERK) 1 and 2] and phosphatidylinositol 3-kinase (PI3K). The activation of ERK1/2 and PI3K is generally accepted to promote cell survival by activating anti-apoptotic signaling pathways, whereas the activation of JNK and p38 MAPK is associated with cell death (Burdette *et al.*, 2002; Traister *et al.*, 2005; Deguil *et al.*, 2007). MAPK cascades are modulated by reactive intermediates including ROS, which may be critical in the regulation of phase II genes during cellular homeostasis through ARE/electrophile response element-containing activator protein-1 DNA enhancer elements (Ross *et al.*, 2000).

In this study, rutaecarpine was isolated and purified from *Evodia rutaecarpa*, as described in a previous report (Ahn *et al.*, 2008). The aim of this study was to investigate the effect of rutaecarpine with respect to the induction of phase II enzymes, as assessed by measuring quinone reductase (QR) activity and resistance to H₂O₂-induced DNA damage in Hepa-1c1c7 cells.

www.biomolther.org

Open Access <http://dx.doi.org/10.4062/biomolther.2012.20.5.487>

pISSN: 1976-9148 eISSN: 2005-4483

Copyright © 2012 The Korean Society of Applied Pharmacology

Received Oct 28, 2011 Revised Jul 24, 2012 Accepted Jul 30, 2012

*Corresponding Author

E-mail: mars@snu.ac.kr

Tel: +82-2-880-2473, Fax: +82-2-3672-5488

The first two authors contributed equally to this work.

MATERIALS AND METHODS

Cell culture

Mouse hepatoma Hepa-1c1c7 cells were cultured in minimum essential medium (Eagle), alpha modification (Sigma, St. Louis, MO, USA) that did not contain nucleosides but was supplemented with 2 mmol/L glutamine and 10 ml/L antibiotics (Sigma).

DAPI staining

After treatment with rutaecarpine and H₂O₂, Hepa-1c1c7 cells were washed with cold phosphate-buffered saline (PBS) and fixed through incubation with 4% formaldehyde for 10 min at room temperature. Following incubation with Triton X-100 (0.2% in PBS) for 5 min, Hepa-1c1c7 cells were incubated with 4',6-diamidino-2-phenylindole (DAPI, 1 μg/ml; Sigma) for 15 min at room temperature. Hepa-1c1c7 cells were viewed under a fluorescence microscope (Deguil *et al.*, 2007).

Fluorescence-activated cell sorting analysis

Hepa-1c1c7 cells were collected after rutaecarpine and H₂O₂ treatment, centrifuged, resuspended in PBS at a concentration of 25×10⁴ cells/ml, and then incubated with 70% ethyl alcohol for 1 h at -20°C. Propidium iodide (PI) was added at a concentration of 50 mg/ml and the cells were kept on ice until analysis (Traister *et al.*, 2005). Apoptosis was analyzed using an EPICS XL fluorescence-activated cell sorter (Coulter, Fullerton, CA, USA).

Comet assay

A single-cell gel electrophoresis assay was carried out according to the manufacturer's suggested procedure (Trevigen, Gaithersburg, MD, USA) with minor modifications (Burdette *et al.*, 2002). Hepa-1c1c7 cells were plated at a density of 2×10⁴ cells/ml and grown for 24 h prior to treatment with sample. After the sample had been added, the cells were incubated for 48 h to induce QR and then treated with H₂O₂ for 6 h.

Cells were washed with PBS, harvested, combined with molten low-melting agarose at a density of 1×10⁵ cells/ml, and maintained at 42°C at a ratio of 1:10 (v/v). The agarose-cell mixture (50 μl) was immediately pipetted onto Comet slides

(Trevigen). The slides were then immersed in pre-chilled lysis solution for 30 min at 4°C in the dark. After lysis, horizontal electrophoresis was performed for 30 min at 300 mA. Slides were fixed through incubation with 70% ethanol for 5 min, dried, stained with 50 μl SYBR Green solution, and viewed under a fluorescence microscope.

RNA extraction and reverse-transcriptase polymerase chain reaction

Total RNA was extracted from hepatic constituent cells and liver tissues using Easy Blue (Intron Biotechnology, Seong-




Fig. 1. Effects of rutaecarpine on Hepa-1c1c7 cell apoptosis (DAPI staining). Hepa-1c1c7 cells were preincubated with rutaecarpine for 48 h before being treated with 500 μM H₂O₂ for 3 h. Cells were then fixed, permeabilized, and stained with DAPI (1 μg/ml). Nuclear fragmentation and condensation were visualized by nuclear fluorescence (arrow). (A) H₂O₂ treatment over a time course. (B) Treatment with various concentrations of rutaecarpine (0.16, 0.8, and 4 μM).




Fig. 2. Effects of different rutaecarpine concentrations on apoptosis of Hepa-1c1c7 cells. Cells were incubated with the indicated concentrations of rutaecarpine and exposed to 500 μM H₂O₂. The percentage of apoptotic cells was determined by flow cytometry after propidium iodide labeling.

nam, Korea). mRNA expression was determined by reverse-transcriptase polymerase chain reaction (RT-PCR) using the One-Step RT-PCR Premix (Intron Biotechnology). The amplification cycles comprised 94°C for 60 s, 55°C for 60 s, and 72°C for 60 s. After 30 cycles, the PCR products were subjected to electrophoresis on a 1% agarose gel. The electrophoresis products were visualized by ethidium bromide staining. The PCR products were quantified by measuring radioactivity using an LAS1000 image analyzer (Fuji, Tokyo, Japan) and image analysis software (Multi Gauge, ver. 3.0; Fuji).

Western blotting

Total protein was extracted from cultured cells using Pro-Pep reagent (Intron Biotechnology). The protein sample concentrations were adjusted so that they were equivalent using a bovine serum albumin (BSA) standard curve and diluted with Pro-Pep reagent. Samples (10 µg/lane) were separated by 9% sodium dodecyl sulfate polyacrylamide gel electrophoresis and then electrophoretically transferred to polyvinylidene fluoride (PVDF) membranes. The PVDF membranes were dried at 37°C for 1 h, blocked in blocking buffer (1% BSA) for 1 h, incubated overnight with primary antibodies at 4°C, washed, and incubated with second antibodies at 37°C for 1 h. The membranes were developed with enhanced chemilu-




Fig. 3. Evaluation of DNA damage by Comet assay. H₂O₂-induced DNA strand breaks in Hepa-1c1c7 cells were assessed by Comet assay. The effect of preincubation with different concentrations of rutaecarpine for 48 h on the number of DNA strand breaks induced by 500 µM H₂O₂ was investigated. **p*<0.05 vs. H₂O₂-treated controls.

minescence substrate (Intron Biotechnology) and the bands were quantified using an LAS1000 image analyzer and image analysis software (Multi Gauge, ver. 3.0).

RESULTS AND DISCUSSION

Oxidative stress induces cell damage that is mediated by ROS, including H₂O₂ and superoxide and hydroxyl radicals. Transient exposure of Hepa-1c1c7 cells to H₂O₂ (500 µM) triggered typical apoptosis, as evidenced by chromatin condensation, nuclear fragmentation, and DNA laddering. Apoptosis is programmed cell death characterized by a series of distinct morphological and biochemical alterations (Cohen, 1997).

Fig. 1 shows the condensation state of nuclei, as assessed by DAPI staining, after treatment with H₂O₂ over a time course (Fig. 1A) and in cells preincubated with rutaecarpine and then treated with 500 µM H₂O₂ for 3 h (Fig. 1B). Nuclear fragmentation and condensation were characterized by nuclear fluorescence (arrows), suggesting apoptosis in cells treated with H₂O₂ or pretreated with rutaecarpine before exposure to H₂O₂. As shown in Fig. 1, treatment with rutaecarpine reduced nuclear fragmentation and condensation.

To determine whether rutaecarpine protects Hepa-1c1c7 cells from apoptosis, cells were exposed to various concentrations of rutaecarpine (0.16, 0.8, and 4 µM) and then treated with the oxidative stress inducer H₂O₂ (500 µM). Cells were harvested at different time points and examined for DNA content. Apoptosis was investigated by assessing DNA distribution. Flow cytometry analysis demonstrating hypodiploid DNA. As shown in Fig. 2, 500 µM H₂O₂ induced apoptotic death in >60% of cells after incubation for 3 h, whereas cells pretreated with 4 µM rutaecarpine for 48 h were found to be detached from the plate, with a survival rate of 37.18 ± 1.76%.

A Comet assay was performed to evaluate the protective




Fig. 4. RT-PCR analysis of quinone reductase mRNA expression. RT-PCR was performed to examine the effect of rutaecarpine on QR mRNA expression in Hepa-1c1c7 cells. 1, control cells; 2, cells treated with 500 µM H₂O₂ alone; 3, cells treated with H₂O₂ and 0.16 µM rutaecarpine; 4, cells treated with H₂O₂ and 0.8 µM rutaecarpine; 5, cells treated with H₂O₂ and 4 µM rutaecarpine. **p*<0.05 vs. H₂O₂-treated controls.




Fig. 5. Western blot analysis of quinone reductase (QR) expression. Western blotting was performed to examine the effect of rutaecarpine on QR protein expression in Hepa-1c1c7 cells. 1, control cells; 2, cells treated with 500 μM H₂O₂ alone; 3, cells treated with H₂O₂ and 0.16 μM rutaecarpine; 4, cells treated with H₂O₂ and 0.8 μM rutaecarpine; 5, cells treated with H₂O₂ and 4 μM rutaecarpine. **p*<0.05 vs. H₂O₂-treated controls.

potential of rutaecarpine against H₂O₂-induced DNA damage. Cells were treated with 500 μM H₂O₂ for 3 h to induce DNA strand breaks. Pretreatment of cells with rutaecarpine for 48 h significantly reduced H₂O₂-induced DNA damage in a concentration-dependent manner (Fig. 3). Bao *et al.* (2011) reported that rutaecarpine prevents hypoxia/reoxygenation-induced myocardial cell injury and apoptosis through inhibition of nicotinamide adenine dinucleotide phosphate oxidases and the reactive oxygen species pathway.

QR is one of several enzymes that inactivate electrophilic compounds and is an important phase II enzyme in the detoxification of quinone. QR catalyzes the detoxification of unstable semiquinones by two-electron reduction to hydroquinone (Ross *et al.*, 2000). Induction of QR protects DNA against oxidative damage. In this study, QR gene expression in Hepa-1c1c7 cells after H₂O₂-induced DNA damage was determined by RT-PCR, and was found to be significantly reduced in cells incubated with 500 μM H₂O₂. However, pretreatment with 4 μM rutaecarpine for 48 h significantly increased QR gene expression up to 2–3-fold (Fig. 4, 5). These findings provide evidence that the protective effect of rutaecarpine against DNA damage is mediated by increased levels of QR.

Protein kinase C (PKC) plays a role in H₂O₂-induced apoptosis by regulating the PI3K/Akt pathway (Zhou *et al.*, 2005). PKC may play an important role in H₂O₂-induced apoptosis through negative regulation of PI3K/Akt and may act upstream of mitochondria (Zhang *et al.*, 2001). The major mechanism involved in the anti-apoptotic effect of PI3K/Akt is the phosphorylation of Bad, which is then sequestered in the cytosol and prevented from interacting with Bcl-2/Bcl-xl and disrupting the mitochondrial membrane (Alessi and Cohen, 1998).

To study the mechanism underlying the protective effect of rutaecarpine against DNA damage, the regulation of PI3K/Akt, which plays a critical role in cell survival, was assessed. As shown in Fig. 6, Hepa-1c1c7 cells undergo apoptosis when




Fig. 6. Western blot analysis of PI3K and p-Akt. (A) Activation of signal transduction pathways by H₂O₂. Hepa-1c1c7 cells were treated with H₂O₂ over a time course and analyzed by western blotting. (B) Hepa-1c1c7 cells were pretreated with rutaecarpine for 48 h, treated with H₂O₂ for 3 h, and analyzed by western blotting.

exposed to H₂O₂ in a time-dependent manner. To identify the signaling mechanisms activated by H₂O₂ in Hepa-1c1c7 cells, we assessed the phosphorylation of Akt and expression of PI3K. pAkt was detected as soon as 10 min after the addition of 500 μM H₂O₂, peaked at 30 min and 1 h, and returned to basal levels by 3 h (Fig. 6A). Treatment with 500 μM H₂O₂ resulted in a marked increase in the PI3K level. Inhibition of apoptosis was assayed at 3 h after the addition of 500 μM H₂O₂. To explore the H₂O₂ responsiveness of the PI3K/Akt signal transduction pathway, Hepa-1c1c7 cells were pretreated with dimethyl sulfoxide (vehicle control) or rutaecarpine for 48 h before treatment with H₂O₂ for 3 h. Protein extracts were analyzed for levels of PI3K and pAkt by western blotting with appropriate antibodies. The results demonstrated increased PI3K and pAkt protein levels in rutaecarpine-treated cells compared with untreated control cells (Fig. 6B).

In conclusion, rutaecarpine reduced the condensation and fragmentation of nuclei, as assessed by DAPI staining; inhibited apoptosis, as assessed by fluorescence-activated cell sorting and PI staining; and reduced the number of oxidative stress-induced DNA strand breaks, as assessed by Comet assay. Rutaecarpine also increased QR gene expression in cells subjected to H₂O₂-induced DNA damage. These results suggest that rutaecarpine protects Hepa-1c1c7 cells against oxidative stress-induced DNA damage.

ACKNOWLEDGMENTS

This research was supported by the Basic Science Research Program of the National Research Foundation of Korea, funded by the Ministry of Education, Science and Technology (grant number 120120392).

REFERENCES

Ahlgren-Beckendorf, J. A., Reising, A. M., Schander, M. A., Herdler,

- J. W. and Johnson, J. A. (1999) Coordinate regulation of NAD(P)H:quinone oxidoreductase and glutathione-S-transferases in primary cultures of rat neurons and glia: role of the antioxidant/electrophile responsive element. *Glia* **25**, 131-142.
- Ahn, H., Nam, J. W., Seo, E. K. and Mar, W. (2008) Induction of NAD(P)H:quinone reductase by rutaecarpine isolated from the fruits of *Evodia rutaecarpa* in the murine hepatic Hepa-1c1c7 cell line. *Planta Med.* **74**, 1387-1390.
- Alessi, D. R. and Cohen, P. (1998) Mechanism of activation and function of protein kinase B. *Curr. Opin. Genet. Dev.* **8**, 55-62.
- Bao, M. H., Dai, W., Li, Y. J. and Hu, C. P. (2011) Rutaecarpine prevents hypoxia-reoxygenation-induced myocardial cell apoptosis via inhibition of NADPH oxidases. *Can. J. Physiol. Pharmacol.* **89**, 177-186.
- Burdette, J. E., Chen, S. N., Lu, Z. Z., Xu, H., White, B. E., Fabricant, D. S., Liu, J., Fong, H. H., Farnsworth, N. R., Constantinou, A. I., Van Breemen, R. B., Pezzuto, J. M. and Bolton, J. L. (2002) Black cohosh (*Cimicifuga racemosa* L.) protects against menadione-induced DNA damage through scavenging of reactive oxygen species: bioassay-directed isolation and characterization of active principles. *J. Agric. Food Chem.* **50**, 7022-7028.
- Cohen, G. M. (1997) Caspases: the executioners of apoptosis. *Biochem. J.* **326**, 1-16.
- Deguil, J., Jailloux, D., Page, G., Fauconneau, B., Houeto, J. L., Philippe, M., Muller, J. M. and Pain, S. (2007) Neuroprotective effects of pituitary adenylate cyclase-activating polypeptide (PACAP) in MPP⁺-induced alteration of translational control in Neuro-2A neuroblastoma cells. *J. Neurosci. Res.* **85**, 2017-2025.
- Finkel, T. and Holbrook, N. J. (2000) Oxidants, oxidative stress and the biology of ageing. *Nature* **408**, 239-247.
- Finley, J. W. (2003) The antioxidant responsive element (ARE) may explain the protective effects of cruciferous vegetables on cancer. *Nutr. Rev.* **61**, 250-254.
- Loft, S. and Poulsen, H. E. (1996) Cancer risk and oxidative DNA damage in man. *J. Mol. Med.* **74**, 297-312.
- Martindale, J. L. and Holbrook, N. J. (2002) Cellular response to oxidative stress: signaling for suicide and survival. *J. Cell Physiol.* **192**, 1-15.
- Ross, D., Kepa, J. K., Winski, S. L., Beall, H. D., Anwar, A. and Siegel, D. (2000) NAD(P)H:quinone oxidoreductase 1 (NQO1): chemoprotection, bioactivation, gene regulation and genetic polymorphisms. *Chem. Biol. Interact.* **129**, 77-97.
- Talalay, P., Fahey, J. W., Holtzclaw, W. D., Prester, T. and Zhang, Y. (1995) Chemoprotection against cancer by phase 2 enzyme induction. *Toxicol. Lett.* **82-83**, 173-179.
- Traister, A., Breitman, I., Bar-Lev, E., Zvibel, I., Harel, A., Halpern, Z. and Oren, R. (2005) Nicotinamide induces apoptosis and reduces collagen I and pro-inflammatory cytokines expression in rat hepatic stellate cells. *Scand. J. Gastroenterol.* **40**, 1226-1234.
- Zhang, L., Himi, T., Morita, I. and Murota, S. (2001) Inhibition of phosphatidylinositol-3 kinase/Akt or mitogen-activated protein kinase signaling sensitizes endothelial cells to TNF- α cytotoxicity. *Cell Death Differ.* **8**, 528-536.
- Zhou, Y., Wang, Q., Evers, B. M. and Chung, D. H. (2005) Signal transduction pathways involved in oxidative stress-induced intestinal epithelial cell apoptosis. *Pediatr. Res.* **58**, 1192-1197.