



Bactericidal Efficacies of an Aquatic Disinfectant Tablet Composed to Calcium Hypochlorite Against *Vibrio anguillarum* and *Streptococcus iniae*

Chun-Nam Cha¹, Yeo-Eun Lee², In-Jin Kang², Chang-Yeul Yoo³, Hyunju Choi⁴, Suk Kim, and Hu-Jang Lee*

Institute of Animal Medicine, College of Veterinary Medicine, Gyeongsang National University, Chinju 600-701, Korea

¹*Engineering Research Institute, Department of Industrial Systems Engineering, Gyeongsang National University, Chinju 600-701, Korea*

²*Department of Environmental Health, Graduate School of Public Health, Gyeongsang National University, Chinju 660-751, Korea*

³*Department of Computer Information, Gyeongnam Provincial Namhae College, Namhae 668-801, Korea*

⁴*Elderly Life Redesign Institute, Department of Biomedical Laboratory Science, Inje University, Gimhae 621-749, Korea*

(Received June 16, 2012/Revised June 24, 2012/Accepted July 8, 2012)

ABSTRACT - *Vibrio* spp. and *Streptococcus* spp. have caused a considerable disease of farmed fish and economic loss in fish farming and seafood industry. In this study, the efficacy of an aquatic disinfectant tablet composed to calcium hypochlorite was evaluated against *V. anguillarum* and *S. iniae*. A bactericidal efficacy test by broth dilution method was used to determine the lowest effective dilution of the disinfectant following exposure to test bacteria for 30 min at 4°C. An aquatic disinfectant tablet and test bacteria were diluted with distilled water (DW), hard water (HW) or organic matter suspension (OM) according to treatment condition. *V. anguillarum* on the DW, HW and OM condition was completely inactivated with 16,000 15,000 and 13,000 fold dilutions of the disinfectant, respectively. On the DW, HW and OM condition, *S. iniae* was absolutely inactivated with 17,000 16,000 and 14,000 fold dilutions of the disinfectant, respectively. As an aquatic disinfectant tablet possesses bactericidal efficacy against fish pathogenic bacteria such as *V. anguillarum* and *S. iniae* this disinfectant solution can be used to control the spread of fish infective bacterial diseases.

Key words: An aquatic disinfectant tablet, *Vibrio anguillarum*, *Streptococcus iniae*, Calcium hypochlorite, Disinfectant efficacy

The genus *Vibrio* includes gram-negative, facultative anaerobic, non-spore-forming bacilli that are oxidase positive and halophilic¹. Several species of *Vibrio* can cause foodborne infection, usually associated with eating undercooked seafood². Most disease-causing *Vibrio* strains are associated with gastroenteritis, but can also infect open wounds and cause septicemia^{3,4}. It can be carried by numerous sea-living animals, such as crabs or prawns, and has been known to cause fatal infections in humans during exposure⁵.

Vibrio anguillarum (*V. anguillarum*) is the causative agent of vibriosis, a deadly hemorrhagic and septicemic disease affecting various marine and fresh/brackish water fish, bivalves and crustaceans. In both aquaculture and larviculture, this disease is responsible for severe economic losses worldwide⁶.

Streptococcus iniae (*S. iniae*) is a species of Gram-positive,

sphere-shaped bacterium belonging to the genus *Streptococcus*⁷. Since its isolation from an Amazon freshwater dolphin in 1976, *S. iniae* has emerged as a leading fish pathogen in aquaculture operations worldwide, resulting in over \$150 million in annual losses^{8,9}. *S. iniae* infections have been reported in at least 27 species of cultured or wild fish from around the world¹⁰. *S. iniae* is highly pathogenic in cultured fishes with 30-50% mortality¹¹. *S. iniae* can cause opportunistic infections in weakened or immunocompromised humans¹².

As *V. anguillarum* and *S. iniae* infections are becoming harder to control because of resistance to commonly used antibiotics, the effective cleaning and disinfection regimes are essential for the prevention of infections and outbreaks^{13,14}. The cleaning and disinfectant regimes depend on the proper use of biocides, and there is the concern that the resulting increased use of biocides in fish farming, food production, and hospital settings, and the home could contribute to the selection of antibiotic-resistant strains as some mechanisms of biocide resistance also confer antibiotic resistance¹⁵. Biocides are often composed of a mixture of ingredients that act upon

*Correspondence to: Hu-Jang Lee, DVM, MPH, PhD, College of Veterinary Medicine, Gyeongsang National University, 900 Gajwadong, Chinju 660-701, Korea

Tel: 82-55-772-2352, Fax: 82-55-772-2308

E-mail: hujang@gnu.ac.kr

a wide range of cellular mechanisms and targets, which makes it difficult for bacteria to become resistant to biocides¹⁶).

Vibriosis and *S. iniae* infection in cultured marine fishes and human may cause enormous economic loss^{6,9}). The stress on cultured marine fishes caused by intensive farming systems, and the development of antibiotic-resistant bacteria is among the major reasons for the increased frequency of bacterial disease outbreaks¹⁷). Highly hygienic measures including the use of disinfectant are very effective for successful control of diseases from potential pathogens in cultured marine fishes¹⁸). Various disinfectants have been used for decontamination of cultured marine fishes and biological agents¹⁹). However, there is not the efficacy test for the disinfectant composed of calcium hypochlorite against *V. anguillarum* and *S. iniae*. Therefore, this study was carried out to examine bactericidal efficacy of a disinfectant tablet against *V. anguillarum* and *S. iniae*.

Materials and Methods

Bacteria and culture

The test bacteria, *V. anguillarum* (KCTC 2711) and *S. iniae* (KCTC 3657) were obtained from the National Fisheries Research and Development Institute (Busan, Korea). The strains were maintained as frozen glycerol stock. *V. anguillarum* cells were cultured in tryptic soy broth (TSB) containing 1.5% agar and 1.0% NaCl at 25°C for 24–28 h. *S. iniae* cells were spread in brain heart infusion (BHI) broth containing 1.5% NaCl and incubated at 28°C for 24–48 h.

Disinfectant

The active ingredient for Easy-Wash[®], the tested disinfectant tablet, is calcium hypochlorite (70% w/w). Easy-Wash[®] was provided by Dae Han New Pharm Co. (Seoul, Korea). The disinfectant tablet was stored in room temperature and prepared for dilution on the day of evaluation. Determination of the antimicrobial efficacy of the disinfectant was based on Animal, Plant and Fisheries Quarantine and Inspection Agency Regulation No. 2011-26, Korea.

Diluents and treatment condition

Testing was based on bactericidal effects of disinfectant diluents in three treatment conditions (distilled water (DW) condition, standard hard water (HW) condition, and organic matter (OM) condition), pathogen control (disinfectant negative control) and DW control (both disinfectant and pathogen negative control) in Table 1. HW, an ingredient of HW treatment condition, was made by adding anhydrous CaCl₂ 0.305 g and MgCl₂·6H₂O 0.139 g into one liter distilled water. Organic suspension, an ingredient of OM treatment condition, is a solution of 5% (w/v) yeast extract in HW. The test organisms

Table 1. Experimental design for the determination of the bactericidal efficacy of an aquatic disinfectant tablet

Treatment condition*	Contents according to treatment condition**				
	DM	HW	OM	Disinfectant	Bacteria
DW condition	+	-	-	+	+
HW condition	-	+	-	+	+
OM condition	-	-	+	+	+
Bacteria control	-	+	-	-	+
DW control	+	-	-	-	+

*DW, distilled water; HW, standard hard water; OM, organic matter.

**+, presence; -, absence

were prepared by titration of each cultural broth into at least 10⁸ CFU/ml viable organisms with the same kind of diluents of treatment condition.

Experimental procedures

For the efficacy test against *V. anguillarum*, an aquatic disinfectant tablet was diluted 12,800, 14,400, 16,000, 17,600 and 19,200 times with DW, 12,000, 13,500, 15,000, 16,500 and 18,000 times with HW, and 10,400, 11,700, 13,000, 14,300 and 15,600 times with OM, respectively. For the efficacy test against *S. iniae*, Easy-Wash[®] was also diluted 13,600, 15,300, 17,000, 18,700 and 20,400 times with DW, 12,800, 14,400, 16,000, 17,600 and 19,200 times with HW, and 11,200, 12,600, 14,000, 15,400 and 16,800 times with OM, respectively.

To verify the lowest effective dilution of the disinfectant, five serial dilutions of the disinfectant were prepared and placed at 4°C prior to test reaction. 2.5 ml of each disinfectant dilution was mixed with the same amount of test organism followed by contact time of 30 min at 4°C.

During this period, the mixture was shaken at 10 min interval. At the end of 30 min contact period, one ml of the mixture was neutralized with 9 ml of nutrient broth containing 5% inactivated horse serum (Becton Dickinson & Co., MD, USA) at 37°C. 0.1 ml of the neutralized reaction mixture was subcultured into 10 ml of recovery each cultural broth at 37°C for 48 h in incubator. The valid dilution was determined that the greatest dilution showing no growth in two or more in the five replicates was confirmed. The final dilution time was statistically determined by a median value among three valid dilution of the triplicate test, but each value of which should be within 20% experimental error.

Results

Table 2 shows the final valid dilution of an aquatic disinfectant tablet composed to calcium hypochlorite. On DW condition, *V. anguillarum* and *S. iniae* were completely inactivated with 16,000 and 17,000 fold dilutions of the

Table 2. Final valid dilution of an aquatic disinfectant tablet against *V. anguillarum* and *S. iniae*

Bacterial strains	Treatment condition*											
	DW				HW				OM			
	DT	1	2	3	DT	1	2	3	DT	1	2	3
<i>V. anguillarum</i>	12,800	×	×	×	12,000	×	×	×	10,400	×	×	×
	14,400	×	×	×	13,500	×	×	×	11,700	×	×	×
	16,000	×	×	○	15,000	×	×	×	13,000	○	×	×
	17,600	○	○	○	16,500	○	×	○	14,300	○	○	○
	19,200	○	○	○	18,000	○	○	○	15,600	○	○	○
	Valid	16,000			Valid	15,000			Valid	13,000		
<i>S. iniae</i>	13,600	×	×	×	12,800	×	×	×	11,200	×	×	×
	15,300	×	×	×	14,400	×	×	×	12,600	×	×	×
	17,000	×	○	×	16,000	○	×	×	14,000	×	×	○
	18,700	○	○	○	17,600	○	○	○	15,400	○	○	○
	20,400	○	○	○	19,200	○	○	○	16,800	○	○	○
	Valid	17,000			Valid	16,000			Valid	14,000		

*DW, distilled water; HW, standard hard water; OM, organic matter; DT, dilution time.

○, growth; ×, growth inhibition

disinfectant, respectively. When the bactericidal effect on HW condition was evaluated, *V. anguillarum* and *S. iniae* were completely inactivated with 15,000 and 16,000 fold dilutions of the disinfectant, respectively. With the investigation of the bactericidal effect of the disinfectant on OM condition, *V. anguillarum* and *S. iniae* were completely inactivated with 13,000 and 14,000 fold dilutions of the disinfectant, respectively. Because the organic material interferes with efficacy by either inactivating the disinfectant or blocking it from surface contact, the bactericidal activity of the disinfectant on the OM condition was lowered against fish pathogenic bacteria compared with DM or HW conditions.

Comparing the results of the disinfectant against two pathogenic bacteria in the present study, the bactericidal effect of an aquatic disinfectant tablet against *S. iniae* was higher than that against *V. anguillarum* on all experimental conditions.

Discussion

An aquatic disinfectant tablet is a potential antibacterial disinfectant which was composed of calcium hypochlorite. Calcium hypochlorite is widely used for water treatment and as a bleaching agent. And this compound is considered to be relatively stable and has greater available chlorine than sodium hypochlorite²⁰.

Calcium hypochlorite has been used for water sanitization as an alternative of chlorine-based sanitizer. Aqueous calcium hypochlorite has a much higher oxidation capacity than chlorine due to the formation of hypochlorous acid²¹. Bacterial inactivation by hypochlorous acid is the result of

inhibition of bacterial DNA replication²². Yousuf *et al.*²³ previously reported that calcium hypochlorite was the most effective than trisodium phosphate, sodium carbonate, and organic acids in reduction of total bacterial load as well as elimination of harmful bacteria in shrimp and prawn, and after treatment of 10 ppm calcium hypochlorite, the count of *Vibrio* spp. in shrimp and prawn reduced 0.23log₁₀ CFU/g to zero. Kim *et al.*²⁴ reported that antimicrobial effects against *Vibrio* spp., *Edwardsiella tarda*, *Streptococcus* spp., and *Staphylococcus* spp. were tested using 10 different disinfectants including hypochlorite salts. Hypochlorite salts resulted in 99.99% bactericidal activity against four strains of fish pathogenic bacteria. Abdel-Rahman *et al.*²⁵ reported that when Sprague-Dawley rats were administered hypochlorous acid at 0, 1, 10 or 100 mg/l daily in drinking water for one year, no significant chloroform concentrations were observed in rat blood at 12 months. In addition, OECD SIDS²⁶ reported that the acute oral toxicological test of calcium hypochlorite was carried out using Wistar-derived albino rats dosed at 890, 1000, 1120, and 1260 mg/kg body weight, and with the results of the study, the LD₅₀ of calcium hypochlorite was calculated to be 790 mg/kg.

With the consideration of previous studies, an aquatic disinfectant tablet is a more effective and safe disinfectant than chlorine and other treatments like ozone and beam irradiation against fish pathogenic bacteria.

In this study, disinfectant efficacy of an aquatic disinfectant tablet has limitation that the results are based on *in vitro* test. Organic material in suspension (OM condition) could not represent all possible parameters of *Vibrio* spp. and *Streptococcus* spp. contaminated fish farm and seafood-industry

environments.

As the efficacy of an aquatic disinfectant tablet against *V. anguillarum* and *S. iniae* was investigated *in vitro*, a controlled field trial is required to determine whether use of an aquatic disinfectant tablet will be able to reduce new pathogenic bacteria infection in fish farm and seafood industry area.

Conclusions

In cultured marine fish farm and seafood industry, vibriosis and *S. iniae* infection were very important diseases because of high mortality for farmed fish, zoonoses and economic loss. In the study of the bactericide efficacy test of an aquatic disinfectant tablet, the results suggest that an aquatic disinfectant tablet has potential bactericidal activity against *V. anguillarum* and *S. iniae*. So, an aquatic disinfectant tablet composed to calcium hypochlorite can be used to control the spread of fish pathogenic bacterial diseases.

Acknowledgements

This work was financially supported by Dae Han New Pharm Co., LTD. (Seoul, Korea).

References

1. Raissy, M., Moumeni, M., Ansari, M. and Rahimi, E.: Occurrence of *Vibrio* spp. in lobster and crab from the Persian gulf. *J. Food Safety* **32**, 198-203 (2012).
2. Amaro, C., Fouz, B., Biosca, E.G., Marconoales, E. and Collado, R.: The lipopolysaccharide O side chain of *Vibrio vulnificus* serogroup E is a virulence determinant for eels. *Infect. Immun.* **65**, 2475-2479 (1997).
3. Chung, P.H., Chuang, S.K., Tsang, T., Wai-man, L., Yung, R. and Lo, J.: Cutaneous injury and *Vibrio vulnificus* infection. *Emerg. Infect. Dis.* **12**, 1302-1303 (2006).
4. Chou, T.K., Lee, Y.T., Lai, Y.Y., Chao, W.N., Yang, C., Chen, C.C., Wang, P.H., Lin, D.B., Wong, R.H. and Chen, S.C.: Prognostic factors for primary septicemia and wound infection caused by *Vibrio vulnificus*. *Am. J. Emerg. Med.* **28**, 424-431 (2010).
5. Daniels, N.A. and Shafaie, A.: A review of pathogenic *Vibrio* infections for clinicians. *Infect. Med.* **17**, 335-685 (2000).
6. Frans, I., Michiels, C.W., Bossier, P., Willems, K.A., Lievens, B. and Rediers, H.: *Vibrio anguillarum* as a fish pathogen: virulence factors, diagnosis and prevention. *J. Fish Dis.* **34**, 643-661 (2011).
7. Peir, G.B. and Madin, S.H.: Isolation and characterization of a second isolate of *Streptococcus iniae*. *Int. J. Syst. Bacteriol.* **28**, 311-314 (1978).
8. Pier, G.B. and Madin, S.H.: *Streptococcus iniae* sp. nov., a beta-hemolytic streptococcus isolated from an Amazon freshwater dolphin, *Inia geoffrensis*. *Int. J. Syst. Bacteriol.* **26**, 545-553 (1976).
9. McNulty, S.T., Klesius, P.H., Shoemaker, C.A. and Evans, J.J.: *Streptococcus iniae* infection and tissue distribution in hybrid striped bass (*Morone chrysops* × *Morone saxatilis*) following inoculation of the gills. *Aquaculture* **220**, 165-173 (2003).
10. Zhou, S.M., Xie, M.Q., Zhu, X.Q., Ma, Y., Tan, Z.L. and Li, A.X.: Identification and genetic characterization of *Streptococcus iniae* strains isolated from diseased fish in China. *J. Fish Dis.* **31**, 869-875 (2008).
11. Weinstein, M.R., Litt, M., Kertesz, D.A., Wyper, P., Rose, D., Coulter, M., McGeer, A., Facklam, R., Ostach, C., Willey, B.M., Borczyk, A., Low, D.E.: Invasive infections due to a fish pathogen, *Streptococcus iniae*. *S. iniae* study group. *N. Engl. J. Med.* **337**, 589-594 (1997).
12. Agnew, W. and Barnes, A.C.: *Streptococcus iniae*: an aquatic pathogen of global veterinary significance and a challenging candidate for reliable vaccination". *Vet. Microbiol.* **122**, 1-15 (2007).
13. Pedersen, K., Tiainen, T. and Larsen, J.L.: Antibiotic resistance of *Vibrio anguillarum*, in relation to serovar and plasmid contents. *Acta Vet. Scand.* **36**, 55-64 (1995).
14. Facklam, R., Elliott, J., Shewmaker, L. and Reingold, A.: Identification and characterization of sporadic isolates of *Streptococcus iniae* isolated from humans. *J. Clin. Microbiol.* **43**, 933-937 (2005).
15. Russell, A.D.: Biocide use and antibiotic resistance: the relevance of laboratory findings to clinical and environmental situations. *Lancet Infect. Dis.* **3**, 794-803 (2003).
16. Cha, C.N., Lee, Y.E., Kang, I.J., Yoo, C.Y., An, S., Kim, S. and Lee, H.J.: Bactericidal efficacy of Vital-Oxide[®], disinfectant solution against *Salmonella typhimurium* and *Brucella ovis*. *J. Fd. Hyg. Safety* **27**, 50-54 (2012).
17. Maqsood, S., Singh, P., Samoon, M.H. and Munir, K.: Emerging role of immunostimulants in combating the disease outbreak in aquaculture. *Int. Aquat. Res.* **3**, 147-163 (2011).
18. Kasai, H., Yoshimizu, M. and Ezura, Y.: Disinfection of water for aquaculture. *Fish. Sci.* **68**, 821-824 (2002).
19. Danner, G.R. & Merrill, P.: Aquaculture biosecurity, prevention, control, and eradication of aquatic animal. In *Disinfectants, disinfection, and biosecurity in aquaculture* (Scarfe, A.D., Lee, C.S. and O'Bryen, P.J. eds.) Blackwell Publishing, Oxford, pp. 90-128 (2006).
20. Epshtein, A.E., Polosina, Y.A. and Ioirish, A.N.: Studies on the stability of disinfectant solutions containing active chlorine. *Hyg. Sanit.* **3**, 86-87 (1986).
21. Willians, D.E., Worley, S.D., Wheatley, W.B. and Swango, L.J.: Bactericidal properties of a new water disinfectant. *Appl. Environ. Microbiol.* **49**, 637-643 (1985).
22. McKenna, S.M. and Davies, K.J.: The inhibition of bacterial growth by hypochlorous acid. Possible role in the bactericidal activity of phagocytes. *Biochem. J.* **254**, 685-692 (1988).
23. Yousuf, A.H.M., Ahmed, M.K., Yeasmin, S., Ahsan, N., Rahman, M.M. and Islam, M.M.: Reduction of bacterial pathogens in *Penaeus monodon* and *Macrobrachium rosenbergii* using several chemical interventions. *World J. Agri. Sci.* **4**, 856-861 (2013).

- (2008).
24. Kim, S.R., Park, K.H., Kim, D., Jung, S.J., Kang, S.Y. and Oh, M.J.: Antimicrobial effects of chemical disinfectants on fish pathogenic bacteria. *Food Sci. Biotechnol.* **17**, 971-975 (2008).
25. Abdel-Rahman, M.S., Suh, D.H. and Bull. R.J.: Pharmacodynamics and toxicity of chlorine in drinking water in the rat. *J. Appl. Toxicol.* **4**, 82-86 (1984).
26. OECD SIDS: Calcium hypochlorite (CAS No. 7778-54-3). UNEP-Publications, Paris, pp. 10-15 (2004).