

Bone thickness of the infrazygomatic crest area in skeletal Class III growing patients: A computed tomographic study

Hyub-Soo Lee¹, Hang-Moon Choi², Dong-Soon Choi¹, Insan Jang¹, Bong-Kuen Cha^{1,*}

¹Department of Orthodontics, College of Dentistry and Research Institute of Oral Science, Gangneung-Wonju National University, Gangneung, Korea

²Department of Oral and Maxillofacial Radiology, College of Dentistry and Research Institute of Oral Science, Gangneung-Wonju National University, Gangneung, Korea

ABSTRACT

Purpose: This study was performed to investigate the bone thickness of the infrazygomatic crest area by computed tomography (CT) for placement of a miniplate as skeletal anchorage for maxillary protraction in skeletal Class III children.

Materials and Methods: CT images of skeletal Class III children (7 boys, 9 girls, mean age: 11.4 years) were taken parallel to the Frankfurt horizontal plane. The bone thickness of the infrazygomatic crest area was measured at 35 locations on the right and left sides, perpendicular to the bone surface.

Results: The bone was thickest (5.0 mm) in the upper zygomatic bone and thinnest (1.1 mm) in the anterior wall of the maxillary sinus. Generally, there was a tendency for the bone to be thicker at the superior and lateral area of the zygomatic process of the maxilla. There was no clinically significant difference in bone thickness between the right and left sides; however, it was thicker in male than in female subjects.

Conclusion: In the infrazygomatic crest area, the superior and lateral area of the zygomatic process of the maxilla had the most appropriate thickness for placement of a miniplate in growing skeletal Class III children with a retruded maxilla. (*Imaging Sci Dent* 2013; 43: 261-6)

KEY WORDS: Computed Tomography, Maxillary Sinus, Zygoma, Orthodontic Anchorage Techniques

Introduction

Protraction headgear therapy is widely used as a treatment modality for growing patients with skeletal Class III malocclusion and maxillary retrusion. Traditionally, the maxillary teeth have been used as an anchorage for applying forces of protraction to the craniofacial complex. However, indirect application of force to the maxilla often causes undesirable tooth movement, such as proclination of the maxillary incisors, or extrusion and mesial movement of the maxillary posterior teeth.¹⁻³ These side effects

increase as children grow older because ossification of the circumaxillary sutures progresses and the resistance to protraction increases.⁴ To minimize undesirable movement of the teeth and apply the force directly to the maxilla, skeletal anchorage has been suggested as an alternative to conventional tooth-borne anchorage.⁵⁻⁸ We previously reported the successful protraction of the maxilla with miniplates in the infrazygomatic crest area as an anchorage in patients with skeletal Class III malocclusion and maxillary retrusion.⁹⁻¹¹ Miniplates installed in the infrazygomatic crest area would also be helpful as an anchorage for a range of other orthodontic tooth movements.¹²⁻¹⁴

In a few cases, however, the miniplates lost primary stability, requiring a reoperation for fixation of the miniplate or modification of the treatment plan. Bone thickness and density are considered to be important factors in

Received April 28, 2013; Revised May 14, 2013; Accepted June 2, 2013

*Correspondence to : Prof. Bong-Kuen Cha

Department of Orthodontics, Gangneung-Wonju National University Dental Hospital, 7 Jukheon-gil, Gangneung city, Gangwon province 210-702, South Korea
Tel) 82-33-640-3192, Fax) 82-33-640-3057, E-mail) korth@gwnu.ac.kr

Copyright © 2013 by Korean Academy of Oral and Maxillofacial Radiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Imaging Science in Dentistry · pISSN 2233-7822 eISSN 2233-7830




Fig. 1. Construction of the horizontal base plane (HB) and antero-posterior line. A. HB is perpendicular to the midsagittal plane and passes through the most inferior border of the zygomatic process of the maxilla or zygomatic bone. B. The antero-posterior line (AP) is drawn on each horizontal plane parallel to the midsagittal plane and passes through the most anterior point of the infratemporal fossa (arrow).

the stability of such miniplates.^{15,16} However, to date, no study has evaluated the bone thickness of the infrazygomatic crest area. The purpose of this study was to evaluate the cortical bone thickness of the infrazygomatic crest area of the maxilla to identify a more suitable region for miniplate placement in growing skeletal Class III patients.

Materials and Methods

The sample consisted of the CT images (Shimadzu Co., Tokyo, Japan) obtained from 16 children (7 boys and 9 girls) with a retruded maxilla and ANB angle of less than 0 degrees. The age of patients ranged from 10 to 13 years (mean age: 11.4 ± 1.7 years), and they had no facial asymmetry or general medical anamnesis. Placement of miniplates on the infrazygomatic crest area was planned for protraction of the maxilla. A spiral CT was taken at 120 kVp, 230 mA with a 1 : 1.2 pitch, a scanning time of 1.5 s, and a slice thickness of 1 mm to obtain the axial images between the maxillary occlusal plane and the inferior margin of the orbit. The window width was 0 Hounsfield units (HU) with a center of 500 HU. The CT images were taken in parallel to the Frankfurt horizontal (FH) plane.

Measurement

The horizontal base plane (HB) was defined as a plane perpendicular to the midsagittal plane and passing the most inferior border of the zygomatic process of the maxilla or zygomatic bone (Fig. 1A). Six more horizontal planes

(HB-2, HB+2, HB+4, HB+6, HB+8, HB+10) parallel with the HB were constructed at 2-mm intervals. The bone located lower than HB-2 was not estimated in the present study because the developing tooth germ or root apex was observed in several patients with late mixed dentition or early permanent dentition.

On each horizontal plane of HB, HB+2, HB+4, HB+6, HB+8, and HB+10, the antero-posterior line (AP) was drawn parallel with the midsagittal plane and passing through the most anterior point of the infratemporal fossa (Fig. 1B, arrow). Five more antero-posterior lines (AP-2, AP-4, AP-6, AP-8, and AP-10) were constructed at 2-mm intervals. On the horizontal plane of HB-2, AP was defined as the line parallel with the midsagittal plane and passing through the most lateral point of the maxilla. The vertical lines of AP-2, AP-4, AP-6, AP-8, and AP-10 were also drawn on HB-2 at 2-mm intervals (Fig. 1A). The bone thickness was measured at the points where the antero-posterior lines and outer surface of the bone intersect at each horizontal plane. All of the measurements were performed by one investigator.

Method errors

The reliability of the measurements was determined on two arbitrarily selected samples. The bone thickness at each point was measured twice, separated by a one-week interval. The method error was calculated from Dahlberg's formula (method error = $\sqrt{\sum d^2 / 2n}$ where d is the difference between 2 measurements of a pair, and n is the number of subjects). The method error of the measure-

Table 1. The minimum (Min), median, maximum (Max) values of the bone thickness in the right and left sides (unit: mm)

	Right			Left			P value	Total		
	Min	Median	Max	Min	Median	Max		Min	Median	Max
HB-2, VL-2	0.3	3.0	6.2	0.5	3.0	6.0	N/S	0.3	3.0	6.2
HB-2, VL-4	0.3	1.9	5.8	0.5	1.8	4.8	N/S	0.3	1.8	5.8
HB-2, VL-6	0.3	1.6	3.5	0.5	1.4	5.0	N/S	0.3	1.4	5.0
HB-2, VL-8	0.3	1.2	2.1	0.5	1.3	3.1	N/S	0.3	1.2	3.1
HB-2, VL-10	0.3	1.1	2.1	0.3	1.1	2.7	N/S	0.3	1.1	2.7
HB, VL-2	1.2	3.2	7.4	0.5	3.6	6.4	N/S	0.5	3.4	7.4
HB, VL-4	0.5	2.4	6.9	0.5	2.8	6.2	N/S	0.5	2.6	6.9
HB, VL-6	0.5	1.7	4.4	0.5	1.8	6.2	N/S	0.5	1.7	6.2
HB, VL-8	0.5	1.4	2.3	0.5	1.4	5.6	N/S	0.5	1.4	5.6
HB, VL-10	0.5	1.2	1.8	0.5	1.1	4.3	N/S	0.5	1.1	4.3
HB+2, VL-2	1.6	3.7	7.6	1.7	3.9	8.4	N/S	1.6	3.7	8.4
HB+2, VL-4	0.8	2.1	8.8	1.3	2.3	7.9	N/S	0.8	2.1	8.8
HB+2, VL-6	0.3	1.6	8.9	0.8	1.5	5.6	N/S	0.3	1.5	8.9
HB+2, VL-8	0.3	1.2	4.5	0.6	1.2	3.2	N/S	0.3	1.2	4.5
HB+2, VL-10	0.3	1.1	2.7	0.3	1.1	3.0	N/S	0.3	1.1	3.0
HB+4, VL-2	2.0	4.0	9.6	1.5	3.9	10.6	N/S	1.5	4.0	10.6
HB+4, VL-4	0.9	2.4	8.8	1.3	2.4	7.0	N/S	0.9	2.4	8.8
HB+4, VL-6	0.7	1.8	5.4	0.8	1.7	5.6	N/S	0.7	1.7	5.6
HB+4, VL-8	0.5	1.3	4.3	0.6	1.3	3.5	N/S	0.5	1.3	4.3
HB+4, VL-10	0.3	1.0	2.5	0.3	1.3	2.4	N/S	0.3	1.2	2.5
HB+6, VL-2	2.0	3.5	10.2	2.1	4.4	10.8	0.043 *	2.0	4.1	10.8
HB+6, VL-4	0.9	2.6	7.3	1.1	3.1	7.0	0.006**	0.9	2.8	7.3
HB+6, VL-6	0.8	1.7	5.1	0.8	2.0	5.6	0.028*	0.8	1.8	5.6
HB+6, VL-8	0.7	1.4	3.4	0.7	1.6	3.4	N/S	0.7	1.5	3.4
HB+6, VL-10	0.3	1.1	2.6	0.3	1.3	2.1	N/S	0.3	1.2	2.6
HB+8, VL-2	1.5	4.4	10.9	3.1	5.1	9.9	0.020*	1.5	4.6	10.9
HB+8, VL-4	1.1	2.9	6.8	1.6	3.6	6.5	0.008**	1.1	3.6	6.8
HB+8, VL-6	0.6	1.8	4.4	1.0	2.5	5.1	0.017*	0.6	2.2	5.1
HB+8, VL-8	0.5	1.4	3.2	0.8	1.8	3.5	0.008**	0.5	1.6	3.5
HB+8, VL-10	0.3	1.3	3.1	0.3	1.4	2.2	N/S	0.3	1.3	3.1
HB+10, VL-2	1.5	4.7	9.5	3.7	5.2	9.4	0.004**	1.5	5.0	9.5
HB+10, VL-4	0.9	3.6	6.3	2.3	4.1	10.6	0.008**	0.9	3.8	10.6
HB+10, VL-6	0.8	2.0	4.4	1.0	3.1	5.4	0.036*	0.8	2.3	5.4
HB+10, VL-8	0.3	1.4	3.8	0.5	1.9	4.4	N/S	0.3	1.6	4.4
HB+10, VL-10	0.3	1.5	4.7	0.3	1.5	4.3	N/S	0.3	1.5	4.7

*Significant difference $P < .05$. **Highly significant difference $P < .01$, N/S indicates not significant

ment ranged from 0.06 mm to 0.30 mm with a mean of 0.19 mm.

Statistical analysis

The nonparametric Wilcoxon signed rank test was used to compare the bone thickness on the right and left. The Mann-Whitney U test was used to compare the males and females. Results with $P < .05$ were regarded as statistically significant.

Results

The minimum, median, and maximum values of the

right and left sides and statistical difference are summarized in Table 1. Ten out of the 35 locations showed significant differences between the right and left sides, but the differences were less than 1.0 mm. The thickest point (HB+10, AP-2) was 5.0 mm on average, and the thinnest point was 1.1 mm. Figure 2 shows the color maps of the median values of bone thicknesses in the infrazygomatic crest and the zygomatic process of the maxilla. There was a tendency for the bone to be thicker at the superior and lateral area of the zygomatic process of the maxilla than the other locations.

Table 2 shows the bone thickness in the male and female groups. Six out of 35 locations showed significant




Fig. 2. Mean of the bilateral bone thickness (mm) map of the infrazygomatic crest area of the zygomatic process of the maxilla. X indicates the bone thickness at each location.




Fig. 3. Male and female bone thickness (mm) maps. X indicates the bone thickness at each point.

differences between the male and female groups. Generally, the bone thickness of the male group was greater than that of the female group. Figure 3 displays color maps of the median values of bone thickness in the male and female groups.

Discussion

In the present study, the FH plane was used as a horizontal plane from which to measure the bone thickness. For the consistent measurement of bone thickness, CT images were taken in parallel with the FH plane of the subjects. For the consistency and convenience of measurement, the binary image setting (window width: 0 HU, window level: 500 HU) was used. A pilot study had been carried out to determine the appropriate window level of the

binary image setting. A CT had been taken with a cortical bone slice of bovine rib attached by periosteum. At the binary image of the window level of 500 HU, the bone thickness measured on the CT image was close to that measured at the cortical bone. Therefore, the binary image setting of the window level of 500 HU was used in this study.

The zygomaticomaxillary suture had been considered to be the reference area for constructing the horizontal planes and vertical lines; however, it was not clearly identified on the CT image in some children due to ossification. On the other hand, the most anterior point of infratemporal fossa was easily identifiable on the CT images on all of the horizontal planes, and these points were observed to be quite close to the zygomaticomaxillary suture.

The thickest point (5.0 mm) was at the point HB+10 and AP-2, which corresponded to the zygomatic process of the maxilla, and the thinnest point (1.1 mm) was at the point HB+2 and AP-10, which corresponded to the anterior wall of the maxillary sinus. The bone at AP-2 and AP-4 in all of the horizontal planes tended to be thicker than that at more medial points, and the bone thickness of these areas ranged from 1.8 mm to 5.0 mm, which was regarded as the appropriate thickness for the miniplate placement. Considering the thickness of the miniplate (2 mm), miniscrews of 4 to 7 mm in length can be selected. If oblique installation of the miniscrew is possible, a longer miniscrew could be selected for better bone support.

According to our findings, the bone thickness in the growing patients tended to be thicker at the superior and lateral areas of the zygomatic process of the maxilla. This might be related to the development of the maxillary sinus. In previous reports,¹⁷⁻²⁰ the maxillary sinus continued to extend both laterally and inferiorly from the medial orbital wall after birth. Inferiorly, it reached the level of the hard palate at 9 years of age and continued to grow downward and together with the pneumatization of the maxillary alveolar bone, reaching the level of the nasal floor at 12 years of age. The floor of the maxillary sinus ended up extending 4-5 mm inferior to the nasal floor. The shape of the maxillary sinus also changed into a reverse pyramidal shape with the lateral expansion at the superior side.

Our results showed no differences between the right and left side except at 10 out of the 35 locations evaluated, and these differences were very small. The reason might be that the patients included in the present study had no facial asymmetry and the measurements were taken at symmetric locations. Previous studies have reported that the right and left side of the maxillary sinus showed no

Table 2. The minimum (Min), median, maximum (Max) values of the bone thickness in male and female (unit: mm)

Points	n	Male			n	Female			P value
		Min	Median	Max		Min	Median	Max	
HB-2, VL-2	13	0.3	4.2	6.2	18	1.2	2.8	5.8	N/S
HB-2, VL-4	13	0.3	1.8	5.8	18	1.1	1.9	4.8	N/S
HB-2, VL-6	13	0.3	1.3	3.5	18	1.1	1.5	5.0	N/S
HB-2, VL-8	13	0.3	1.2	2.1	16	0.6	1.2	3.1	N/S
HB-2, VL-10	13	0.3	1.2	2.0	16	0.3	1.0	2.7	N/S
HB, VL-2	14	0.5	3.6	6.4	18	0.7	3.1	7.4	N/S
HB, VL-4	14	0.5	3.2	6.0	18	0.5	2.4	6.9	N/S
HB, VL-6	14	0.5	2.0	5.4	18	0.6	1.7	6.2	N/S
HB, VL-8	14	0.5	1.6	5.6	18	0.8	1.4	4.5	N/S
HB, VL-10	14	0.5	1.0	4.3	17	0.8	1.2	3.4	N/S
HB+2, VL-2	14	1.6	3.9	7.6	18	1.8	3.6	8.4	N/S
HB+2, VL-4	14	0.8	2.8	8.8	18	1.3	2.1	7.2	N/S
HB+2, VL-6	14	0.3	1.4	8.9	18	0.7	1.6	4.7	N/S
HB+2, VL-8	14	0.3	1.2	4.5	18	0.6	1.4	3.4	N/S
HB+2, VL-10	14	0.3	1.1	2.2	18	0.3	1.1	3.0	N/S
HB+4, VL-2	14	2.1	4.2	9.6	18	1.5	3.3	10.6	N/S
HB+4, VL-4	14	1.2	2.9	8.8	18	0.9	2.0	5.9	N/S
HB+4, VL-6	14	0.7	1.8	5.6	18	0.7	1.6	4.3	N/S
HB+4, VL-8	14	0.7	1.2	4.3	18	0.5	1.4	3.5	N/S
HB+4, VL-10	14	0.3	1.2	2.0	18	0.3	1.2	2.5	N/S
HB+6, VL-2	14	3.1	4.7	10.8	18	2.0	3.6	7.7	0.030*
HB+6, VL-4	14	1.6	3.4	7.3	18	0.9	2.2	6.0	0.032*
HB+6, VL-6	14	0.8	2.1	5.6	18	0.8	1.7	4.4	N/S
HB+6, VL-8	14	0.8	1.4	3.4	18	0.7	1.6	3.4	N/S
HB+6, VL-10	14	0.3	1.2	2.3	18	0.3	1.2	2.6	N/S
HB+8, VL-2	14	3.7	5.3	10.9	18	1.5	3.9	9.4	0.011*
HB+8, VL-4	14	1.5	3.9	6.8	18	1.1	3.0	6.0	N/S
HB+8, VL-6	14	1.0	2.5	5.1	18	0.6	1.8	4.6	N/S
HB+8, VL-8	14	0.8	1.6	3.5	18	0.5	1.5	3.1	N/S
HB+8, VL-10	14	0.3	1.3	3.1	17	0.5	1.3	2.3	N/S
HB+10, VL-2	14	4.1	5.4	9.5	18	1.5	4.4	9.2	0.023*
HB+10, VL-4	14	2.2	4.4	10.6	18	0.9	3.2	7.4	0.018*
HB+10, VL-6	14	1.2	3.1	5.4	18	0.8	2.0	4.6	0.042*
HB+10, VL-8	14	0.3	1.9	4.4	18	0.5	1.6	3.1	N/S
HB+10, VL-10	13	0.3	1.5	3.3	18	0.5	1.5	4.7	N/S

*Significant difference $P < .05$, **Highly significant difference $P < .01$, N/S indicates not significant

significant differences in the transverse width, antero-posterior width, height, or volume.^{17,18}

Our results showed that only 6 out of 35 locations demonstrated a statistically significant gender difference. This is in concordance with many previous studies showing no significant difference in maxillary sinus development between males and females.¹⁷⁻²⁰ However, there was a tendency for the bone to be thicker in males than females in the present study. This was probably related to the gender difference in occlusal forces because a high concentration of occlusal stress at the zygomatic arch might be related with thick and dense cortical bone of this area, with bony adaptation to the functional load.²¹

In some younger children with mixed dentition in the present study, the tooth germs or root apex of the permanent teeth were observed under HB-2. Therefore, the area superior to HB-2 seemed to be appropriate for placement of miniplates in children with mixed dentition. In a previous study, the distance from the apex of the mesio-buccal root of the maxillary second molar to the buccal bone plate was 4.63 mm.²¹ However, no study has evaluated the buccal bone thickness to the tooth germs in children with mixed dentition. This should be investigated in a future study.

The patients in the present study were diagnosed with skeletal Class III malocclusion with maxillary hypo-

plasia. There might be differences in the bone thicknesses at the infrazygomatic crest area among skeletal Class I, II, and III patients. Further studies are needed to investigate the bone thickness with various skeletal patterns.

The principal conclusions can be summarized as follows. In the infrazygomatic crest area, the superior and lateral area of the zygomatic process of the maxilla are most appropriate for placement of a miniplate for orthodontic treatment in growing children with skeletal class III malocclusion and a retruded maxilla.

There was a tendency for the bone to be thicker at the superior and lateral area of the zygomatic process of the maxilla. There was no clinically significant difference in bone thickness between the right and left sides; however, the bone was thicker in the males than the females.

References

1. De Pauw GA, Dermaut L, De Bruyn H, Johansson C. Stability of implants as anchorage for orthopedic traction. *Angle Orthod* 1999; 69: 401-7.
2. Irie M, Nakamura S. Orthopedic approach to severe skeletal Class III malocclusion. *Am J Orthod* 1975; 67: 377-92.
3. Kambara T. Dentofacial changes produced by extraoral forward force in the *Macaca irus*. *Am J Orthod* 1977; 71: 249-77.
4. Revelo B, Fishman LS. Maturational evaluation of ossification of the midpalatal suture. *Am J Orthod Dentofacial Orthop* 1994; 105: 288-92.
5. Smalley W, Shapiro PA, Hohl TH, Kokich VG, Brånemark PI. Osseointegrated titanium implants for maxillofacial protraction in monkeys. *Am J Orthod Dentofacial Orthop* 1988; 94: 285-95.
6. Roberts WE, Helm FR, Marshall KJ, Gongloff RK. Rigid endosseous implants for orthodontic and orthopedic anchorage. *Angle Orthod* 1989; 59: 247-56.
7. Singer SL, Henry PJ, Rosenberg I. Osseointegrated implants as an adjunct to facemask therapy: a case report. *Angle Orthod* 2000; 70: 253-62.
8. Hong H, Ngan P, Han G, Qi LG, Wei SH. Use of onplants as stable anchorage for facemask treatment: a case report. *Angle Orthod* 2005; 75: 453-60.
9. Cha BK, Lee NK, Choi DS. Maxillary protraction treatment of skeletal Class III children using miniplate anchorage. *Korean J Orthod* 2007; 37: 73-84.
10. Cha BK, Choi DS, Ngan P, Jost-Brinkmann PG, Kim SM, Jang IS. Maxillary protraction with miniplates providing skeletal anchorage in a growing Class III patient. *Am J Orthod Dentofacial Orthop* 2011; 139: 99-112.
11. Cha BK, Ngan PW. Skeletal anchorage for orthopedic correction of growing Class III patients. *Semin Orthod* 2011; 17: 124-37.
12. Sherwood KH, Burch J, Thompson W. Intrusion of supererupted molars with titanium miniplate anchorage. *Angle Orthod* 2003; 73: 597-601.
13. Bengi AO, Karacay S, Akin E, Olmez H, Okçu KM, Mermut S. Use of zygomatic anchors during rapid canine distalization: a preliminary case report. *Angle Orthod* 2006; 76: 137-47.
14. De Clerck H, Geerinckx V, Siciliano S. The zygoma anchorage system. *J Clin Orthod* 2002; 36: 455-9.
15. Seebeck J, Goldhahn J, Städele H, Messmer P, Morlock MM, Schneider E. Effect of cortical thickness and cancellous bone density on the holding strength of internal fixator screws. *J Orthop Res* 2004; 22: 1237-42.
16. Molly L. Bone density and primary stability in implant therapy. *Clin Oral Implants Res* 2006; 17: Suppl 2: 124-35.
17. Park CH, Kim KD, Park CS. Measurement of maxillary sinus volume using computed tomography. *Korean J Oral Maxillofac Radiol* 2000; 30: 63-70.
18. Jun BC, Song SW, Park CS, Lee DH, Cho KJ, Cho JH. The analysis of maxillary sinus aeration according to aging process; volume assessment by 3-dimensional reconstruction by high-resolution CT scanning. *Otolaryngol Head Neck Surg* 2005; 132: 429-34.
19. Lee CH, Rhee CS, Oh SJ, Jun YH, Min YG, Kim IO. Development of the paranasal sinuses in children: a MRI study. *Korean J Otolaryngol-Head Neck Surg* 2000; 43: 507-13.
20. Choi DS, Cha BK, Jang I, Kang KH, Kim SC. Three-dimensional finite element analysis of occlusal stress distribution in the human skull with premolar extraction. *Angle Orthod* 2013; 83: 204-11.
21. Jin GC, Kim KD, Roh BD, Lee CY, Lee SJ. Buccal bone plate thickness of the Asian people. *J Endod* 2005; 31: 430-4.