

Molecular Detection of *Ancylostoma duodenale*, *Ancylostoma ceylanicum*, and *Necator americanus* in Humans in Northeastern and Southern Thailand

Issarapong Phosuk^{1,2}, Pewpan M. Intapan^{1,2,*}, Tongjit Thanchomnang^{1,3}, Oranuch Sanpool^{1,2}, Penchom Janwan^{1,2}, Pornpip Laummaunwai^{1,2}, Witthaya Aamnat⁴, Nimit Morakote⁵ and Wanchai Maleewong^{1,2}

¹Department of Parasitology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand; ²Research and Diagnostic Center for Emerging Infectious Diseases, Khon Kaen University, Khon Kaen 40002, Thailand; ³Department of Medicine, Mahasarakham University, Mahasarakham 44150, Thailand; ⁴Department of Medical Technology, School of Allied Health Sciences and Public Health, Walailak University, Thasala, Nakhon Si Thammarat 80160, Thailand; ⁵Department of Parasitology, Faculty of Medicine Chiang Mai University, Chiang Mai 50200, Thailand

Abstract: The 2 principal species of hookworms infecting humans are *Necator americanus* and *Ancylostoma duodenale*. Case studies on zoonotic hookworm infections with *Ancylostoma ceylanicum* and/or *Ancylostoma caninum* are known mainly from Asian countries. Of these 2 zoonotic species, only *A. ceylanicum* can develop to adulthood in humans. In the present study, we report a molecular-based survey of human hookworm infections present in southern and northeastern Thailand. Thirty larval hookworm samples were obtained from fecal agar plate cultures of 10 patients in northeastern Thailand and 20 in southern Thailand. Partial ITS1, 5.8S, and ITS2 regions of the ribosomal DNA genes were amplified using PCR. The amplicons were sequenced, aligned, and compared with other hookworm sequences in GenBank database. The results showed that, in Thailand, *N. americanus* is more prevalent than *Ancylostoma* spp. and is found in both study areas. Sporadic cases of *A. ceylanicum* and *A. duodenale* infection were seen in northeastern Thailand.

Key words: *Ancylostoma duodenale*, *Ancylostoma ceylanicum*, *Necator americanus*, human, molecular detection

Human hookworm infections commonly cause socioeconomic and public health problems, with approximately 1 billion persons infected worldwide [1]. Hookworm infections cause iron deficiency anemia, resulting in mental retardation and growth insufficiency in children [2]. *Ancylostoma duodenale* and *Necator americanus* are the 2 common species that cause human infections [3]. The former is common in the Middle East, Northern Africa, India, Australia, and Europe, while *N. americanus* is widespread in the western hemisphere, sub-Saharan Africa, eastern Asia, and southeast Asia [4]. In addition, zoonotic hookworms such as *Ancylostoma ceylanicum*, *Ancylostoma braziliense*, and *Ancylostoma caninum* have been reported as potentially significant public health threats in many areas [5]. Hookworm infections are still highly prevalent in Thailand

[6,7] and are generally diagnosed by finding larvae/eggs in fecal preparations. Morphological identification of hookworm larvae to species is difficult and molecular identification is a useful tool in this regard. Zoonotic hookworm disease caused by *A. ceylanicum* was detected by copro-molecular methods in central Thailand [6,8] and Lao PDR [9]. A molecular approach to identify the causative species in other parts of Thailand is still lacking. Here, we report molecular identification of hookworm species that infect humans in the northeastern (NE) and southern Thailand. This genetic data is important as a part of continuing investigations into epidemiology of hookworms in Thailand.

The study was conducted between 2011 and 2013. Hookworm larvae were collected from 30 fecal samples using the agar plate culture technique [10]. Ten fecal samples were collected from patients in Srinagarind Hospital, Faculty of Medicine, Khon Kaen University, NE Thailand and 20 samples from rural villagers, Nakhon Si Thammarat, southern Thailand. A single larval hookworm from each fecal sample was kept in 95% ethanol until used for DNA extraction. Oral or written

•Received 30 May 2013, revised 16 September 2013, accepted 11 October 2013.

*Corresponding author (pewpan@kku.ac.th)

© 2013, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

informed consent was obtained from patients and legal guardians. This study was approved by the Khon Kaen University Ethics Committee for Human Research (HE551247), Khon Kaen, Thailand.

DNA was extracted separately from each larva using a Genomic DNA Mini Kit (Macherey-Nagel GmbH & Co., Düren, Germany) according to the manufacturer's instructions. The partial ITS1, full length 5.8S gene, and partial ITS2 ribosomal DNA regions were amplified from each larva using PCR and the primers RTHW1F (forward): 5'-GAT GAG CAT TGC WIG AAT GCC G-3' and RTHW1R (reverse): 5'-GCA AGT RCC GTT CGA CAA ACA G-3' [8]. PCR conditions were as follows; initial denaturation at 94°C for 5 min, followed by 35 cycles of 95°C for 30 sec (denaturation), 65°C for 30 sec (annealing), and 72°C for 30 sec (extension), and a final extension at 72°C for 10 min. The reaction was carried out in a 25 µl volume containing PCR 1x FastStart High Fidelity Reaction Buffer (Roche Applied Science, Mannheim, Germany), 1.8 mM MgCl₂, 0.2 mM of each deoxyribonucleotide triphosphate, 0.2 µM of each primer, and 0.625 units of FastStart High Fidelity Enzyme Blend (Roche Applied Science), respectively. The PCR product was run on a 1% agarose gel, cut out, and purified for DNA sequencing, which was performed using the MegaBACE™ 1000 DNA Analysis System (GE Healthcare, Piscataway, New Jersey, USA). The specific primers above were used as sequencing primers. The nucleotide sequences were analyzed by BLAST-N search via NCBI and the DNA alignment using Clustal-W [11].

Amplicon sizes were approximately 485 bp (typical of *N. americanus*) or 380 bp (typical of *Ancylostoma* spp.). Sequences showed extremely high similarities (99-100%) with hookworm sequences in the GenBank database. Of the 10 samples from NE Thailand, 6 were *N. americanus* (Khon Kaen, n=3; Mukdahan, n=1; Roi ET, n=1; Loei, n=1) (different from AF217891 at a single base), 3 were *A. ceylanicum*, (Khon Kaen, n=2; Mahasarakham, n=1) (identical with AB501355), and 1 was *A. duodenale* (Loei) different from AB501348 at 2 bases). All 20 samples from Nakhon Si Thammarat, southern Thailand, were *N. americanus* (different from AF217891 at a single base) (n=20). All sequences obtained in this study were deposited in the GenBank data base under the accession no. KC896796-KC896825 (Table 1).

In the present study, molecular analysis was used to confirm human infections with 2 species of human hookworms, namely, *N. americanus* and *A. duodenale* found in NE and southern Thailand. In addition, 1 species of animal hookworm, namely,

Table 1. Hookworm sequences deposited in Genbank database

Parts of Thailand	Provinces	Sequence ID	Accession no.	Hookworm species
Northeast-ern	Khon Kaen	THA FHWKK4	KC896796	<i>N. americanus</i>
		THA FHWKK6	KC896798	<i>A. ceylanicum</i>
		THA FHWKK7	KC896799	<i>N. americanus</i>
		THA FHWKK9	KC896801	<i>N. americanus</i>
		THA HWKK3	KC896805	<i>A. ceylanicum</i>
	Loei	THA FHWKK8	KC896800	<i>A. duodenale</i>
		THA HWKK1	KC896803	<i>N. americanus</i>
	Mahasarakham	THA FHWKK5	KC896797	<i>A. ceylanicum</i>
		Roi Et	THA FHWKK10	KC896802
	Mukdahan	THA HWKK2	KC896804	<i>N. americanus</i>
Southern	Nakhon Si Thammarat	THA HWS1	KC896806	<i>N. americanus</i>
		THA HWS2	KC896807	<i>N. americanus</i>
		THA HWS4	KC896808	<i>N. americanus</i>
		THA HWS5	KC896809	<i>N. americanus</i>
		THA HWS6	KC896810	<i>N. americanus</i>
		THA HWS7	KC896811	<i>N. americanus</i>
		THA HWS8	KC896812	<i>N. americanus</i>
		THA HWS9	KC896813	<i>N. americanus</i>
		THA HWS10	KC896814	<i>N. americanus</i>
		THA HWS11	KC896815	<i>N. americanus</i>
		THA HWS12	KC896816	<i>N. americanus</i>
		THA HWS13	KC896817	<i>N. americanus</i>
		THA HWS15	KC896818	<i>N. americanus</i>
		THA HWS16	KC896819	<i>N. americanus</i>
THA HWS17	KC896820	<i>N. americanus</i>		
THA HWS18	KC896821	<i>N. americanus</i>		
THA HWS19	KC896822	<i>N. americanus</i>		
THA HWS20	KC896823	<i>N. americanus</i>		
THA HWS21	KC896824	<i>N. americanus</i>		
THA HWS22	KC896825	<i>N. americanus</i>		

A. ceylanicum, was found in NE Thailand. In NE Thailand, *N. americanus* was the main hookworm identified, but *A. duodenale* and *A. ceylanicum* were also found. In southern Thailand, only *N. americanus* was detected. Our results complement previous reports from the central part of Thailand [6,7]. A survey of gastrointestinal parasites of dogs and humans in communities in Bangkok revealed *A. ceylanicum* and *A. caninum* in dogs and *N. americanus* and *A. ceylanicum* in humans [6]. Recently, a cohort study to identify the incidence and risk factors of hookworm infections was conducted in a rural community, central Thailand. *N. americanus* was the most common hookworm identified there. *A. duodenale* and *A. ceylanicum* were also detected [7]. We confirmed that the principal hookworm species infecting humans in Thailand is *N. americanus*. However, *A. duodenale* and *A. ceylanicum* infections were also detected in NE Thailand. The finding of the predominance of *N. ameri-*

canus in southern Thailand is consistent with the study in Peninsular Malaysia [12], where *N. americanus* was more common than *A. ceylanicum*, whereas *A. duodenale* infection was not found. In contrast, a study in Lao PDR found that *A. duodenale*, and the animal hookworms, *A. caninum* and *A. ceylanicum* were slightly more prevalent than *N. americanus* [8]. Thailand, Lao PDR, and Malaysia are neighboring countries. The geographical differences in the species of hookworms causing human infections might possibly be associated with parasite behavior, ethnicity, climate, temperature, and environmental factors [13,14].

In conclusion, our study revealed evidence of both human and animal hookworms among people in various areas of Thailand.

ACKNOWLEDGMENTS

This research was funded by grants from the Higher Education Research Promotion and National Research University Project of Thailand, Office of the Higher Education Commission, Thailand; the Research and Diagnostic Center for Emerging Infectious Diseases and the Faculty of Medicine, Khon Kaen University, Thailand. Issarapong Phosuk was supported by the Khon Kaen University grant. Wanchai Maleewong and Pewpan M. Intapan were supported by TRF Senior Research Scholar Grant, Thailand Research Fund grant number RTA5580004. We wish to acknowledge the support of the Khon Kaen University Publication Clinic, Research and Technology Transfer Affairs, Khon Kaen University, Thailand for their assistance.

REFERENCES

- Schneider B, Jariwala AR, Periago MV, Gazzinelli MF, Bose SN, Hotez PJ, Diemert DJ, Bethony JM. A history of hookworm vaccine development. *Hum Vaccin* 2011; 7: 1234-1244.
- Crompton DW. The public health importance of hookworm disease. *Parasitology* 2000; 121 (suppl): 39-50.
- Chan MS, Medley GF, Jamison D, Bundy DA. The evaluation of potential global morbidity attributable to intestinal nematode infections. *Parasitology* 1994; 109: 373-387.
- de Silva NR, Brooker S, Hotez PJ, Montresor A, Engels D, Savioli L. Soil-transmitted helminth infections: updating the global picture. *Trends Parasitol* 2003; 19: 547-551.
- Mahdy MA, Lim YA, Ngui R, Siti Fatimah MR, Choy SH, Yap NJ, Al-Mekhlafi HM, Ibrahim J, Surin J. Prevalence and zoonotic potential of canine hookworms in Malaysia. *Parasit Vectors* 2012; 5: 88.
- Traub RJ, Inpankaew T, Sutthikornchai C, Sukthana Y, Thompson RC. PCR-based coprodiagnostic tools reveal dogs as reservoirs of zoonotic ancylostomiasis caused by *Ancylostoma ceylanicum* in temple communities in Bangkok. *Vet Parasitol* 2008; 155: 67-73.
- Jex AR, Lim YA, Bethony JM, Hotez PJ, Young ND, Gasser RB. Soil-transmitted helminths of humans in Southeast Asia--towards integrated control. *Adv Parasitol* 2011; 74: 231-265.
- Jiraanankul V, Aphijirawat W, Mungthin M, Khositnithikul R, Rangsin R, Traub RJ, Piyaraj P, Naaglor T, Taamasri P, Leelayoova S. Incidence and risk factors of hookworm infection in a rural community of central Thailand. *Am J Trop Med Hyg* 2011; 84: 594-598.
- Sato M, Sanguankiat S, Yoonuan T, Pongvongsa T, Keomoungkhoun M, Phimmayoi I, Boupa B, Moji K, Waikagul J. Copromolecular identification of infections with hookworm eggs in rural Lao PDR. *Trans R Soc Trop Med Hyg* 2010; 104: 617-622.
- Arakaki T, Iwanaga M, Kinjo F, Saito A, Asato R, Ikeshiro T. Efficacy of agar plate culture in detection of *Strongyloides stercoralis* infection. *J Parasitol* 1990; 76: 425-428.
- Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 1994; 22: 4673-4680.
- Ngui R, Ching LS, Kai TT, Roslan MA, Lim YA. Molecular identification of human hookworm infections in economically disadvantaged communities in Peninsular Malaysia. *Am J Trop Med Hyg* 2012; 86: 837-842.
- Beaver PC, Jung RC, Cupp EW. *Clinical Parasitology*. 9th ed. Philadelphia, USA. Lea & Febiger. 1984, p 269-287.
- Hoagland KE, Schad GA. *Necator americanus* and *Ancylostoma duodenale*: life history parameters and epidemiological implications of two sympatric hookworms of humans. *Exp Parasitol* 1978; 44: 36-49.