

The chronology of second molar development in Brazilians and its application to forensic age estimation

Manuella Santos Carneiro Almeida, Andrea dos Anjos Pontual*, Rejane Targino Beltrão, Ricardo Villar Beltrão, Maria Luiza dos Anjos Pontual**

Department of Clinic and Social Dentistry, School of Dentistry, Federal University of Paraíba, João Pessoa, Brazil

**Department of Oral Medicine, Division of Oral Radiology, Pernambuco Dental School, Pernambuco University-UPE, Camaragibe, Brazil*

***Department of Clinical and Social Dentistry, Division of Oral Radiology, Federal University of Pernambuco, Recife, Brazil*

ABSTRACT

Purpose: This study investigated the possible correlation between the mineralization of the second molars and the chronological age of a sample population of the state of Paraíba, Brazil.

Materials and Methods: One thousand eight hundred fifty-four digitized panoramic images using a scanner of a private dental radiology service were obtained in six months. Of these, 457 were selected from patients aged 4.6 to 16 years who met certain criteria. The images were assessed twice by a radiologist with 5 years experience. A minimum interval of 30 days between the evaluations of the same patient was established. To analyze the relationship between chronological age, calcification level proposed by Demirjian et al in 1973, gender, and tooth, a multiple linear regression model was adjusted, taking age as the response variable ($p < 0.05$).

Results: The gender and calcification level were significantly correlated with the age of the patients ($p < 0.05$). There was a significant difference between the average age of the patients' upper teeth compared to the lower ones in both genders ($p < 0.05$). The dental development occurred earlier in female individuals than in male ones, and there was no significant difference between the right and left second permanent molars.

Conclusion: It was observed that ethnic variables are related to certain parameters of age and sex identification in the Brazilian population, providing important information for forensic evaluations. (*Imaging Sci Dent 2013; 43 : 1-6*)

KEY WORDS: Tooth Calcification; Molar; Age Determination by Teeth; Radiography, Panoramic

Introduction

Biological and chronological ages are not completely associated with body development. Therefore, parameters are obtained to evaluate biological development such as dental age, bone age, mental age, and other factors such as menarche, voice change, height, and weight.¹ Dental age assessment plays an important role in forensic medicine, pediatric dentistry, and orthodontic treatment planning.²

Several methods of age estimation have been suggested, and the method by Demirjian et al³ is the most frequently used.^{4,5}

Demirjian's method is based on eight stages of tooth calcification ranging from crown and root formation to apex closure of the seven left permanent mandibular teeth. The score of each stage is allocated, and the sum of the scores gives an evaluation of the subject's dental maturity. These scores can be converted into a dental age by means of available tables and percentile curves from the original study. The difference between the dental and chronological age indicates an advancement or delay in dental maturity.⁶

Estimating dental maturity according to percentile charts has been shown to be more suitable in determining whether

Received October 13, 2012; Revised November 10, 2012; Accepted November 20, 2012
Correspondence to : Dr. Manuella Santos Carneiro Almeida
Department of Oral Diagnosis, Federal University of Paraíba, Campus I, João Pessoa
58035140, Brazil
(Tel) 55-83-3245-4681, (Fax) 55-83-3218-6782, (E-mail) manuellaalmeida@hotmail.com

the dental maturity of a subject with a known age is advanced or delayed rather than to predict an unknown age.^{4,7,8} Demirjian's study was conducted from the data obtained from a group of French Canadian boys and girls. Thereafter, several studies have shown that the estimation of dental age of another population using Demirjian's standards developed for French Canadians was not suitable for their study populations and the ethnic differences of dental maturity had to be taken into account.⁹⁻¹³

The establishment of population-specific standards for third molars is considered important for enhancing the accuracy of age estimation based on the development of these teeth. However, third molars are often missing congenitally or have been extracted, so they cannot be depended upon to indicate age. Moreover, if they are malformed or malpositioned, age estimation may be difficult.^{7,14} To overcome these limitations, the second molars can be analyzed as an age indicator. Many studies have investigated the accuracy of forensic age estimation based on the chronology of second molar development and have concluded that it can be considered a valuable age indicator in populations.^{7,14,15}

Thevissen et al¹⁶ insisted that dental age estimation, especially for young individuals, should be based on the data collected from the appropriate population group. Thus, the aim of this study was to evaluate the chronology of second molar development in Brazilians and suggest its application to forensic age estimation.

Materials and Methods

This study was approved by the Center for Health Sciences Ethics Committee in Human Research of Federal University of Paraíba, Brazil (0178/2009). A cross-sectional analysis was performed for a sample of 457 panoramic radiographs taken from 192 males and 265 females aged between 4.6 and 16 years. A convenience sampling method was used, and the panoramic radiographs had been obtained from July to December 2010 from a private dental radiology service in Paraíba, Brazil. A sample of 457 out of a total of 1,854 panoramic radiographs was selected based on the following criteria: the analysis was restricted to individuals of both genders aged 3-16 years and born in the state of Paraíba, Brazil. Unreadable radiographs, or those showing hypodontia or gross pathologic problems, were excluded.

The panoramic radiographs were taken using a Rotograph Plus panoramic machine (Villa Sistemi Medicali, Milan, Italy), and suitable exposure factors considering the patients'

ages and sizes were applied. The Kodak T-Mat G/RA X-ray film (Eastman Kodak, Rochester, NY, USA) were manually processed according to temperature/time tables using fresh Kodak concentrated processing solutions (Eastman Kodak, Rochester, NY, USA). The radiographs were digitized using a Hewlett-Packard scanner (Hewlett-Packard Co., Palo Alto, CA, USA) equipped with a transparency reader as part of routine clinical activity, with a resolution of 300 dots per inch (dpi). All the images were saved and stored in Tagged Image File Format (TIFF). Each digitized panoramic image was coded with a numerical ID. The examinations were assessed twice by a radiologist with 5 years of experience. Then, all of the images were exported to the computer and viewed, randomly, in a quiet and darkened room with the aid of a 17-inch LCD monitor (HP Compaq LA1751 g, Hewlett-Packard Co., Palo Alto, CA, USA) using the Windows Picture and Fax Viewer (Microsoft, Redmond, WA, USA).

The age and gender of the subjects were thus unknown to the observer. Prior to the examination, verbal and practical instructions and calibration tests were performed. The evaluation method was explained, and training and knowledge of the evaluators were verified. A minimum interval of 30 days between the evaluations of the same patient was established. If the first and second assessments differed, the readings were repeated until a consistent result was obtained. All of the second molars were assessed. For the analysis of the degree of calcification, Demirjian's method (Fig. 1) and a form was filled out for each panoramic radiograph were used. The examiner could use the zoom tool whenever necessary to obtain a maximum magnification of 3 times to avoid loss of resolution. At most, 10 panoramic images were examined each day to avoid compromising the assessments due to eyestrain.

The data were analyzed using Stata[®] 9.2 (StataCorp, Col-




Fig. 1. Classification of the degree of mineralization of molar teeth according to Demirjian et al³

Table 1. Multilinear regression models adjusted for each second molar to predict the age (in months) of patients in the sample based on gender and Demirjian’s score for the second molars of the right maxilla, left maxilla, left mandible, and right mandible

Tooth	Explanatory variable*	B coefficient	SD	P
Right maxillary second molar	Constant	74.3*	6.44	< 0.001
	Demirjian’s stage	16.2	0.46	< 0.001
	Gender	4.3	1.52	0.005
	$R^2=73.2\%$			
Left maxillary second molar	Constant	74.9*	2.11	< 0.001
	Demirjian’s stage	16,0	0.46	< 0.001
	Gender	4.4	1.54	0.005
	$R^2=72.8\%$			
Left maxillary second molar	Constant	62.9*	2.55	< 0.001
	Demirjian’s stage	18.0	0.54	< 0.001
	Gender	3.7	1.59	0.021
	$R^2=70.6\%$			
Right maxillary second molar	Constant	62.2*	2.56	< 0.001
	Demirjian’s stage	18.3	0.55	< 0.001
	Gender	3.0	1.59	0.050
	$R^2=70.8\%$			

*: Estimated age (in months) for Demirjian’s stage of the second molars in female individuals, SD: Standard deviation

lege Station, TX, USA) and Minitab® 15 (Minitab Inc., State College, MA, USA) software. To evaluate the relationship between age, the calcification level proposed by Demirjian, gender, and tooth, a multiple linear regression model was adjusted taking age into consideration as the response variable. In all the tests, a 0.05 significance level was adopted.

Results

The patients’ ages ranged from 4.66 to 16 years (11.7 ± 2.5 years). A total of 192 (42%) panoramic radiographs were taken from male individuals, and 265 (58%) were taken from female individuals.

Table 1 presents the results of the multilinear regression models adjusted for each second molar to predict the age (in months) of patients in the sample based on the gender and Demirjian’s score for each second molar. There was a statistically significant difference between the average ages of the patients in the different stages of calcification of second molars according to the gender ($p < 0.05$). The results of the regression models corresponding to the right maxillary second molar and left maxillary second molar




Fig. 2. Representation of a linear regression model for predicting the age by gender, upper second molars, and Demirjian’s stage of calcification




Fig. 3. Representation of a linear regression model for predicting the age by gender, lower second molars, and Demirjian’s stage of calcification

were found to be identical. The same was observed for the left mandibular second molar and right mandibular second molar. There were no second molars in the sample of patients in stage A of calcification; thus, the first stage of calcification found was stage B (B coefficient). For the upper second molar teeth, stage B had a mean age of 76.6 months, and the lower ones, an average of 65.5 months old. In the upper teeth, there was a variation of 16.1 months for the subsequent stage and in the lower teeth, an interval of 18.1 months.

Figures 2 and 3 present the results of the multilinear regression models adjusted to predict the age of patients in the sample based on gender and the second upper and lower molars, respectively. There was a significant association between age and Demirjian’s calcification stages

Table 2. Descriptive summary of the ages (in months) and standard deviation in the sample, according to Demirjian’s scores, gender, and upper second molars

Stage	N	Right maxillary second molars				Left maxillary second molars			
		Male	N	Female	N	Male	N	Female	
B	4	79.5 ± 14.4	4	64.8 ± 6.0	5	82.2 ± 13.8	5	66.8 ± 6.9	
C	8	94.5 ± 21.6	10	87.0 ± 13.4	7	94.7 ± 23.4	9	89.6 ± 12.8	
D	36	114.3 ± 17.1	48	107.0 ± 17.4	36	114.0 ± 17.1	50	106.9 ± 17.6	
E	30	125.4 ± 12.2	35	125.0 ± 17.4	29	125.6 ± 12.0	31	126.2 ± 17.4	
F	35	139.5 ± 14.4	41	138.7 ± 15.3	36	139.3 ± 14.3	43	137.9 ± 15.5	
G	35	158.9 ± 17.4	61	155.6 ± 14.1	36	159.5 ± 17.6	60	156.0 ± 13.8	
H	44	177.2 ± 13.5	66	170.5 ± 19.3	43	177.1 ± 13.7	67	169.9 ± 19.8	

N: Number of samples

Table 3. Descriptive summary of the averages of the ages (in months) and standard deviation in the sample, according to Demirjian’s scores, gender, and lower second molars

Stage	N	Right mandibular second molars				Left mandibular second molars			
		Male	N	Female	N	Male	N	Female	
B	1	60.0	0	—	1	60.0	2	62.5 ± 9.2	
C	6	93.8 ± 23.9	8	77.3 ± 17.1	6	93.8 ± 23.9	6	82.2 ± 16.6	
D	14	96.1 ± 12.4	32	103.5 ± 22.4	11	98.3 ± 17.5	25	98.2 ± 18.2	
E	48	121.4 ± 14.2	43	113.2 ± 16.5	50	119.1 ± 15.0	55	116.4 ± 18.8	
F	38	138.8 ± 13.5	46	136.4 ± 13.6	39	138.7 ± 13.3	44	137.1 ± 13.9	
G	40	154.5 ± 21.4	68	152.9 ± 15.0	41	154.2 ± 20.8	68	154.0 ± 15.6	
H	45	176.6 ± 13.5	68	170.6 ± 19.7	44	177.3 ± 13.3	65	170.7 ± 19.6	

N: Number of samples

of the second upper and lower molars. A similar situation is observed when gender is the variable compared. In each stage, the average age of the male patients was higher than that of the female patients. These values make possible the development of formulas to predict the patient’s age based on the calcification stage of the second molar evaluated.

Tables 2 and 3 present the descriptive summaries of the participants’ ages, the calcification stage of Demirjian, the upper and lower second molars, and gender.

Discussion

Estimation of age at death of human remains constitutes an important contribution to the identification of missing people, and also provides valuable data in the paleodemographic analysis of historical skeletal remains.¹⁷ In this study, based on the results, it was possible to develop formulas to predict chronological age in a specific Brazilian population, contributing significantly to age estimation in forensic evaluations.

Schmeling et al¹⁸ noted the influence of ethnicity on the development of systems of age estimation and the difficulty when the subjects of forensic examination mostly be-

longed to populations for which no reference studies could be available for forensic purposes. Following this line of thought, Chen et al⁶ observed that the development standards used by Demirjian et al for French-Canadian children were not suitable for western Chinese children of different origins; therefore, it is necessary to establish new tables for these populations.

In the present study, the method proposed by Demirjian et al was chosen as the assessment parameter to be compared to the international literature. In Brazil, the method of Nolla¹⁹ was frequently adopted. Moreover, we considered the need to evaluate dental age in the Brazilian population. In fact, there has been significantly less research on genetic, environmental, and even nutritional characteristics of the Brazilian population compared to other populations.

External influences, such as some environmental factors, socio-economic status, nutrition, dietary habits, and lifestyle, despite having little impact on dental development, should be taken into consideration.² Phillips and van Wyk Kotze¹² suggested that the socio-economic environment of a child might play a role in the calcification of the permanent teeth and influence the radiographic images of

these teeth for aging purposes. This would explain why the pubertal growth spurt did not occur at the same time for all individuals. The sample used in the present study was not evaluated for external and internal modifying factors. However, the participants probably had a high socioeconomic status since they were the patients of a private clinic.

Some studies have suggested restricting the evaluation of dental development to the lower elements because maxillary teeth cannot be easily seen on panoramic radiographs and little data may be available for these teeth.^{8,11} Nevertheless, in the present study, the sample was selected based on the possibility of detailed visualization of the degrees of calcification of the second molars. Therefore, 10 unclear panoramic images of the upper molars were excluded.

It has been postulated that Demirjian's dataset is only applicable for individuals aged from 3 years to 16 years.⁹ Furthermore, the minimum age of patients in the present study was established based on Moyers,²⁰ who explained that the development of permanent second molars began at 3 years of age for both genders. Although a minimum age of 3 and maximum of 16 years was a sample selection criterion in the present study, there were no patients aged less than 4 years and 9 months. This could be explained by the need for a solid justification for taking panoramic radiographs before the mixed dentition period. Furthermore, younger children were usually unable to stay perfectly still for 15 to 20 seconds, which might lead to incorrect patient positioning and distortion of the panoramic radiograph.

In the present study, there was a correlation with chronological age. This result was similar to the findings of others.^{4,5,7,14,15,21}

According to the study by Bagherian and Sadeghi,⁵ dental development, including that of the permanent second molars, was earlier in females than in males. This fact was also observed in the present study, with females showing a precocity of 4.3 to 4.4 months for the upper teeth and 3.0 to 3.7 months for the lower ones. However, Nolla did not find significant differences in the degree of mineralization of teeth between the genders.¹⁹

Nolla¹⁹ noticed that neither the upper nor the lower teeth in either gender showed significant differences in the degree of mineralization on the right and left side. On the other hand, in the present study, the degree of calcification was significantly different between the lower and upper teeth, with the former growing earlier than the latter, which was consistent with the findings of the study by Holtgrave et al²² carried out in the United Kingdom.

Our results indicated that the dental development in fe-

males could be considered to occur earlier than in males; in addition, the lower second molar teeth develop much earlier than the upper ones, and there was no difference between the right and left second permanent molars. In conclusion, it was observed that ethnic variables were related to certain parameters of age and gender in a Brazilian population, contributing important information for forensic evaluations.

References

1. Bagherpour A, Imanimoghaddam M, Bagherpour MR, Einolghozati M. Dental age assessment among Iranian children aged 6-13 years using the Demirjian method. *Forensic Sci Int* 2010; 197 : 121.e1-4.
2. Koshy S, Tandon S. Dental age assessment: the applicability of Demirjian's method in south Indian children. *Forensic Sci Int* 1998; 94 : 73-85.
3. Demirjian A, Goldstein H, Tanner JM. A new system of dental age assessment. *Hum Biol* 1973; 45 : 211-27.
4. Chaillet N, Nyström M, Kataja M, Demirjian A. Dental maturity curves in Finnish children: Demirjian's method revisited and polynomial functions for age estimation. *J Forensic Sci* 2004; 49 : 1324-31.
5. Bagherian A, Sadeghi M. Assessment of dental maturity of children aged 3.5 to 13.5 years using the Demirjian method in an Iranian population. *J Oral Sci* 2011; 53 : 37-42.
6. Chen JW, Guo J, Zhou J, Liu RK, Chen TT, Zou SJ. Assessment of dental maturity of western Chinese children using Demirjian's method. *Forensic Sci Int* 2010; 197 : 119.e1-4.
7. Choi JH, Kim CY. A study of correlation between the development of the third molar and second molar as an aid in age determination. *J Korean Acad Oral Med* 1991; 16 : 121-36.
8. Nyström M, Aine L, Peck L, Haavikko K, Kataja M. Dental maturity in Finns and the problem of missing teeth. *Acta Odontol Scand* 2000; 58 : 49-56.
9. Jayaraman J, King NM, Roberts GJ, Wong HM. Dental age assessment: are Demirjian's standards appropriate for southern Chinese children? *J Forensic Odontostomatol* 2011; 29 : 22-8.
10. Karataş OH, Öztürk F, Dedeoğlu N, Çolak C, Altun O. Dental age assessment: the applicability of Demirjian method in southwestern of eastern Anatolia region Turkish children. *Cumhuriyet Dent J* 2012; 15 : 130-7.
11. Phillips VM, van Wyk Kotze TJ. Testing standard methods of dental age estimation by Moorrees, Fanning and Hunt and Demirjian, Goldstein and Tanner on three South African children samples. *J Forensic Odontostomatol* 2009; 27 : 29-44.
12. Phillips VM, van Wyk Kotze TJ. Dental age related tables for children of various ethnic groups in South Africa. *J Forensic Odontostomatol* 2009; 27 : 20-8.
13. Willems G, Van Olmen A, Spiessens B, Carels C. Dental age estimation in Belgian children: Demirjian's technique revisited. *J Forensic Sci* 2001; 46 : 893-5.
14. Shim WH, Kim KS, Shin KJ, Yoon CL, Kim CY, Choi JH. Second and third molar calcification in relation to chronological age in Koreans. *J Korean Acad Oral Med* 2004; 29 : 329-40.

15. Lee SS, Byun YS, Park MJ, Choi JH, Yoon CL, Shin KJ. The chronology of second and third molar development in Koreans and its application to forensic age estimation. *Int J Legal Med* 2010; 124 : 659-65.
16. Thevissen PW, Pittayapat P, Fieuws S, Willems G. Estimating age of majority on third molars developmental stages in young adults from Thailand using a modified scoring technique. *J Forensic Sci* 2009; 54 : 428-32.
17. Meisl A, Tangl S, Huber C, Maurer B, Watzek G. The chronology of third molar mineralization in the Austrian population - a contribution to forensic age estimation. *Forensic Sci Int* 2007; 169 : 161-7.
18. Schmeling A, Reisinger W, Geserick G, Olze A. Age estimation of unaccompanied minors. Part I. General considerations. *Forensic Sci Int* 2006; 159 Suppl 1 : S61-4.
19. Nolla CM. The development of permanent teeth. *J Dent Child* 1960; 4 : 254-66.
20. Moyers RE. *Handbook of orthodontics for the student and general practitioner*. 3rd ed. Chicago: Year Book Medical; 1973.
21. Lee SE, Lee SH, Lee JY, Park HK, Kim YK. Age estimation of Korean children based on dental maturity. *Forensic Sci Int* 2008; 178 : 125-31.
22. Holtgrave EA, Kretschmer R, Müller R. Acceleration in dental development: fact or fiction. *Eur J Orthod* 1997; 19 : 703-10.