

RIGHT AND LEFT FREDHOLM OPERATOR MATRICES

DRAGAN S. DJORDJEVIĆ AND MILICA Z. KOLUNDŽIJA

ABSTRACT. We consider right and left Fredholm operator matrices of the form $\begin{bmatrix} A & C \\ T & S \end{bmatrix}$, which are linear and bounded on the Banach space $Z = X \oplus Y$.

1. Introduction

Let Z be an infinite dimensional Banach space, such that $Z = X \oplus Y$ for some closed subspaces X and Y . This sum will be also denoted by $\begin{bmatrix} X \\ Y \end{bmatrix}$. If W is a finite dimensional subspace of X , then $\dim W$ denotes its dimension. If W is infinite dimensional, then we simply write $\dim W = \infty$. However, if U is a closed subspace of a Hilbert space, then $\dim_H(U)$ denotes the orthogonal dimension of U .

Let $\mathcal{L}(X, Y)$ denote the set of all linear bounded operators from X to Y . We abbreviate $\mathcal{L}(X) = \mathcal{L}(X, X)$. The set of all finite rank operators from X to Y is denoted by $\mathcal{F}(X, Y)$. For $A \in \mathcal{L}(X, Y)$ we use $\mathcal{R}(A)$ and $\mathcal{N}(A)$ to denote the range and the null-space of A , respectively.

If $Z = X \oplus Y$, then any $M \in \mathcal{L}(Z)$ can be decomposed as the following operator matrix

$$M = \begin{bmatrix} A & C \\ T & S \end{bmatrix} : \begin{bmatrix} X \\ Y \end{bmatrix} \rightarrow \begin{bmatrix} X \\ Y \end{bmatrix}$$

for some $A \in \mathcal{L}(X)$, $C \in \mathcal{L}(Y, X)$, $T \in \mathcal{L}(X, Y)$ and $S \in \mathcal{L}(Y)$. On the other hand, any choice of A, C, T, S (linear and bounded operators on the corresponding subspaces), produces a linear and bounded operator M on the space Z . Moreover, M is finite rank if and only if all A, C, T, S are finite rank operators.

If A and C are fixed, then we use the notation $M_{(T,S)}$ to show that M depends on T and S . For given A and C , we are interested to find T and S , such that $M_{(T,S)}$ is right or left Fredholm operator.

Received June 13, 2012.

2010 *Mathematics Subject Classification.* 47A53.

Key words and phrases. right (left) Fredholm, operator matrices.

The authors are supported by the Ministry of Education and Science, Serbia, grant no. 174007.

For this purpose we need to review some properties of right and left Fredholm operators [9]. An operator $A \in \mathcal{L}(X, Y)$ is right Fredholm, if $\text{def}(A) = \dim Y/\mathcal{R}(A) < \infty$, and $\mathcal{N}(A)$ is complemented in X . Notice that if A is right Fredholm, then it follows that $\mathcal{R}(A)$ has to be a closed and complemented subspace of Y . The set of all right Fredholm operators from X to Y is denoted by $\Phi_r(X, Y)$. It is well-known that $A \in \Phi_r(X, Y)$ if and only if there exist $B \in \mathcal{L}(Y, X)$ and $F \in \mathcal{F}(Y)$ such that $AB = I_Y + F$ holds.

An operator $A \in \mathcal{L}(X, Y)$ is left Fredholm, if $\text{nul}(A) = \dim \mathcal{N}(A) < \infty$, and $\mathcal{R}(A)$ is closed and complemented in Y . The set of all left Fredholm operators from X to Y is denoted by $\Phi_l(X, Y)$. It is well-known that $A \in \Phi_l(X, Y)$ if and only if there exist $B \in \mathcal{L}(Y, X)$ and $F \in \mathcal{F}(X)$ such that $BA = I_X + F$ holds.

If $A \in \Phi_r(X, Y)$ and $B \in \Phi_r(Y, Z)$, then $BA \in \Phi_r(X, Z)$. The similar result holds for the class Φ_l . The set of Fredholm operators is defined as $\Phi(X, Y) = \Phi_r(X, Y) \cap \Phi_l(X, Y)$.

We formulate the following well-known results.

Lemma 1.1. *Let X, Y, Z be Banach spaces and let $A \in \mathcal{L}(X, Y)$, $B \in \mathcal{L}(Y, Z)$. If $BA \in \Phi(X, Z)$, then the following holds: $A \in \Phi(X, Y)$ if and only if $B \in \Phi(Y, Z)$.*

Lemma 1.2. *Let X, Y be Banach spaces, and let $A \in \Phi_r(X, Y)$, $P \in \mathcal{F}(X, Y)$. Then $A + P \in \Phi_r(X, Y)$. The analogous result holds for classes Φ_l and Φ .*

Lemma 1.3. *Let M_1, M_2 and N be the vector subspaces of the vector space X . If $M_1 \subseteq M_2$, then $\dim M_1/(M_1 \cap N) \leq \dim M_2/(M_2 \cap N)$.*

Properties of right (left) Fredholm and related operators can be found in [6] and [9]. For the importance and applications of operator matrices we refer to [1], [2], [3], [4], [5], [7], [8] and [10]. Particularly, this paper is related to the research in [4] and [7], where the left and right invertibility of $M_{(T,S)}$ is considered.

2. Right Fredholm operators

We consider right Fredholm properties of $M_{(T,S)}$.

Theorem 2.1. *Let $A \in \mathcal{L}(X)$ and $C \in \mathcal{L}(Y, X)$ be given. The following statements are equivalent:*

- (a) $[A \ C] \in \Phi_r(X \oplus Y, X) \setminus \Phi(X \oplus Y, X)$, and there exists an operator $J \in \Phi_l(Y, \mathcal{N}([A \ C]) \setminus \Phi(Y, \mathcal{N}([A \ C])))$.
- (b) $M_{(T,S)} \in \Phi_r(X \oplus Y) \setminus \Phi(X \oplus Y)$ for some $T \in \mathcal{L}(X, Y)$ and $S \in \mathcal{L}(Y)$.

Proof. (a) \implies (b): Suppose that $[A \ C] \in \Phi_r(X \oplus Y, X) \setminus \Phi(X \oplus Y, X)$. It follows that $\mathcal{N}([A \ C])$ is infinite dimensional. By the assumption, there exists an operator $J \in \Phi_l(Y, \mathcal{N}([A \ C]) \setminus \Phi(Y, \mathcal{N}([A \ C])))$, so $\mathcal{N}(J)$ is finite

dimensional and $\mathcal{N}([A \ C])/R(J)$ is infinite dimensional. The operator J has the form

$$J = \begin{bmatrix} E \\ G \end{bmatrix} : Y \rightarrow \begin{bmatrix} X \\ Y \end{bmatrix}.$$

Since $\mathcal{R}(J)$ is closed and complemented in $\mathcal{N}([A \ C])$, and $\mathcal{N}([A \ C])$ is closed and complemented in $X \oplus Y$, we obtain that there exist closed subspaces V and W such that $\mathcal{N}([A \ C]) = R(J) \oplus V$ and $X \oplus Y = \mathcal{N}([A \ C]) \oplus W = R(J) \oplus V \oplus W$. Notice that V is infinite dimensional.

There exists a closed subspace Y_1 such that $Y = \mathcal{N}(J) \oplus Y_1$. Now, the reduction operator $J : Y_1 \rightarrow \mathcal{R}(J)$ is invertible, so let $K_1 : \mathcal{R}(J) \rightarrow Y_1$ denote its inverse. Define the operator $K \in \mathcal{L}(X \oplus Y, Y)$ in the following way:

$$Kx = \begin{cases} K_1x, & x \in \mathcal{R}(J), \\ 0, & x \in V \oplus W. \end{cases}$$

Then $K \in \mathcal{L}(X \oplus Y, Y)$ is a right Fredholm operator, such that $\mathcal{N}(K) = V \oplus W$. The operator K has the matrix form

$$K = [T \ S] : \begin{bmatrix} X \\ Y \end{bmatrix} \rightarrow Y.$$

We also have

$$(1) \quad KJ = [T \ S] \begin{bmatrix} E \\ G \end{bmatrix} = I_Y - P_1,$$

where P_1 is the projection from Y onto the finite dimensional subspace $\mathcal{N}(J)$, parallel to Y_1 .

From $\mathcal{R}(J) \subset \mathcal{N}([A \ C])$ we get that

$$(2) \quad [A \ C] \begin{bmatrix} E \\ G \end{bmatrix} = 0.$$

Since $[A \ C] \in \Phi_r(X \oplus Y, X)$, we have the following decompositions of spaces: $X \oplus Y = \mathcal{N}([A \ C]) \oplus W$ and $X = \mathcal{R}([A \ C]) \oplus U$, where U is finite dimensional. Since the reduction $[A \ C] : W \rightarrow \mathcal{R}([A \ C])$ is invertible, define $L_1 : \mathcal{R}([A \ C]) \rightarrow W$ to be its inverse. Then consider the operator $L \in \mathcal{L}(X, X \oplus Y)$, which is defined as follows:

$$Lx = \begin{cases} L_1x, & x \in \mathcal{R}([A \ C]) \\ 0, & x \in U. \end{cases}$$

The operator L has the matrix form

$$L = \begin{bmatrix} D \\ F \end{bmatrix} : X \rightarrow \begin{bmatrix} X \\ Y \end{bmatrix}.$$

Then $L \in \Phi_l(X, X \oplus Y)$, $\mathcal{R}(L) = W$, and

$$(3) \quad [A \ C]L = [A \ C] \begin{bmatrix} D \\ F \end{bmatrix} = I_X - P_2,$$

where P_2 is the projection from X onto the finite dimensional subspace U , parallel to $\mathcal{R}([A \ C])$. Since $\mathcal{N}([T \ S]) = V \oplus W$, we conclude that

$$(4) \quad [T \ S] \begin{bmatrix} D \\ F \end{bmatrix} = 0.$$

Finally, from (1), (2), (3) and (4), we get that for $M = \begin{bmatrix} A & C \\ T & S \end{bmatrix}$, $N = \begin{bmatrix} D & E \\ F & G \end{bmatrix}$ the following holds:

$$(5) \quad MN = \begin{bmatrix} A & C \\ T & S \end{bmatrix} \begin{bmatrix} D & E \\ F & G \end{bmatrix} = \begin{bmatrix} I_X & 0 \\ 0 & I_Y \end{bmatrix} + \begin{bmatrix} -P_2 & 0 \\ 0 & -P_1 \end{bmatrix}.$$

Since $\begin{bmatrix} -P_2 & 0 \\ 0 & -P_1 \end{bmatrix}$ is finite rank, we conclude that M is right Fredholm. Moreover, we notice that

$$\begin{aligned} \mathcal{N}(M) &= \mathcal{N}([A \ C]) \cap \mathcal{N}([T \ S]) = V, \\ \mathcal{R}(N) &= \mathcal{R}\left(\begin{bmatrix} D \\ F \end{bmatrix}\right) + \mathcal{R}\left(\begin{bmatrix} E \\ G \end{bmatrix}\right) = W \oplus \mathcal{R}(J), \\ X \oplus Y &= \mathcal{R}(J) \oplus V \oplus W. \end{aligned}$$

Since V is infinite dimensional, we obtain that both M and N are not Fredholm operators.

(b) \implies (a): Suppose that there exist some $T \in \mathcal{L}(X, Y)$ and $S \in \mathcal{L}(Y)$ such that $M_{(T, S)} \in \Phi_r(X \oplus Y) \setminus \Phi(X, Y)$. Then there exist operators $N \in \mathcal{L}(X \oplus Y)$ and $P \in \mathcal{F}(X \oplus Y)$ such that $MN = I + P$. The last equality holds in the matrix form as follows:

$$\begin{bmatrix} A & C \\ T & S \end{bmatrix} \begin{bmatrix} D & E \\ F & G \end{bmatrix} = \begin{bmatrix} I_X & 0 \\ 0 & I_Y \end{bmatrix} + \begin{bmatrix} P_{11} & P_{12} \\ P_{21} & P_{22} \end{bmatrix},$$

where all P_{ij} are finite rank operators. It also follows that $N = \begin{bmatrix} D & E \\ F & G \end{bmatrix} \in \Phi_l(X \oplus Y)$.

In particular, we obtain

$$[A \ C] \begin{bmatrix} D \\ F \end{bmatrix} = I_X + P_{11},$$

so $[A \ C]$ is right Fredholm. The operator $I_X + P_{11}$ is Fredholm. If we suppose that $[A \ C]$ is Fredholm, by Lemma 1.1 it follows that $\begin{bmatrix} D \\ F \end{bmatrix}$ is also Fredholm. Since

$$\mathcal{R}\left(\begin{bmatrix} D & E \\ F & G \end{bmatrix}\right) = \mathcal{R}\left(\begin{bmatrix} D \\ F \end{bmatrix}\right) + \mathcal{R}\left(\begin{bmatrix} E \\ G \end{bmatrix}\right) \supset \mathcal{R}\left(\begin{bmatrix} D \\ F \end{bmatrix}\right),$$

it follows that $\begin{bmatrix} D & E \\ F & G \end{bmatrix}$ belongs to $\Phi_r(X \oplus Y)$, so $\begin{bmatrix} D & E \\ F & G \end{bmatrix}$ is Fredholm. By Lemma 1.1 again, we obtain that $\begin{bmatrix} A & C \\ T & S \end{bmatrix}$ is Fredholm (since $I + P$ is Fredholm from Lemma 1.2). The last statement is not possible, so we obtain that $[A \ C] \in \Phi_r(X \oplus Y, X) \setminus \Phi(X \oplus Y, X)$.

Denote with $L = \begin{bmatrix} E \\ G \end{bmatrix} \in \mathcal{L}(Y, X \oplus Y)$. We have $[T \ S]L = I_Y + P_{22}$, so $L \in \Phi_l(Y, X \oplus Y) \setminus \Phi(Y, X \oplus Y)$. Otherwise, if L is Fredholm, then also $\begin{bmatrix} D & E \\ F & G \end{bmatrix}$ is Fredholm, so $\begin{bmatrix} A & C \\ T & S \end{bmatrix}$ is Fredholm.

Since we have the following decomposition of space $X \oplus Y = \mathcal{N}([A \ C]) \oplus W$, the operator L has the matrix form

$$L = \begin{bmatrix} J \\ K \end{bmatrix} : Y \rightarrow \begin{bmatrix} \mathcal{N}([A \ C]) \\ W \end{bmatrix}.$$

From the fact that

$$\mathcal{R}(P_{12}) = \mathcal{R}([A \ C]L) = \mathcal{R}\left([A \ C] \begin{bmatrix} J \\ K \end{bmatrix}\right) = [A \ C](\mathcal{R}(K))$$

is a finite space and the reduction $[A \ C] : W \rightarrow \mathcal{R}([A \ C])$ is a bijection, we obtain that $\mathcal{R}(K)$ is a finite dimensional subspace of W .

Since $L \in \Phi_l(Y, X \oplus Y) \setminus \Phi(Y, X \oplus Y)$, we have the following decompositions of spaces $Y = \mathcal{N}(L) \oplus U$ and $X \oplus Y = \mathcal{R}(L) \oplus U_1$, where $\dim \mathcal{N}(L) < \infty$ and $\dim U_1 = \infty$. The reduction operator $L : U \rightarrow \mathcal{R}(L)$ is invertible, so let $L_1 : \mathcal{R}(L) \rightarrow U$ be its inverse.

As it was shown, $\mathcal{R}(K)$ is a finite dimensional subspace, so $Y_1 = L_1(\mathcal{R}(K))$ have to be a finite dimensional subspace of U and there exists a closed subspace Y_2 such that $U = Y_1 \oplus Y_2$.

Now, the operator L has the following matrix form

$$L = \begin{bmatrix} J & 0 & 0 \\ 0 & K & 0 \end{bmatrix} : \begin{bmatrix} Y_2 \\ Y_1 \\ \mathcal{N}(L) \end{bmatrix} \rightarrow \begin{bmatrix} \mathcal{N}([A \ C]) \\ W \end{bmatrix},$$

where Y_1 is finite dimensional. We obtain that $\mathcal{N}(J) = Y_1 \oplus \mathcal{N}(L)$, so $\dim \mathcal{N}(J) < \infty$.

From the fact that $[T \ S]L = I_Y + P_{22}$ follows that

$$L_1(\mathcal{N}([T \ S]) \cap \mathcal{R}(L)) \subseteq \mathcal{N}(I_Y + P_{22}).$$

Since $I_Y + P_{22}$ is a Fredholm operator, we have that $L_1(\mathcal{N}([T \ S]) \cap \mathcal{R}(L))$ is finite dimensional, so $\mathcal{N}([T \ S]) \cap \mathcal{R}(L)$ is also a finite dimensional subspace.

Denote with $V = \mathcal{N}([A \ C]) \cap \mathcal{N}([T \ S]) \cap \mathcal{R}(J)$. Further,

$$V \subseteq \mathcal{N}([T \ S]) \cap \mathcal{R}(J) \subseteq \mathcal{N}([T \ S]) \cap \mathcal{R}(L),$$

so it follows that $\dim V < \infty$. Then, there exists a closed subspace V_1 such that $\mathcal{N}(M_{(T,S)}) = \mathcal{N}([A \ C]) \cap \mathcal{N}([T \ S]) = V \oplus V_1$. Since $\mathcal{N}(M_{(T,S)})$ is infinite dimensional, then V_1 is also an infinite dimensional subspace.

Now, applying Lemma 1.3 on the spaces $\mathcal{N}([A \ C]) \cap \mathcal{N}([T \ S])$, $\mathcal{N}([A \ C])$ and $\mathcal{R}(J)$, we obtain

$$\dim V_1 = \dim(\mathcal{N}([A \ C]) \cap \mathcal{N}([T \ S]))/V \leq \dim \mathcal{N}([A \ C])/\mathcal{R}(J).$$

We conclude that $\dim \mathcal{N}([A \ C])/\mathcal{R}(J) = \infty$.

Lastly, we proved for the operator $J : Y \rightarrow \mathcal{N}([A \ C])$ that $\dim \mathcal{N}(J) < \infty$ and $\dim \mathcal{N}([A \ C])/\mathcal{R}(J) = \infty$.

So, there exists the operator $J \in \Phi_l(Y, \mathcal{N}([A \ C]) \setminus \Phi(Y, \mathcal{N}([A \ C])))$. \square

3. Left Fredholm operators

Now we investigate the left Fredholm properties of $M_{(T,S)}$. We consider two separate cases according to the dimension of Y .

Theorem 3.1. *Let X be infinite dimensional, and let Y be finite dimensional. For given $A \in \mathcal{L}(X)$ and $C \in \mathcal{L}(Y, X)$, the following statements are equivalent:*

- (a) $M_{(T,S)} \in \Phi_l(X \oplus Y) \setminus \Phi(X \oplus Y)$ for every $T \in \mathcal{L}(X, Y)$ and every operator $S \in \mathcal{L}(Y)$;
- (b) $A \in \Phi_l(X) \setminus \Phi(X)$.

Proof. Before the proof of the equivalence, note that

$$\mathcal{N}\left(\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}\right) = \mathcal{N}(A) \oplus Y, \quad \mathcal{R}\left(\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}\right) = \mathcal{R}(A) \oplus \{0\}.$$

Since Y is finite dimensional, we have that $A \in \Phi_l(X) \setminus \Phi(X)$ if and only if $\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix} \in \Phi_l(X \oplus Y) \setminus \Phi(X \oplus Y)$.

(a) \implies (b): Suppose that $M_{(T,S)}$ is left Fredholm but not Fredholm, for every $T \in \mathcal{L}(X, Y)$ and every $S \in \mathcal{L}(Y)$. We have that $\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} A & C \\ T & S \end{bmatrix} + \begin{bmatrix} 0 & -C \\ -T & -S \end{bmatrix}$ where $\begin{bmatrix} 0 & -C \\ -T & -S \end{bmatrix}$ is a finite rank operator. Applying Lemma 1.2, we obtain that $\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}$ is a left Fredholm operator.

Suppose that $\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}$ is Fredholm. Applying Lemma 1.2 to $\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}$ we conclude that $M_{(T,S)}$ has to be Fredholm, which does not hold. Hence, $\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}$ is left Fredholm but not Fredholm, so we have that $A \in \Phi_l(X) \setminus \Phi(X)$.

(b) \implies (a): Suppose that A is left Fredholm but not Fredholm, so the operator $\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}$ is also left Fredholm but not Fredholm.

Let $T \in \mathcal{L}(X, Y)$ and $S \in \mathcal{L}(Y)$ be arbitrary operators. Then the operator $M_{(T,S)}$ is a finite-rank perturbation of $\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}$. Indeed, $\begin{bmatrix} A & C \\ T & S \end{bmatrix} = \begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & C \\ T & S \end{bmatrix}$, where $\begin{bmatrix} 0 & C \\ T & S \end{bmatrix}$ is a finite rank operator because Y is a finite dimensional space. Applying Lemma 1.2 to $\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}$ we get that $M_{(T,S)}$ is a left Fredholm operator. If we suppose that $M_{(T,S)}$ is Fredholm, from Lemma 1.2, we conclude that $\begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}$ have to be Fredholm, which does not hold. We obtain that $M_{(T,S)}$ is left Fredholm but not Fredholm operator. \square

Theorem 3.2. *Let X and Y be infinite dimensional, such that Y is isomorphic to $Z = X \oplus Y$. Let $A \in \mathcal{L}(X)$ and $C \in \mathcal{L}(Y, X)$ be arbitrary. Then $M_{(T,S)} \in \Phi_l(X \oplus Y) \setminus \Phi(X \oplus Y)$ for some $T \in \mathcal{L}(X, Y)$ and $S \in \mathcal{L}(Y)$.*

Proof. Since Y is isomorphic with Z , then $Y = Y_1 \oplus Y_2$, where X is isomorphic to Y_1 , and Y is isomorphic to Y_2 . Let $T \in \mathcal{L}(X, Y_1)$ and $S \in \mathcal{L}(Y, Y_2)$ be those isomorphisms. Then $T \in \mathcal{L}(X, Y)$ is left invertible with a left inverse $K \in \mathcal{L}(Y, X)$ and $\mathcal{N}(K) = Y_2$. Also, $S \in \mathcal{L}(Y, Y_2)$ is left invertible with a left inverse L and $\mathcal{N}(L) = Y_1$. Then

$$\begin{bmatrix} 0 & K \\ 0 & L \end{bmatrix} \begin{bmatrix} A & C \\ T & S \end{bmatrix} = \begin{bmatrix} I_X & 0 \\ 0 & I_Y \end{bmatrix},$$

so $M_{(T,S)}$ is left invertible. It follows that $M_{(T,S)}$ is left Fredholm for chosen operators T and S . Suppose that $M_{(T,S)}$ is Fredholm. Since $\begin{bmatrix} I_X & 0 \\ 0 & I_Y \end{bmatrix}$ is Fredholm, from Lemma 1.1 it follows that N is also Fredholm. However, we notice $\mathcal{N}(N) = X$, which is infinite dimensional. Hence, N is not Fredholm. Then $M_{(T,S)}$ is not Fredholm also, i.e., $M_{(T,S)} \in \Phi_l(X \oplus Y) \setminus \Phi(X \oplus Y)$. \square

We formulate a corollary for Hilbert space operators.

Corollary 3.1. *Let X and Y be infinite dimensional and mutually orthogonal subspaces of a Hilbert space $Z = X \oplus Y$. Suppose that $\dim_H Y = \dim_H Z$. Let $A \in \mathcal{L}(X)$ and $C \in \mathcal{L}(Y, X)$ be arbitrary. Then $M_{(T,S)} \in \Phi_l(X \oplus Y) \setminus \Phi(X \oplus Y)$ for some $T \in \mathcal{L}(X, Y)$ and $S \in \mathcal{L}(Y)$.*

References

- [1] D. S. Djordjević, *Perturbations of spectra of operator matrices*, J. Operator Theory **48** (2002), no. 3, 467–486.
- [2] H. Du and J. Pan, *Perturbation of spectrums of 2×2 operator matrices*, Proc. Amer. Math. Soc. **121** (1994), no. 3, 761–766.
- [3] B. P. Duggal, *Upper triangular operator matrices, SVEP and Browder, Weyl theorems*, Integral Equations Operator Theory **63** (2009), no. 1, 17–28.
- [4] G. Hai and A. Chen, *Perturbations of the right and left spectra for operator matrices*, J. Operator Theory **67** (2012), no. 1, 207–214.
- [5] J. K. Han, H. Y. Lee, and W. Y. Lee, *Invertible completions of 2×2 upper triangular operator matrices*, Proc. Amer. Math. Soc. **128** (2000), no. 1, 119–123.
- [6] R. Harte, *Invertibility and Singularity for Bounded Linear Operators*, Marcel Dekker, New York, 1988.
- [7] M. Kolundžija, *Right invertibility of operator matrices*, Funct. Anal. Approx. Comput. **2** (2010), no. 1, 1–5.
- [8] W. Y. Lee, *Weyl spectra of operator matrices*, Proc. Amer. Math. Soc. **129** (2001), no. 1, 131–138.
- [9] V. Müller, *Spectral Theory of Linear Operators and Spectral Systems in Banach Algebras*, Birkhäuser Verlag, Basel-Boston-Berlin, 2007.
- [10] C. Tretter, *Spectral Theory of Block Operator Matrices and Applications*, Imperial College Press, London, 2008.

DRAGAN S. DJORDJEVIĆ
 FACULTY OF SCIENCES AND MATHEMATICS
 UNIVERSITY OF NIŠ
 VIŠEGRADSKA 33, P.O. BOX 224, 18000 NIŠ, SERBIA
 E-mail address: dragan@pmf.ni.ac.rs

MILICA Z. KOLUNDŽIJA
 FACULTY OF SCIENCES AND MATHEMATICS
 UNIVERSITY OF NIŠ
 VIŠEGRADSKA 33, P.O. BOX 224, 18000 NIŠ, SERBIA
 E-mail address: mkolundzija@pmf.ni.ac.rs