



Growth, Feed Utilization and Blood Metabolic Responses to Different Amylose-amylopectin Ratio Fed Diets in Tilapia (*Oreochromis niloticus*)

Meng-Yao Chen, Ji-Dan Ye*, Wei Yang, and Kun Wang

Xiamen Key Laboratory for Feed Quality Testing and Safety Evaluation, Jimei University, Xiamen 361021, China

ABSTRACT: A feeding trial was conducted in tilapia to determine the growth performance, nutrient digestibility, digestive enzymes, and postprandial blood metabolites in response to different dietary amylose-amylopectin ratios. Five isonitrogenous and isolipidic diets containing an equal starch level with different amylose-amylopectin ratios of 0.11 (diet 1), 0.24 (diet 2), 0.47 (diet 3), 0.76 (diet 4) and 0.98 (diet 5) were formulated using high-amylose corn starch (as the amylose source) and waxy rice (as the amylopectin source). Each diet was hand-fed to six tanks of 15 fish each, three times a day over a 6-wk period. After the growth trial, a postprandial blood metabolic test was carried out. Fish fed diet 2 exhibited the highest percent weight gain and feed efficiency and protein efficiency ratio, whereas fish fed with diet 5 showed the lowest growth and feed utilization among treatments. The digestibility for starch in fish fed diet 1 and 2 was higher than those in fish fed with other diets ($p < 0.05$). The highest activities for protease, lipase and amylase were found in fish fed the diet 2, diet 1, and diet 1 respectively among dietary treatments, while the lowest values for these indexes were observed in fish fed the diet 3, diet 5 and diet 4, respectively. The liver glycogen concentrations in fish fed diets 4 and 5 were found higher than in fish fed other diets ($p < 0.05$). The feeding rate, hepatosomatic index, condition factor, and plasma parameters (glucose, triglyceride, total cholesterol, high-density lipoprotein cholesterol, and low-density lipoprotein cholesterol) did not differ across treatments. In terms of postprandial blood responses, peak blood glucose and triglycerides were lower after 3 or 6 h in the fish fed with diets 3-5 than in the fish fed diet 1, but delayed peak blood total amino acid time was observed in fish fed with the diets 1 or 2. The lowest peak values for each of the three blood metabolites were observed in fish fed diet 5. The results indicate that high-dietary amylose-amylopectin ratio could compromise growth, but help in reducing the blood glucose stress on fish caused by postprandial starch load. (**Key Words:** Performance, Amylose, Amylopectin, Postprandial Response, Tilapia)

INTRODUCTION

Carbohydrate and particular cereal starch are the main sources for energy production of most terrestrial warm-blooded animals including human beings, but its impact is less in fish (Wilson, 1994). Although fish have the same carbohydrate metabolic pathways as mammals, they have a lower carbohydrate tolerance than mammals, as evidenced by a prolonged postprandial hyperglycemia after feeding carbohydrate-rich diets (Wilson, 1994; Moon, 2001). As a result, the use of carbohydrate for energy is limited and a high dietary inclusion level of carbohydrates could lead to growth retardation in fish (Wilson, 1994). Fish depend more than most other animals on amino acids as precursors to glucose for tissue protein and glucose synthesis as well as

energy (Mommensen and Plisetskaya, 1991; Moon, 2001). Nevertheless, from an economic point of view, it is desirable to reduce the protein and lipid use for energy consumption by increasing the carbohydrate use in the aquafeeds where carbohydrates are more readily available and less expensive than other macro-nutrients. The nutritional value of starches varies widely depending on its sources (Granfeldt et al., 1994; Gutierrez Del Alamo et al., 2009; Glencross et al., 2012), processing (Mahadevamma and Tharanathan, 2007), hardness (Denardin et al., 2012), non-starch polysaccharides (Glencross et al., 2012) and fish species (Wilson, 1994) as well as its complexity (Jun et al., 2010). Furthermore, previous studies have shown that the metabolic responses induced by dietary starches differ (Hemre et al., 2002; Stone, 2003; Enes et al., 2009). Starch is composed of two types of glucose polymers, amylose, a linear chain of glucose units joined by α -1,4-glucosidic linkages, and amylopectin, a highly branched amylose in

* Corresponding Author: Ji-Dan Ye. Tel: +86-592-6181420, Fax: +86-592-6181476, E-mail: yjdwk@sina.com
Submitted Jan. 8, 2013; Accepted Apr. 6, 2013; Revised May 9, 2013

which the branches are tied by α -1,6 bonds (Zobel, 1988). Amylopectin-rich starches are easily digested, which may induce a faster and higher glycemic and insulinemic responses compared to those high in amylose (Behall et al., 1989; Weststrate and van Amelsvoort, 1993; Granfeldt et al., 1995; Behall et al., 2002; Rawles and Lochmann, 2003). Although these metabolic responses were previously observed both in mammals and fish, the mechanisms behind these responses are not well understood.

In terms of growth effects in relation to starch type, most of the research has focused on terrestrial animals, but contradictory results have occurred in different studies. The positive effects on growth in response to dietary amylose were observed in pigs (Cohen and Tanksley, 1973), poultry (Pirgozliev et al., 2000; Gutierrez Del Alamo et al., 2009) and goats (Kyuma and Ishida, 1989), but the negative effects were observed in rats (Aziz et al., 2009; Denardin et al., 2012), pigs (Perez and Aumaitre, 1979; Camp et al., 2003), and poultry (Ertl and Dale, 1997; Bergh et al., 1999; Pirgozliev et al., 2002; Collins et al., 2003; Pirgozliev et al., 2010). No effect of starch type on growth was also observed in rats (Mongeau et al., 1995; Zhou and Kaplan, 1997; Kabir et al., 1998) and pigs (Shelton et al., 2004). In fish, Rawles and Lochmann (2003) first investigated the starch utilization related to starch type and observed an enhanced growth of sunshine bass (*Morone chrysops* ♀ × *M. saxatilis* ♂) when fed with a diet at the higher ratio of amylose-amylopectin starch. To the best of our knowledge, the impact of dietary amylose-amylopectin ratio in the starch fraction of the diet on other fish species has not yet been determined. The omnivorous fish such as tilapia (*Oreochromis niloticus*) have greater ability to digest starch than carnivorous fish (Krogdahl et al., 2005). We hypothesized that the tilapia fed diets with increasing amylose-amylopectin ratio would exhibit better growth. Therefore, we investigated the effects of dietary amylose-amylopectin ratio on the growth, feed utilization, digestibility, digestive enzyme activity, and blood intermediary metabolites in the tilapia.

MATERIALS AND METHODS

Diets

High-amylose corn starch (Hi-Maize 260, National Starch and Chemical Co., Ltd., China, 66.9% amylose) and waxy rice (the local glutinous rice, 81.0% amylopectin and 0.1% amylose) were used as the main amylose and amylopectin sources, respectively. Steam dried fish meal was obtained from Shine Biotech Co., Ltd., China. Fish oil, soybean oil, and other feed ingredients were obtained from Baisuihang Feed Co. Ltd., China. Based on the proximate composition of feed ingredients, five isonitrogenous (32% crude protein) and isolipidic (6.5% crude fat) diets

containing an equal starch level (35% starch) were formulated. Diets differing in ratios of amylose-amylopectin (diet 1, 0.11; diet 2, 0.24; diet 3, 0.47; diet 4, 0.76; and diet 5, 0.98) were achieved by adjusting the proportion of corn starch to waxy rice, i.e. by incremental levels of amylose from 2.58% to 14.12% at the expense of amylopectin. The ingredients and proximate composition of the experimental diets are shown in Table 1. The coarse dry ingredients were ground using a hammer mill and then passed through a 60 mesh sieve, weighed and blended together into a homogeneous mixture. The liquid ingredients (mixed oil and water) were then added, and the mixture was made into a mash. The dough was pelleted through a 2.5-mm die using multifunctional spiral extrusion machinery (CD4XITS, South China University of Technology, China). The pellets were dried in a forced-air oven at 50°C for 24 h and then sealed in plastic bags and kept at -20°C until further use for feeding.

Feeding procedures

The tilapia juveniles obtained from a local commercial farm in Xiamen were transported to the aquaculture laboratory of Jimei University. The fish were stocked into a closed recirculating system consisting of two circular fiber glass tanks (0.85 m height × 1.22 m upper diameter, 1.04 m lower diameter) with a polygeyser bead filter (Aquaculture Systems Technologies, LLC., USA) and were initially maintained on a commercial tilapia feed for 15 d. A total of four hundred and fifty animals, initially weighing an average of 9.81 ± 0.42 g (means ± SD) were randomly divided into five experimental groups, each with six replicate 100-L tanks at a density of 15 fish per tank. During a feeding period of 42 d, the fish in each group were hand-fed on one of the diets to satiation at 08:00, 13:00 and 18:00 h per day under a natural photoperiod. After 7 d of feeding, feces were collected from wk 2, until the end of the feeding trial. Excess feed was collected one hour after each feeding and then feces were collected by siphoning. To minimize the nutrient leaching in feces, only fresh and intact feces were collected. Feces collected from a single tank were pooled each day and dried at 70°C. Dissolved oxygen was measured daily and nitrite-N was monitored twice a week with a multi-parameter photometer (HI83200, Hanna Instruments, USA). During the feeding trial, the water temperature, dissolved oxygen and nitrite-N ranged from 27.5 to 32°C, 4.9 to 5.6 mg/L and 0 to 0.2 mg/L, respectively.

Sampling procedures

At the termination of the feeding trial, fish in each tank were bulk weighed and counted after 24 h starvation. Three fish were randomly taken from each of four replicate tanks per treatment and killed with an overdose of MS 222

Table 1. Ingredients and composition of the experimental diets (as-fed basis)

Ingredients (%)	Diets at different amylose-amylopectin ratios				
	Diet 1 (0.11)	Diet 2 (0.24)	Diet 3 (0.47)	Diet 4 (0.76)	Diet 5 (0.98)
Fish meal	45.77	46.28	46.9	47.34	47.67
Fish and soybean oil (1:1)	2.43	2.39	2.34	2.31	2.28
High-amylose corn starch ¹	3.85	8.47	14.12	18.15	21.17
Waxy rice	31.15	26.53	20.88	16.85	13.83
Microcrystalline cellulose ²	14.70	14.23	13.66	13.25	12.95
Choline chloride	0.1	0.1	0.1	0.1	0.1
Vitamin premix ³	0.4	0.4	0.4	0.4	0.4
Mineral premix ⁴	0.5	0.5	0.5	0.5	0.5
Sodium alginate	0.1	0.1	0.1	0.1	0.1
Ca(H ₂ PO ₄) ₂	1.0	1.0	1.0	1.0	1.0
Analyzed composition					
Dry matter (%)	91.59	91.21	91.93	91.18	91.30
Crude protein (%)	32.77	32.54	32.80	32.58	32.71
Crude fat (%)	6.85	6.41	6.83	6.56	6.72
Ash (%)	7.30	7.49	7.59	7.72	7.66
Amylose (%)	2.58	5.56	9.17	12.17	14.12
Amylopectin (%)	25.81	23.18	19.51	16.02	14.41
Amylose-amylopectin	0.11	0.24	0.47	0.76	0.98

¹ High-amylose corn starch (Hi-Maize 260) obtained from National Starch and Chemical Co., Ltd., Shanghai, China.

² Microcrystalline cellulose obtained from Baisuihang Feed Co. Ltd., Xiamen, China.

³ Vitamin premix (per kg diet): vitamin C (as Stay-C 35%) 400 mg; vitamin A (as vitamin A palmitate) 15,000 IU; vitamin D₃ 7,500 IU; vitamin E (as D, L- α -tocopheryl acetate) 150 mg; vitamin K₃ (as menadione sodium bisulphite) 7.5 mg; vitamin B₁ (as thiamin hydrochloride) 75 mg; riboflavin 60 mg; vitamin B₆ (as pyridoxine hydrochloride) 100 mg; D-calcium pantothenate 120 mg; nicotinic acid 150 mg; biotin 1.0 mg; inositol 450 mg.

⁴ Mineral premix (per kg diet): Fe (from Ferrous sulfate) 80 mg; Zn (from Zinc sulfate) 40 mg; Cu (from Copper sulfate) 5 mg; Mn (from Manganese sulfate) 20 mg; Se (from Sodium selenite) 0.2 mg; I (from Potassium iodine) 30 μ g; Co (from Cobalt chloride) 50 μ g; Mg (from Magnesium sulphate) 300 mg; P (from Sodium dihydrogen phosphate) 600 mg; K (from Potassium chloride) 740 mg.

(tricaine methanesulfonate) solution, pooled, weighed, minced and then dried at 70°C for whole-body composition analysis. Additionally, sixteen fish from four replicate tanks in each treatment were randomly sampled, anesthetized (MS 222, 100 mg/L) and weighed individually. Blood was collected via venipuncture using a heparinized syringe and immediately centrifuged at 3,000 \times g for 10 min at 4°C. Plasma samples from each tank were pooled and stored at -80°C for subsequent analysis of blood biochemical parameters. After blood collection, the liver and intestine were immediately removed and weighed. Liver and intestine samples were frozen in liquid nitrogen and stored at -80°C before use. The remaining fish in each treatment were used for the subsequent postprandial blood response test.

Postprandial blood metabolic response test

After sample collection, the remaining fish in each treatment were readjusted and distributed into the same tanks as used in the growth trial with three tanks of 22 fish per tank. Fish were continued to be maintained on their respective diets for 7 d. The next morning, fish received a single meal to apparent satiety within 30 min. Blood samples were collected from three fish in each tank with a

total of nine fish for each treatment at each sampling time, according to the above described method and blood were collected at 0, 1, 3, 6, 9, 12 and 24 h after a meal. Plasma aliquots were stored at -80°C for subsequent analysis of blood biochemical parameters.

Plasma components and liver glycogen analysis

Plasma glucose, triglyceride (TG), total cholesterol (T-CHO), high-density lipoprotein cholesterol (HDL-C) and low-density lipoprotein cholesterol (LDL-C) were measured by using kits (BioSino Bio-technology and Science, Inc., China). All the blood parameters were determined by using an automatic biochemical analyzer (Hitachi 7020, Japan) as described by Ye et al. (2011). Total plasma amino acid and liver glycogen were analyzed using the L-amino acid assay kit and glycogen assay kit, respectively (Nanjing Jiancheng Bioengineering Institute, China) according to the manufacturer's protocols by using UV-2800S spectrophotometer (UNICO Instruments Co., Ltd., China).

Digestive enzyme analysis

The pooled intestines of each sample were homogenized in chilled saline at a ratio of 1:9 (w/v) for 10 min. The supernatant was collected after centrifugation at 6,261 \times g for

15 min at 4°C. Aliquots of the supernatant, designated as the crude extract, were used to estimate the activities of amylase and protease, according to the procedure described by Ye et al. (2011), and the lipase activity was assayed with the lipase activity assay kit (NJBI, China) using a UV-2800S spectrophotometer.

Proximate composition analysis

Crude protein, crude fat, moisture and ash contents in feed ingredients, diets, feces and whole-body fish samples were determined according to the methods of the Association of Official Chemists (AOAC, 1995). Crude protein content was determined by the Kjeldahl method ($N \times 6.25$) using KjeltacTM 8400 Auto Sample Systems (Foss Tecator AB, Sweden). Crude fat content was determined by the Soxhlet method by using Soxtec Avanti 2050 (Foss Tecator AB). Moisture content was determined by the drying method, using an oven at 105°C. Ash content was determined by the combustion method, using a muffle furnace at 550°C. Acid-insoluble ash content was determined by using the same muffle furnace at 550°C, after acid digestion. Acid-insoluble ash was used as an endogenous indicator, instead of chromium oxide as an exogenous marker, to determine the nutrient digestibility because the dietary chromium oxide as a glucose tolerance factor is believed to affect starch utilization (Shiau and Chen, 1993). The total starch content in corn starch, waxy rice, diets and feces was calorimetrically determined by using the total starch assay kit (Megazyme International Ireland Ltd., Ireland). Using the amylose assay kit (Megazyme International Ireland Ltd., Ireland), the amylose content in high-amylose corn starch, sticky rice and diets was analyzed.

Calculations

Percent weight gain, feed conversion ratio, protein efficiency ratio, feeding rate, survival, hepatosomatic index and condition factor were calculated as follows:

$$\text{Percent weight gain (WG, \%)} = 100 \times (W_f - W_i) / W_i$$

$$\text{Feed efficiency (FE, \%)} = 100 \times (W_f - W_i) / W_c$$

$$\text{Protein efficiency ratio (PER)} = (W_f - W_i) / (W_c \times P)$$

$$\text{Feeding rate (FR, \%)} = 100 \times W_f / [(W_i + W_f) / 2] / t$$

$$\text{Hepatosomatic index (HSI, \%)} = 100 \times W_l / W_b$$

$$\text{Condition factor (CF)} = 100 \times W_b / L^3$$

where W_i (g) and W_f (g) are the initial body weight

(IBW) and final body weight (FBW) respectively; W_c (g) is the average feed intake per fish (on an as-fed basis); t is the time duration of the experiment (days); P is the protein content of the diet; W_l (g) is the liver weight of the sampled fish; W_b (g) is the final body weight of the sampled fish and L (cm) is the final body length of the sampled fish.

The apparent digestibility coefficients (ADCs) of the experimental diets for dry matter, crude protein, crude fat, ash and starch were calculated according to the following equation (Spyridakis et al., 1989):

$$\text{ADC (\%)} = 100 - 100 \times (N_f) / (N_d) \times (M_d) / (M_f)$$

where N_f is the percentage of a specific nutrient in feces, N_d is the percentage of the same nutrient in the diet, M_d is the percentage of acid-insoluble ash in the diet and M_f is the percentage of acid-insoluble ash in feces.

Statistical analysis

Data are presented as mean \pm SD. The normality and variance's homogeneity of all data were systematically checked before applying One-way ANOVA, and the mean values were compared by using the Student-Neuman-Keuls multiple comparison tests, at the 5% level of significance. The results of the measured variables were examined by regression. A repeated measures ANOVA was carried out to examine mean differences in postprandial concentrations of blood glucose, TG and total amino acids among treatments at time intervals. All statistical tests were calculated using SPSS13.0 software (SPSS Inc., Chicago, USA).

RESULTS

Growth performance

The fish in each treatment exhibited proper growth and survived across the feeding period. Fish fed diet 2 exhibited significantly ($p < 0.05$) higher WG (%), FE (%) and PER compared with fish fed diet 5 (Table 2). A strong negative correlation was observed between WG%, FE% and PER and dietary amylose-amylopectin ratio ($R = -0.8907$, -0.9347 and -0.9031 , respectively) and the values for WG (%), FE (%) and PER in fish fed diet 2 were significantly higher than those in fish fed diet 5 ($p < 0.05$). Although HSI (%) did not differ across treatments, the parameter was linearly ($R = -0.9728$) correlated with dietary amylose-amylopectin ratio. No influences of dietary amylose-amylopectin ratio on FR% and CF could be detected ($p > 0.05$).

Nutrient digestibility

As shown in Table 3, the digestibility values for dry matter, crude protein and lipid did not differ among dietary treatments ($p > 0.05$). However, there was a negatively linear

Table 2. Effects of dietary amylose-amylopectin ratios on the growth performance of tilapia over a feeding period of 6 wks

Diets ¹	Growth parameters						
	FAW (g) ²	WG (%) ²	FE (%) ²	PER ²	FR (%)	HSI (%) ³	CF ³
Diet 1 (0.11)	69.48 ^{ab}	606.7 ^{ab}	79.4 ^{ab}	2.42 ^{ab}	5.28	2.64	3.98
Diet 2 (0.24)	75.38 ^a	659.7 ^a	82.6 ^a	2.55 ^a	5.13	2.49	4.03
Diet 3 (0.47)	62.56 ^b	530.0 ^b	76.3 ^{ab}	2.33 ^{ab}	5.26	2.39	3.96
Diet 4 (0.76)	58.69 ^b	490.9 ^b	73.5 ^{ab}	2.25 ^{ab}	5.59	2.20	3.83
Diet 5 (0.98)	57.63 ^b	479.7 ^b	69.4 ^b	2.15 ^b	5.57	2.18	3.97
Pooled SE	2.045	20.541	1.399	0.043	0.083	0.078	0.038
p-value	0.009	0.006	0.015	0.015	0.334	0.028	0.592

WG (Y)-amylose-amylopectin ratio (X) correlation: $Y = 651.62 - 191.84X$; $R = -0.8907$; $p = 0.0427$.
 FE (Y)-amylose-amylopectin ratio (X) correlation: $Y = 83.05 - 13.29X$; $R = -0.9347$; $p = 0.0199$.
 PER (Y)-amylose-amylopectin ratio (X) correlation: $Y = 2.54 - 0.386X$; $R = -0.9031$; $p = 0.0357$.
 HSI (Y)-amylose-amylopectin ratio (X) correlation: $Y = 2.65 - 0.527X$; $R = -0.9728$; $p = 0.0054$.

Values with different superscripts in each column indicate significant difference ($p < 0.05$).

FAW = Final average weight; WG % = Percent weight gain; FE = Feed efficiency; PER = Protein efficiency ratio; FR = Feeding rate; HIS = Hepatosomatic index; CF = Condition factor.

¹ Values in parentheses are amylose-amylopectin ratios in diets. ² Data are means of four replicate tanks per dietary treatment.

³ Data are means of 24 fish per dietary treatment.

correlation between dry matter digestibility and dietary amylose-amylopectin ratio ($R = -0.8948$; $p = 0.0403$). The starch digestibility showed a decreasing trend with the increase in dietary amylose-amylopectin ratio ($R = -0.9893$, $p = 0.0013$), and the values of fish fed diets 1-2 were significantly ($p < 0.05$) higher than those of fish fed with other diets. When considering WG%, FE% and PER in relation to nutrient digestibility, the FE% and PER were linearly correlated with the values for dry matter ($R = 0.973$, $p = 0.005$; $R = 0.928$, $p = 0.023$, respectively) and for starch ($R = 0.956$, $p = 0.011$; $R = 0.903$, $p = 0.036$, respectively). The WG% was also correlated linearly with apparent digestibility of starch ($R = 0.911$, $p = 0.031$), but protein

and lipid digestibility did not show this relationship (Table 3).

Digestive enzyme activity

Highest protease activity was observed in fish fed diet 2, and was significantly ($p < 0.05$) higher than those in fish fed the other diets (Table 4). A decreasing trend of lipase ($R = -0.9080$, $p = 0.0330$) and amylase ($R = -0.8617$, $p = 0.0604$) activities was observed as dietary amylose-amylopectin ratio increased and the values of amylase and lipase in fish fed with diet 1 were significantly ($p < 0.05$) higher than those of fish fed other diets. The amylase activity was linearly correlated ($R = 0.910$; $p = 0.032$) with starch digestibility in

Table 3. Effects of dietary amylose-amylopectin ratios on nutrient digestibility for tilapia over a feeding period of 6 wks.

	Apparent digestibility coefficients (%) ²			
	Dry matter	Crude protein	Crude fat	Starch
Diet 1 (0.11)	82.29	90.85	94.44	72.53 ^a
Diet 2 (0.24)	82.95	91.53	93.06	70.76 ^a
Diet 3 (0.47)	82.19	91.14	93.96	66.47 ^b
Diet 4 (0.76)	81.59	90.98	94.89	64.25 ^b
Diet 5 (0.98)	80.49	90.45	93.97	61.59 ^b
Pooled SE	0.755	0.607	0.618	1.191
p-value	0.912	0.992	0.940	0.001

Dry matter (Y)-amylose-amylopectin ratio (X) correlation: $Y = 83.08 - 2.30X$; $R = 0.8948$; $p = 0.0403$.

Starch (Y)-amylose-amylopectin ratio (X) correlation: $Y = 73.48 - 12.43X$; $R = -0.9893$; $p = 0.0013$.

Dry matter (Y)-feed efficiency (X) correlation: $Y = 0.6849 + 0.18X$; $R = 0.973$; $p = 0.005$.

Dry matter (Y)-protein efficiency ratio (X) correlation: $Y = 4.59 + 0.82X$; $R = 0.928$; $p = 0.023$.

Starch (Y)-feed efficiency (X) correlation: $Y = 68.46 + 5.75X$; $R = 0.956$; $p = 0.011$.

Starch (Y)-protein efficiency ratio (X) correlation: $Y = 5.10 + 26.51X$; $R = 0.903$; $p = 0.036$.

Starch (Y)-percent weight gain (X) correlation: $Y = 37.0 + 0.05X$; $R = 0.911$; $p = 0.031$.

Values with different superscripts in each column indicate significant difference ($p < 0.05$).

¹ Values in parentheses are amylose-amylopectin ratios in diets. ² Data are means of four replicate tanks per dietary treatment.

Table 4. Effects of dietary amylose-amylopectin ratios on digestive enzyme activity in the gut of tilapia over a feeding period of 6 wks

Diets ¹	Digestive enzyme activity in gut (U/mg protein) ²		
	Protease	Lipase	Amylase
Diet 1 (0.11)	1,803 ^b	207.9 ^a	6.51 ^a
Diet 2 (0.24)	2,020 ^a	152.2 ^b	4.93 ^b
Diet 3 (0.47)	1,221 ^c	137.9 ^b	3.43 ^c
Diet 4 (0.76)	1,253 ^c	123.2 ^{bc}	3.09 ^c
Diet 5 (0.98)	1,361 ^c	104.2 ^c	3.24 ^c
Pooled SE	76.21	8.50	0.35
p-value	<0.001	<0.001	<0.001

Lipase (Y)-amylose-amylopectin ratio (X) correlation: $Y = 195.94 - 99.34X$; $R = -0.9080$; $p = 0.0330$.

Amylase (Y)-amylose-amylopectin ratio (X) correlation: $Y = 6.04 - 3.51X$; $R = -0.8617$; $p = 0.0604$.

Amylase (Y)-starch digestibility (X) correlation: $Y = -15.89 + 0.3X$; $R = 0.910$; $p = 0.032$.

Values with different superscripts in each column indicate significant difference ($p < 0.05$).

¹ Values in parentheses are amylose-amylopectin ratios in diets. ² Data are means of four replicate tanks per dietary treatment.

response to dietary amylose-amylopectin ratio (Table 4).

related to the ratio ($R = -0.9642$, $p = 0.0081$).

Whole-body composition

The values of moisture, crude protein, crude fat and ash of the fish did not vary across dietary treatments (Table 5). In case of the relationship between body lipid and dietary amylose-amylopectin ratio, lipid retention was inversely

Plasma components and liver glycogen

There were no significant differences in plasma glucose, TG, T-CHO, LDL-C, and HDL-C concentrations between dietary treatments (Table 6). However, liver glycogen content showed significant difference between different

Table 5. Effects of different dietary amylose-amylopectin ratios on whole body composition of tilapia over a feeding period of 6 wks

Diets	Whole-body composition (%) ¹			
	Moisture ²	Crude protein ²	Crude fat ²	Ash ²
Diet 1 (0.11)	74.03	12.68	8.02	3.66
Diet 2 (0.24)	73.83	13.11	7.78	3.66
Diet 3 (0.47)	72.96	13.66	7.68	3.93
Diet 4 (0.76)	74.77	13.07	6.89	4.11
Diet 5 (0.98)	74.47	13.39	6.86	3.86
Pooled SE	0.398	0.212	0.217	0.089
p-value	0.702	0.705	0.326	0.495

Crude fat (Y)-amylose-amylopectin ratio (X) correlation: $Y = 8.18 - 1.44X$; $R = -0.9642$; $p = 0.0081$.

Values with different superscripts in each column indicate significant difference ($p < 0.05$).

¹ Crude protein, crude fat and ash are expressed on a wet weight basis. ² Data are means of four replicate tanks per dietary treatment.

Table 6. Effects of dietary amylose-amylopectin ratios on the concentrations of plasma components and liver glycogen in tilapia over a feeding period of 6 wks

Diets ¹	Plasma components (mmol/L) ²					Liver glycogen (mg/g)
	Glucose	TG	T-CHO	LDL-C	HDL-C	
Diet 1 (0.11)	7.22	4.75	5.19	2.21	2.42	70.3 ^b
Diet 2 (0.24)	7.36	4.70	5.79	2.56	2.72	56.7 ^b
Diet 3 (0.47)	6.45	4.58	5.56	2.52	2.61	54.6 ^b
Diet 4 (0.76)	6.17	4.42	4.50	2.14	2.57	98.3 ^a
Diet 5 (0.98)	6.87	4.53	4.43	2.38	2.63	97.7 ^a
Pooled SE	0.132	0.037	0.151	0.047	0.034	4.856
p-value	0.121	0.339	0.158	0.627	0.983	0.003

TG (Y)-amylose-amylopectin ratio (X) correlation: $Y = 4.470 - 0.135 \log(X)$; $R = 0.9054$; $p = 0.0344$.

Values with different superscripts in each column indicate significant difference ($p < 0.05$).

TG = Triglyceride; T-CHO = Total cholesterol; HDL-C = High-density lipoprotein cholesterol; LDL-C = Low-density lipoprotein cholesterol, LDL-C.

¹ Values in parentheses are amylose-amylopectin ratios in diets. ² Data are means of four replicate tanks per dietary treatment.




Figure 1. Effect of amylose-amylopectin ratios on the plasma glucose concentrations in the fish at different diets. Plasma glucose concentrations were measured at various time intervals after a meal in fish fed diets varying in amylose-amylopectin ratios (0.11, 0.24, 0.47, 0.76 and 0.98, respectively) over the 6-wk feeding period. Data are means±SD (n = 3), represents three replicate tanks used at each sampling time. Plasma samples of three fish were pooled for tank. Plasma glucose was significantly affected by dietary treatments and time after a meal.

dietary treatments ($p < 0.05$), and the highest value was seen in fish fed diet 4 and lowest in fish fed diet 3.

Postprandial blood metabolic response

Plasma glucose concentrations of fish fed diets 1, 2 and 5 reached a peak 3 h after a meal (Figure 1), but the peak was reached in fish fed diets 3-4 only at 6 h, and thereafter the values gradually decreased to about the preprandial levels till 24 h. However, the fish fed diet 1 exhibited a significant difference ($p < 0.05$), when compared with the lower peaks achieved in diets 3-5.

A decreasing trend was observed with the increase in dietary amylose-amylopectin ratio, and the fish fed diet 1, showed the highest value, which was significantly different ($p < 0.05$) from the fish fed diets 3-5 at 1, 3 or 9 h, while lowest value was observed in the fish fed diet 5. Highest peak was observed in fish fed diets 1-3 and diet 5 3 h after a meal, but it reached its peak in fish fed diet 4 only after 6 h (Figure 2). However, the peak in fish fed diet 1 was significantly ($p < 0.05$) higher than fish fed diets 3 and 5 at 3 h. The values in fish fed diet 1 maintained at a higher level than those of fish fed other diets for the 9 h after a meal.




Figure 2. Effect of amylose-amylopectin ratios on the plasma triglyceride concentrations in the fish at different diets. Plasma triglyceride concentrations were measured at various time intervals after a meal in fish fed diets varying in amylose-amylopectin ratios (0.11, 0.24, 0.47, 0.76 and 0.98, respectively) over the 6-wk feeding period. Data are means±SD (n = 3), represents three replicate tanks used at each sampling time. Plasma samples of three fish were pooled for tank. Plasma triglyceride was significantly affected by dietary treatments and time after a meal.




Figure 3. Effect of amylose-amylopectin ratios on the plasma total amino acid concentrations in the fish at different diets. Plasma total amino acid concentrations were measured at various time intervals after a meal in fish fed diets varying in amylose-amylopectin ratios (0.11, 0.24, 0.47, 0.76 and 0.98, respectively) over the 6-wk feeding period. Data are means \pm SD (n = 3), represents three replicate tanks used at each sampling time. Plasma samples of three fish were pooled for tank. Plasma total amino acid was significantly affected by dietary treatments and time after a meal.

The values across all treatments were reduced to below the preprandial value till 24 h.

Contrary to the plasma glucose changes, total amino acid concentrations of fish fed diets 1 and 2 reached a peak 6 h after a meal, whereas the peak of fish fed diets 3-5 was reached at 3 h (Figure 3). The peak of fish fed diet 5 at 3 h was at the lowest and was not different from the initial value and those at other sampling times.

DISCUSSION

The results of our study demonstrated that amylose-amylopectin ratio in the starch fraction of the diet affects the growth performance of the tilapia. The fish fed with the diet containing amylose-amylopectin at ratio of 0.24 had superior performance (WG% and FE%) relative to those fed the diets at higher or lower ratio, which was not accordance with the results of Rawles and Lochmann (2003) who observed enhanced growth of sunshine bass that were fed a 70% amylose diet. Our finding is consistent with what has been reported for broilers (Dai et al., 2008) and piglets (Deng et al., 2010) wherein the dietary amylose-amylopectin ratio at 0.23 is suitable for this production. The observation in our study and the previous reports are closer to the average value of amylose-amylopectin in cereal starch fraction (Shibanuma et al., 1994; Hruby, 2005). The digestion of starch depends on its amylose-amylopectin ratio and their molecular weight (Granfeldt et al., 1994; Englyst and Hudson, 1996). Amylose has more hydrogen bonds and undergoes more rapid retrogradation than does amylopectin (Zhou and Kaplan, 1997), while the amylopectin, having a branched chain, allows a greater

access to digestive enzymes. For this reason, amylose is more resistant to hydrolytic enzymes compared with amylopectin (Kabir et al., 1998; Weurding et al., 2001), which may be partly attributed to the presence or development of resistance starch (Behall and Hallfrisch, 2002). Therefore, the ratio of amylose to amylopectin affects the digestion of starch and other macronutrients and its metabolic responses (Le Leu et al., 2009; Deng et al., 2010; Fugui et al., 2010; Jun et al., 2010). In the present study, a linear correlation was found between amylase activity and starch digestibility in response to dietary amylose-amylopectin ratio, which reflects the direct impact of dietary amylose-amylopectin ratio on starch digestion at a higher level. Furthermore, FE%, PER or/and WG% correlated linearly with the apparent digestibility of dry matter and starch. Our results agreed with what was reported by Denardin et al. (2012) for rats. However, the mismatch between protease activity and protein digestibility and between lipase activity and lipid digestibility in response to dietary amylose-amylopectin ratio indicated that protein and lipid digestibility was less affected by starch type compared with starch digestibility. This indicates that nutrient digestibility is not the only reason for the differences in starch utilization in terms of starch type. Previous studies showed that rats and poultry were able to adapt to dietary changes by increasing their feed intake and dietary energy consumption to compensate the lower digestibility values of diets caused by starch type (Zhou and Kaplan, 1997; Pirgozliev et al., 2010). Similar responses also exist in some carnivorous fish (Bergot, 1979; Yamamoto et al., 2001; Boujard et al., 2004), that is, such fish have the ability to compensate for imbalances in

nutrient density through increasing feed intake. The tilapia, however, exhibited a partial ability in this regard in our current study, because although the feed intake increased with an increasing dietary ratio of amylose-amylopectin, the fish fed higher amylose still exhibited lower nutrient digestibility and growth. Berger and Halver (1987) showed that liver enlargement is associated with increased lipid or glycogen deposition. An increased HSI was found in high amylose-fed rats (Zhou and Kaplan, 1997). However, the opposite result was observed in high amylose-fed sunshine bass (Rawles and Lochmann, 2003), consistent with our current results. This indicates that liver was in a good state, though liver glycogen concentration was higher in fish fed a diet at a high amylose-amylopectin ratio compared with those at the low ratio. In the present study, liver glycogen deposition indicates that starch not used for energy may accumulate in the liver of the tilapia fed higher-amylose-amylopectin ratio diets, supporting the observations for rainbow trout *Oncorhynchus mykiss* (Brauge et al., 1994). These observations may partly account for the inferior growth and feed utilization of fish that were fed higher-amylose diets.

The whole-body composition was not affected by dietary treatments, which was similar to the observations by Solomon et al. (2007). Although not significant, there was a progressive reduction in whole-body lipid retention to dietary amylose-amylopectin ratio. The inverse relationship between dietary amylose-amylopectin ratio and whole-body lipid retention appears interesting, since an increased dietary amylose level did not lead to undesirable lipid accumulation in the body of the fish, though the whole-body protein retention was not related with the dietary amylose-amylopectin ratio.

Rawles and Lochmann (2003) reported a reduced liver glycogen deposit in higher amylose-fed sunshine bass. The opposite response was observed in our current study and in previous work with gibel carp (*Carassius auratus gibelio*; Tan et al., 2006). Regardless of starch type, high dietary starch levels could elevate liver glycogen deposits in rainbow trout (Bergot, 1979; Suárez et al., 2002), eel (*Angilla Anguilla*; Suárez et al., 2002) and Atlantic cod (*Gadus morhua*; Hemre et al., 1993), indicating the relative low ability of these carnivorous fish to use glucose by the glycolytic pathway. The variations may be caused by the difference in fish species.

The plasma intermediary metabolites were not affected by dietary treatments, which is consistent with those of Camp et al. (2003), who reported no effect of starch type on plasma metabolites (TG, LDL-C and HDL-C) in growing pigs. However, decreased serum T-CHO and TG levels in mammals was observed by Behall et al. (1988) and Denardin et al. (2012), whereas the opposite effect was observed by Yin et al. (2011) in weaned pigs after the

consumption of a diet rich in amylose compared to a diet rich in amylopectin. The reason for this difference is unclear in that the dietary starch level was identical for each experiment besides protein and lipid level. Some previous studies have also shown that starch type is not the sole determinant factor affecting starch digestion and postprandial glycemic responses (Zhou and Kaplan, 1997) as well as growth performance (Pirgozliev et al., 2010), but there are other physiochemical characteristics of cereals like genotype (Granfeldt et al., 1994; Gutierrez Del Alamo et al., 2009), processing (Mahadevamma and Tharanathan, 2007; Glencross et al., 2012), hardness (Denardin et al., 2012) and non-starch polysaccharides (Glencross et al., 2012), which could also exert some effect on the growth.

Because of the high enzyme resistance of amylose, mammals ingesting high dietary amylose have low glycemic and insulinemic postprandial responses compared with amylopectin (Granfeldt et al., 1994; Noakes et al., 1996; Zhou and Kaplan, 1997; Behall and Hallfrisch, 2002; Aziz et al., 2009). This effect is beneficial for the control of diabetes and also to reduce the risk of diabetes development (Behall and Hallfrisch, 2002; Aziz et al., 2009). Contrary to terrestrial animals, tolerance to carbohydrate is somewhat lower in fish (Wilson, 1994; Enes et al., 2010), and its underlying mechanism remains unclear (Moon, 2001; Polakof et al., 2010). Few studies have observed fish glycemic changes in response to amylose-amylopectin ratio. In the present study, we measured the postprandial changes in several blood intermediary metabolites of fish fed diets varying in amylose-amylopectin ratios. The weak peak level of blood glucose in the tilapia in our study and in sunshine bass in the previous study conducted by Rawles and Lochmann (2003) was observed after the high-amylose load when compared with those with the low-amylose load, a fact suggesting that feeding higher-amylose diet might lead to higher tolerance to starch. Meanwhile, peak blood TG paralleled with the glucose change of the tilapia in response to dietary amylose-amylopectin ratio. In contrast with peak blood glucose, peak total amino acid time was delayed when fish were fed the low-amylose diet when compared with those fed the high-amylose diet. We also observed that the highest dietary amylose-amylopectin ratio produced the weakest metabolic responses, regardless of blood glucose, TG or total amino acids. The lowered metabolic responses in the presence of high-amylose starch could be associated with a lowered rate of amylolysis and this may be due to the increased time of gastrointestinal transit after ingestion of high-amylose (Goddard et al., 1984; Granfeldt et al., 1994; Weurding et al., 2001).

CONCLUSIONS

There is considerable variation in growth and feed

utilization with the dietary use of starch type in terrestrial animals and fish. Our results in the present study showed that there were negative linear relationships between percent weight gain, feed efficiency, protein efficiency ratio, starch digestibility, the activities of protease, lipase and amylase and the dietary amylose-amylopectin ratio. After a meal, peak blood glucose and TG were lower in fish fed higher-amylose diets than those in fish fed lower-amylose diets, and peak total amino acid time was delayed in the fish fed the low-amylose diet when compared with those fed the high-amylose diet. The decreased growth performance, along with the lower-peak blood glucose and TG as well as the delayed peak total amino acid time in fish fed higher-amylose-amylopectin diets let us to conclude that higher dietary amylose compromises the growth but helps in reducing the blood glucose stress on the fish caused by postprandial starch load.

ACKNOWLEDGEMENTS

This study was funded by the Fujian Provincial Natural Science Foundation (2012J01141), the Foundation for Innovative Research Team of Jimei University (2011A001), and the Research Foundation of Jimei University (F04008).

REFERENCES

- AOAC. 1995. Association of Official Analytical Chemists. In: Official Methods of Analysis (Ed. P. A. Cunniff). 16th edn. AOAC International, Arlington, VA, USA.
- Aziz, A. A., L. S. Kenney, B. Goulet, and E. Abdel-Aal. 2009. Dietary starch type affects body weight and glycemic control in freely fed but not energy-restricted obese rats. *J. Nutr.* 139: 1881-1889.
- Behall, K. M. and J. Hallfrisch. 2002. Plasma glucose and insulin reduction after consumption of breads varying in amylose content. *Eur. J. Clin. Nutr.* 56: 913-920.
- Behall, K. M., D. J. Scholfield, and J. Canary. 1988. Effect of starch structure on glucose and insulin responses in adults. *Am. J. Clin. Nutr.* 47:428-432.
- Behall, K. M., D. J. Scholfield, I. Yuhaniak, and J. Canary. 1989. Diets containing high amylose vs amylopectin starch: effects on metabolic variables in human subjects. *Am. J. Clin. Nutr.* 49:337-344.
- Berger, A., and J. E. Halver. 1987. Effect of dietary protein, lipid and carbohydrate content on the growth, feed efficiency and carcass composition of striped bass, *Morone saxatilis* (Walbaum), fingerlings. *Aquac. Res.* 18:345-356.
- Bergh, M., A. Razdan, and P. Aman. 1999. Nutritional influence of broiler chicken diets based on covered normal, waxy and high amylose barleys with or without enzyme supplementation. *Anim. Feed Sci. Technol.* 78:215-226.
- Bergot, F. 1979. Carbohydrate in rainbow trout diets: effects of the level and source of carbohydrate and the number of meals on growth and body composition. *Aquaculture* 18:157-167.
- Boujard, T., A. Gelineau, D. Coves, G. Corraze, G. Dutto, E. Gasset, and S. Kaushik. 2004. Regulation of feed intake, growth, nutrient and energy utilisation in European sea bass (*Dicentrarchus labrax*) fed high fat diets. *Aquaculture* 231: 529-545.
- Brauge, C., F. Medale, and G. Corraze. 1994. Effect of dietary carbohydrate levels on growth, body composition and glycaemia in rainbow trout, *Oncorhynchus mykiss*, reared in seawater. *Aquaculture* 123:109-120.
- Camp, L. K., L. Southern, and T. D. Bidner. 2003. Effect of carbohydrate source on growth performance, carcass traits, and meat quality of growing-finishing pigs. *J. Anim. Sci.* 81:2488-2495.
- Cohen, R. S., and T. D. Jr. Tanksley. 1973. Energy and protein digestibility of sorghum grains with different endosperm textures and starch types by growing swine. *J. Anim. Sci.* 37: 931-935.
- Collins, N. E., E. T. Moran Jr., and H. L. Stilborn. 2003. Performance of broilers fed normal and waxy corn diets formulated with chick and rooster derived apparent metabolizable energy values for the grains. *J. Appl. Poult. Res.* 12:196-206.
- Dai, Q., X. Li, S. Zhang, G. Jiang, Y. Hu, and F. Chen. 2008. Effects of different dietary amylose-amylopectin ratios on production performance and nutrient availability of yellow broilers. *Chinese J. Anim. Nutr.* 20:249-255.
- Denardin, C. C., N. Bouffleur, P. Reckziege, L. P. da Silva, and M. Walter. 2012. Amylose content in rice (*Oryza sativa*) affects performance, glycemic and lipidic metabolism in rats. *Ciênc. Rural* 42:381-387.
- Deng, J., X. Wu, S. Bin, T. J. Li, R. Huang, Z. Liu, Y. Liu, Z. Ruan, Z. Deng, Y. Hou, and Y. L. Yin. 2010. Dietary amylose and amylopectin ratio and resistant starch content affects plasma glucose, lactic acid, hormone levels and protein synthesis in splanchnic tissues. *J. Anim. Physiol. Anim. Nutr.* 94:220-226.
- Enes, P., S. Panserat, S. Kaushik, and A. Oliva-Teles. 2009. Nutritional regulation of hepatic glucose metabolism in fish. *Fish Physiol. Biochem.* 35:519-539.
- Enes, P., H. Peres, A. Couto, and A. Oliva-Teles. 2010. Growth performance and metabolic utilization of diets including starch, dextrin, maltose or glucose as carbohydrate source by gilthead sea bream (*Sparus aurata*) juveniles. *Fish Physiol. Biochem.* 36:903-910.
- Englyst, H. N., and G. J. Hudson. 1996. The classification and measurement of dietary carbohydrates. *Food Chem.* 57:15-21.
- Ertl, D., and N. Dale. 1997. The metabolizable energy of waxy vs. normal corn for poultry. *J. Appl. Poult. Res.* 6:432-435.
- Fugui, Y., Z. Zhenzhen, H. Ju, and Y. Yulong. 2010. Digestion rate of dietary starch affects systemic circulation of amino acids in weaned pigs. *Br. J. Nutr.* 103:1404-1412.
- Glencross, B., D. Blyth, S. Tabrett, N. Bourne, S. Irvin, M. Anderson, T. Fox-Smith, and R. Smullen. 2012. An assessment of cereal grains and other starch sources in diets for barramundi (*Lates calcarifer*) – implications for nutritional and functional qualities of extruded feeds. *Aquacult. Nutr.* 18: 388-399.
- Goddard, M. S., G. Young, and R. Marcus. 1984. The effect of amylose content on insulin and glucose responses to ingested rice. *Am. J. Clin. Nutr.* 39:388-392.
- Granfeldt, Y., A. Drews, and I. Björck. 1995. Arepas made from

- high amylose corn flour produce favorably low glucose and insulin response in healthy humans. *J. Nutr.* 125:459-465.
- Granfeldt, Y., H. Liljeberg, A. Drews, R. Newman, and I. Björck. 1994. Glucose and insulin responses to barley products: influence of food structure and amylose-amylopectin ratio. *Am. J. Clin. Nutr.* 59:1075-1082.
- Gutierrez Del Alamo, A., M. W. A. Verstegen, L. A. den Hartog, P. Perez de Ayala, and M. J. Villamide. 2009. Wheat starch digestion rate affects broiler performance. *Poult. Sci.* 88:1666-1675.
- Hemre, G. I., Ø. Lie, and A. Sundby. 1993. Dietary carbohydrate utilization in cod (*Gadus morhua*): metabolic responses to feeding and fasting. *Fish Physiol. Biochem.* 10:455-463.
- Hemre, G.-I., T. P. Mommsen, and Å. Krogdahl. 2002. Carbohydrates in fish nutrition: effects on growth, glucose metabolism and hepatic enzymes. *Aquacult. Nutr.* 8:175-194.
- Hruby, M. 2005. Challenge of corn variability. *Feed Int.* 26:5-11.
- Jun, H., C. Daiwen, and Y. Bing. 2010. Metabolic and transcriptomic responses of weaned pigs induced by different dietary amylose and amylopectin ratio. *PLoS One.* 5:e15110.
- Kabir, M., S. W. Rizkalla, M. Champ, J. Luo, J. Boillot, F. Bruzzo, and G. Slama. 1998. Dietary amylose-amylopectin starch content affects glucose and lipid metabolism in adipocytes of normal and diabetic rats. *J. Nutr.* 128:35-43.
- Krogdahl, Å., G.-I. Hemre, and T. P. Mommsen. 2005. Carbohydrates in fish nutrition: digestion and absorption in postlarval stages. *Aquac. Nutr.* 11:103-122.
- Kyuma, T., and M. Ishida. 1989. Growth and digestion of goats fed high-amylose corn and waxy corn. *Asian-Aust. J. Anim. Sci.* 2:359-360.
- Le Leu, R. K., Y. Hu, I. L. Brown, and G. P. Young. 2009. Effect of high amylose maize starches on colonic fermentation and apoptotic response to DNA-damage in the colon of rats. *Nutr. Metab.* 6:11.
- Mahadevamma, S., and R. N. Tharanathan. 2007. Processed rice starch characteristics and morphology. *Eur. Food Res. Technol.* 225:603-612.
- Mommsen, T. P., and E. M. Plisetskaya. 1991. Insulin in fishes and agnathans: history, structure and metabolic regulation. *Rev. Aquat. Sci.* 4:225-259.
- Mongeau, R., G. Sarwar, R. Brassard, and H. G. Botting. 1995. Effects of amylose and wheat bran on the levels of blood serum urea nitrogen (BUN), other blood parameters, growth and fecal characteristics in rats. *Plant Foods Hum. Nutr.* 48:95-105.
- Moon, T. W. 2001. Glucose intolerance in teleost fish: fact or fiction? *Comp. Biochem. Physiol. B, Biochem. Mol. Biol.* 129:243-249.
- Noakes, M., P. M. Clifton, P. J. Nestel, R. L. Leu, and G. McIntosh. 1996. Effect of high-amylose starch and oat bran on metabolic variables and bowel function in subjects with hypertriglyceridemia. *Am. J. Clin. Nutr.* 64:944-951.
- Perez, J. M., and A. Aumaitre. 1979. Waxy versus regular maize: Energy value for growing pigs and utilization in piglet diets. *Anim. Feed Sci. Technol.* 4:109-115.
- Pirgozliev, V. R., S. P. Rose, and M. R. Bedford. 2010. The effect of amylose-amylopectin ratio in dietary starch on growth performance and gut morphology in broiler chickens. *Arch. Geflügelkd.* 74:S21-S29.
- Pirgozliev, V. R., S. P. Rose, and R. A. Graybosch. 2002. Energy and amino acid availability to chickens of waxy wheat. *Arch. Geflügelkd.* 66:108-113.
- Pirgozliev, V. R., S. P. Rose, P. S. Kettlewell, and M. R. Bedford. 2000. Relationship between chemical composition of wheat and broiler chicken growth performance. *Brit. Poult. Sci.* 41: S697-S698.
- Polakof, S., R. Álvarez, and J. L. Soengas. 2010. Gut glucose metabolism in rainbow trout: implications in glucose homeostasis and glucosensing capacity. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.* 299:R19-R32.
- Rawles, S., and R. Lochmann. 2003. Effects of amylopectin-amylose starch ratio on growth, body composition and glycemic response of sunshine bass *Morone chrysops* ♀ × *M. saxatilis* ♂. *J. World Aquacult. Soc.* 34:278-288.
- Shelton, J. L., J. O. Matthews, L. L. Southern, A. D. Higbie, T. D. Bidner, J. M. Fernandez, and J. E. Pontif. 2004. Effect of nonwaxy and waxy sorghum on growth, carcass traits, and glucose and insulin kinetics of growing-finishing barrows and gilts. *J. Anim. Sci.* 82:1699-1706.
- Shiau, S.-Y., and M.-J. Chen. 1993. Carbohydrate utilization by tilapia (*Oreochromis niloticus* × *O. aureus*) as influenced by different chromium sources. *J. Nutr.* 123:1747-1753.
- Shibanuma, K., Y. Takeda, S. Hizukuri, and S. Shibata. 1994. Molecular structure of some wheat starches. *Carbohydr. Polym.* 25:111-116.
- Solomon, S. G., L. O. Tihamiyu, and U. J. Agaba. 2007. Effect of feeding different grain sources on the growth performance and body composition of tilapia, (*Oreochromis niloticus*) fingerlings fed in outdoor hapas. *Pakistan J. Nutr.* 6:271-275.
- Spyridakis, P., R. Metailler, J. Gabaudan, and A. Riaza. 1989. Studies on nutrient digestibility in European sea bass (*Dicentrarchus labrax*): 1. Methodological aspects concerning faeces collection. *Aquaculture* 77:61-70.
- Stone, D. A. 2003. Dietary carbohydrate utilization by fish. *Rev. Fish. Sci.* 11:337-369.
- Suárez, M. D., A. Sanz, J. Bazoco, and M. García-Gallego. 2002. Metabolic effects of changes in the dietary protein: carbohydrate ratio in eel (*Angilla anguilla*) and trout (*Oncorhynchus mykiss*). *Aquacult. Int.* 10:143-156.
- Tan, Q., S. Xie, X. Zhu, W. Lei, and Y. Yang. 2006. Effect of dietary carbohydrate sources on growth performance and utilization for gibel carp (*Carassius auratus gibelio*) and Chinese longsnout catfish (*Leiocassis longirostris* Günther). *Aquacult. Nutr.* 12:61-70.
- Weststrate, J. A., and J. M. M. van Amelsvoort. 1993. Effects of the amylose content of breakfast and lunch on postprandial variables in male volunteers. *Am. J. Clin. Nutr.* 58:180-186.
- Weurding, R. E., A. Veldman, W. A. G. Veen, P. J. van der Aar, and M. W. A. Verstegen. 2001. *In vitro* starch digestion correlates well with rate and extent of starch digestion in broiler chickens. *J. Nutr.* 131:2336-2342.
- Wilson, R. P. 1994. Utilization of dietary carbohydrate by fish. *Aquaculture* 124:67-80.
- Yamamoto, T., T. Shima, H. Furuita, N. Suzuki, F. J. Sánchez-Vázquez, and M. Tabata. 2001. Self-selection and feed consumption of diets with a complete amino acid composition and a composition deficient in either methionine or lysine by rainbow trout *Oncorhynchus mykiss* (Walbaum). *Aquac. Res.*

32 (Suppl. 1):83-91.

- Ye, J., K. Wang, F. Li, and Y. Sun. 2011. Single or combined effects of fructo- and mannan oligosaccharide supplements and *Bacillus clausii* on the growth, feed utilization, body composition, digestive enzyme activity, innate immune response and lipid metabolism of the Japanese flounder *Paralichthys olivaceus*. Aquac. Nutr. 17:e902-e911.
- Yin, F., Y. Yin, Z. Zhang, M. Xie, J. Huang, R. Huang, and T. Li. 2011. Digestion rate of dietary starch affects the systemic circulation of lipid profiles and lipid metabolism-related gene expression in weaned pigs. Br. J. Nutr. 106:369-377.
- Zhou, X., and M. L. Kaplan. 1997. Soluble amylose cornstarch is more digestible than soluble amylopectin potato starch in rats. J. Nutr. 127:1349-1356.
- Zobel, H. F. 1988. Molecules to granules: a comprehensive starch review. Starch-Stärke 40:44-50.