



Effect of FTO Expression and Polymorphism on Fat Deposition in Suzhong Pigs

Yanfeng Fu^a, Lan Li^{1,2,a}, and Shouwen Ren*

Institute of Animal Science, Jiangsu Academy of Agricultural Sciences, 210014 Nanjing, China

ABSTRACT: Fat mass and obesity associated gene (FTO) plays an important role in appetite control and energy consumption in human and mice. In order to examine FTO expression influence on fat deposition in Suzhong pigs, FTO mRNA expression was detected in 16 tissues by RT-PCR, FTO protein expression was detected in 5 tissues by western blot, and association of FTO polymorphism with meat quality traits was analyzed in Suzhong populations with 714 records. RT-PCR results revealed that FTO mRNA was expressed in all sixteen tissues with significant differences ($p < 0.05$), expression in backfat was significantly higher than that of any other tissue ($p < 0.05$), and expression in longissimus dorsi muscle had the second highest significance level ($p < 0.05$). Western blot results demonstrated that FTO protein was highly expressed in backfat and longissimus dorsi muscle. Furthermore, FTO mRNA and protein expression in tissues of high-fat pigs was significantly higher than that of low-fat pigs ($p < 0.05$), suggesting FTO expression had advantageous effects on fat deposition. FTO polymorphism results evidenced that at A227G locus, G allele seemed to have advantageous effects on fat deposition, indicating it could be a significant candidate gene for improving pork quality in Suzhong pigs. (**Key Words:** FTO, Fat Deposition, Expression, Polymorphism, Suzhong Pig)

INTRODUCTION

Pork quality is an increasing concern for swine producers due to the development of export markets and increased consumer demands (Van Wijk et al., 2005; 2006). Fat deposition, whether in adipose tissue or muscle, contributes importantly to various aspects of meat quality and are central to the nutritional value of meat (Wood et al., 2008), and fat deposition could influence some economic meat quality traits, such as backfat thickness, Intramuscular fat (IMF) content and lean meat percentage. It was reported that each type of fat deposition can have a different developmental profile (Kouba et al., 1999; Hausman and Poulos, 2004). Additionally, obesity in humans is an increasing problem worldwide, and research on candidate genes in good animal models is urgently required. The pig is an excellent model as its metabolism,

organ size, and eating habits resemble humans (Madsen et al., 2009). Therefore, for these reasons it's urgent to study the molecular mechanism of fat deposition in pigs.

Many major genes affecting meat quality traits in pigs have been successfully identified with the development of candidate gene and comparative mapping approaches, such as fatty acid-binding protein (FABP) (Gerbens et al., 2000) and proliferator-activated receptors (PPARs) (Mandard et al., 2004), Lipoprotein lipase (LPL) (Shen et al., 2010). Fat mass and obesity associated gene (FTO) was discovered in 2007, and FTO SNPs were associated with obesity of children and adults. A genome-wide study found that the FTO gene was located on chromosome 6 (Du et al., 2008). The region on chromosome 6 was first indicated as the site of the FTO gene as quantitative trait loci (QTLs) linked to regulation of some fat traits in pigs were mapped to the region (Hu and Reecy, 2007). In addition, swine FTO polymorphism was associated with pork quality, such as intramuscular fat deposition, analysis (Dina et al., 2007; Frayling et al., 2007). In humans and mice, some studies reported that FTO played an important role in appetite control and energy consumption (Klötting et al., 2008; Fischer et al., 2009; Olszewski et al., 2009).

Suzhong pig is a new breed with excellent reproductive performance and meat quality originating from Taihu pig.

* Corresponding Author: Shouwen Ren. Tel: +86-25-84390349, Fax: +86-25-84390349, E-mail: shouwenren@gmail.com

¹ College of Veterinary Medicine, China Agricultural University, 100193 Beijing China.

² National Research Center of Veterinary Biologicals engineering and technology, 210014 Nanjing China.

^a Yanfeng Fu and Lan Li contributed equally to this work.

Submitted Jan. 16, 2013; Accepted Apr. 9, 2013; Revised May 26, 2013

Table 1. Meat quality traits related with fat deposition in Suzhong pigs for FTO expression trial (Mean±SE)

Item	Longissimus dorsi muscle area	Lean meat percentage	Backfat thickness			Marbling score
			Shoulder at the thickest joint	Thoracic and lumbar spine junction	Lumbar and sacral spine junction	
Low-fat pigs	61.86±2.46 ^a	56.62±1.35 ^a	3.9±0.53 ^a	2.4±0.46	2.1±0.48 ^a	4.71±1.12
High-fat pigs	48.52±3.57 ^b	49.42±2.39 ^b	5.2±1.21 ^b	2.7±0.56	3.2±0.8 ^b	4.72±1.33

Values with different superscripts show significant levels within columns (^{a,b} p<0.05).

The objective of this study is to examine the role of FTO expression on regulation of fat deposition in Suzhong pigs. To achieve this target, we cloned the coding sequence (CDS) of swine FTO gene in Suzhong pigs, analyzed the FTO mRNA expression by RT-PCR, detected the FTO protein expression by western blot, and examined the association of FTO polymorphism with meat quality traits.

MATERIALS AND METHODS

Animals and tissues

Animals were all healthy half-siblings fed under the same conditions from the seed stock farms of Hongze Xixiang Swine Breeding Company (Jiangsu, China). In order to investigate FTO expression, ten Suzhong pigs (five high-fat pigs and five low-fat pigs) were slaughtered on the same day by electrical stunning when their body-weight was 95±2 kg at six-month age. Afterwards, 16 tissues of heart, liver, kidney, longissimus dorsi muscle, backfat, lung, stomach, bladder, spleen, uterus, cervix, ovary, large intestine, oviduct, endometrium and small intestine were collected and prepared according to the procedure proposed by Lord with minor modifications (Lord et al., 2006). For RT-PCR and western blot, these tissues were immediately saved into liquid nitrogen. For meat quality traits, longissimus dorsi muscle area was tested using a planimeter (Haguang, China), and lean meat percentage was tested using a lean meat percentage analyzer (SFK, Denmark) (Table 1).

To detect FTO polymorphism, there were 714 records of 51 pigs killed, these records contained such data as: ear number, birth date, breed, parity, longissimus dorsi muscle area and lean meat percentage. Ear tissues of pigs were collected in centrifuge tubes (1.5 mL) with 70% ethanol,

stored at 4°C until the genomic DNA of each ear sample was extracted using a Genomic DNA Rapid Isolation Kit (Sangon, China) (Fu et al., 2012b).

Primer design

Primers for swine FTO (GenBank: NM_001112692) and housekeeping gene (GAPDH) were designed by Oligo 6 software and synthesized by Introvigen company (Shanghai, China). Primer FTO_1 and FTO_2 was used in RT-PCR, and Primer FTO_3 was used in detection of FTO polymorphism. Both Primer FTO_1 and FTO_2 were designed as cross-exons primers (Table 2).

RNA extraction and RT-PCR

Total RNA was extracted from swine tissues using a modified Trizol/chloroform (Invitrogen, USA) extraction according to the protocol (Sambrook and Russell, 2001). RNA samples were quantified spectrophotometrically (Shimadzu, Japan) and the integrity was confirmed using 1.5% agarose gel.

In reverse transcriptase-polymerase chain reaction (RT-PCR), total RNA was reverse transcribed using a First Strand cDNA Synthesis Kit (Fermentas, Canada) and was carried out in a 25 µL reaction volume by two steps according to the kit manual. The amplifying reactions were carried out in a 15 µL reaction volume containing 1 µL cDNA, 0.2 mmol/L dNTPs, 1.5 mmol/L MgCl₂, 0.5 µM of each PCR primer, 1 U Taq DNA polymerase and reaction buffer (Takara, Japan). The amplifying conditions were heating at 94°C for 5 min, followed by 35 cycles of denaturation at 94°C for 30 s, annealing at 60°C for 30 s, extension at 72°C for 50 s, after that, extension at 72°C for 7 min. Amplification of the house-keeping gene GAPDH was performed as the control for the reaction. The PCR

Table 2. Primer pairs and PCR conditions used in RT-PCR and SNPs detection analysis of FTO

Primers	Sequences (5'→3')	Temperature (°C)	Product size (bp)
FTO_1	AGCAGCAGCATGAAGCGAAC GTGAGATCAAACGGCAGAGG	62	1478
FTO_2	GGTGCACCCATGCTGTGCT ACGGCCGGCATTCTGGCTTC	60	190
FTO_3	TACCCTAAGCTAATTCTCCGA GCTATAATTGTACTGACCAC	60	609
GAPDH	GGTGAAGGTCGGAGTGAACG TGGGTGGAATCATACTGGAACA	62	150

products were confirmed by electrophoresis on 1.5% agarose gel.

Real-time PCR

The Real-time PCR amplification was performed in a 20 μ L reaction volume, containing 10 μ L 2 \times Real-time PCR Master Mix (SYBR Green), 1 μ L cDNA, 1 μ L of each PCR primer (10 μ M), 1 μ L *Taq* DNA polymerase (10 u/ μ L) and 6 μ L sterilized water (Toyobo, Japan). The reaction was run on a Line-gene Fluorescence PCR detection system (Bori, China) heating at 95°C for 5 min for initial enzyme activation, followed by 45 cycles amplification including denaturation at 95°C for 20 s, annealing at 60°C for 25 s, and extension at 72°C for 30 s. Adequate precautions were given to prevent cross-contamination and negative controls were performed routinely with each reaction. The standard curves were prepared for both tested and housekeeping gene, GAPDH (GenBank: U48832).

Western blot

Frozen sections of endometrial specimens were prepared and western-blot was performed as previously described (Wong and Medrano, 2005) with minor modification. Tissues were cracked in tissue protein extracts (0.05 mol/L Tris-HCl, NaCl 8.76 mg/mL, 1% TritonX-100 and 100 μ g/mL PMSF) (Keygen, China) by vortex meter (Huxi, China). Total protein concentrations were detected using the BCA Protein Assay Kit (Keygen, China) according to the manufacturer's recommendations.

There was a 60 μ g sample separated from a 10% Tris-HCl polyacrylamide gel using electrophoresis system (Bio-Rad, America), and protein from the gel was transferred onto a single PVDF membrane (Bio-Rad, America). After rinsing in TBST for 5 min, the membrane was soaked in 5% skim milk (in TBST) for 1 h. Next, the membrane was immersed into 1:200 dilution of the primary antibody (mouse anti-human FTO mAb, Santa Cruz, USA) at 4°C for 1 h. After rinsing in TBST for 5 min, the membrane was immersed into 1:5,000 dilution of the secondary antibody (HRP) (Santa Cruz, USA) for 2 h. Finally, the membrane was colored using the DAB kit (Invitrogen, USA) and exposed using Chemiluminescence Detection Kit for HRP (Keygen, China). β -Actin was used as a housekeeping gene. Scanned images were quantified using LabWorks gray image analysis software (Fu et al., 2012a).

cSNPs detection by PCR-sequencing

For purpose of detecting single nucleotide polymorphisms in the coding sequence (cSNPs), a 609-bp cDNA sequence was scanned within Exon 3 of the swine FTO gene by PCR-sequencing technology. The

amplification reactions were performed in a 15 μ L reaction volume containing 50 ng genomic DNA, 1.5 μ L of 10 \times PCR buffer (containing 100 mM Tris-HCl (pH 8.0), 500 mM KCl, 10 mM of MgCl₂ and 0.1% Triton X-100), 200 μ M of each dNTPs, 0.5 μ L of each primer (10 pM) and 3 U of *Taq* DNA polymerase (Takara, China). The PCR conditions were heating at 94°C for 5 min, followed by 35 cycles of reactions containing denaturation at 94°C for 30 s, annealing at 59°C for 30 s, and extension at 72°C for 60 s, finally, a further heating at 72°C for 7 min. The PCR products were confirmed by electrophoresis on 1.5% agarose gel and then purified using a DNA fragment gel purification and extraction kit (Sangon, China).

Direct DNA sequencing was implemented using purified PCR products on an ABI 3730XL automated DNA sequencer (Applied Biosystems, USA) by a professional gene sequencing company (Invitrogen, Shanghai, China). Each sample was sequenced twice and all bases of both strands were sequenced. The sequence data were analyzed using DNAMAN (Version 5.2) and aligned with the consensus sequence of swine FTO sequence available in the GenBank database.

Statistical analysis

Data analysis was performed using GLM procedures of SAS (Ver. 8.2). Significant differences were established utilizing PDIFF option of GLM with least significant differences (LSD) post hoc test and assumed as statistically significant for $p \leq 0.05$, and Lsmeans \pm SE are presented. The additive and dominance effects were estimated using Estimate step of GLM procedure in SAS according to the expressions $a = (AA-BB)/2$ and $d = (AB-(AA+BB))/2$, respectively, where AA, AB, and BB are the genotypic values (Tang et al., 2010; Fu et al., 2012b).

RT-PCR data were examined by gray values ratio (FTO/GAPDH) of scanned agarose lane images by LabImage software, real-Time PCR data were analyzed using delta-delta C_T method (also known as the 2^{- $\Delta\Delta$ C_T} method) (Livak and Schmittgen, 2001; Schmittgen and Livak, 2008), GAPDH gene is as positive control.

Association analysis between FTO polymorphism and meat quality traits were evaluated with the GLM procedure of SAS (Version 8.2). Genotype effects of SNPs were analyzed by the established model (Fu et al., 2012b),

$$\text{Model: } Y_{ijklm} = \mu + HYS_i + B_j + S_k + G_l + e_{ijklm}$$

Where Y is the meat quality trait, μ is the overall mean, HYS_i is the effect of herd-year-season ($i = 1$), B_j is the effect of breed ($j = 1$), S_k is the effect of sex ($k = 1, 2$), G_l is the effect of genotype ($l = 1, 2, 3$) and e is the random residual.




Figure 1. The 1.5% agarose gel electrophoresis atlas for the expression of swine fat mass and obesity associated gene (FTO) in different tissues of Suzhong pigs, and gray values of scanned agarose lane images by LabImage software, GAPDH gene is as positive control. (A) agarose gel electrophoresis atlas, (B) Gray values of FTO/GAPDH. Values with different superscripts show significant levels within columns ($p < 0.05$).

RESULTS

FTO mRNA expression by RT-PCR

According to OD values measured by spectrophotometer, the OD260/OD280 values of total RNA extracted from swine tissues ranged from 1.8 to 2.1, and RNA content ranged from 0.60 to 2.22 $\mu\text{g}/\mu\text{L}$, suggesting the quality of RNA extracted was good and it could be used for subsequent experiments.

Expression analysis showed that FTO mRNA was expressed in 16 tissues in Suzhong pigs. Ranking by mRNA

expression intensity (FTO/GAPDH) in these tissues, from highest to lowest, it was backfat (9.44), heart (1.10), kidney (0.92), ovary (0.91), lung (0.89), oviduct (0.86), endometrium (0.83), longissimus dorsi muscle (0.81), large intestine (0.66), uterus (0.65), cervix (0.64), spleen (0.57), bladder (0.50), stomach (0.42), liver (0.20) and small intestine (0.04). Among all these tissues, mRNA expression in backfat was extremely significantly ($p < 0.01$) higher than any other tissue and expression in longissimus dorsi muscle of high-fat pigs was significantly ($p < 0.05$) higher than that of low-fat pigs (Figures 1 and 2).




Figure 2. The mRNA expression intensity of fat mass and obesity associated gene (FTO) in different tissues. Real-time PCR detected mRNA expression in heart, liver, kidney, backfat and longissimus dorsi muscle (muscle) of Suzhong pigs. GAPDH was used as the housekeeping gene. The mRNA expression intensity of FTO varied significantly in different tissues.




Figure 3. Western-blot results of FTO expression in different tissues of Suzhong pigs, including heart, liver, kidney, backfat and longissimus dorsi muscle (muscle). (A) Western blot lanes (B) Gray values of FTO/ β -actin for scanned western blot lane images. Lowercase letters on columns indicate significant levels within columns ($p < 0.05$).

This suggested that a higher FTO expression was in favor of a higher fat content, a lower longissimus dorsi muscle area and a lower lean meat percentage (Table 1), FTO expression seemed to have advantageous effects on fat deposition.

FTO protein expression by western blot

The western blot results showed that FTO protein (GenBank: NP_001106162) was expressed in heart, liver, kidney, longissimus dorsi muscle and backfat of Suzhong pigs. Ranking by FTO protein expression intensity, from highest to lowest, it was backfat, longissimus dorsi muscle, kidney, heart, liver in low-fat pigs, and its backfat, longissimus dorsi muscle, liver, kidney, heart in high-fat pigs. FTO protein expression in backfat was significantly ($p < 0.05$) higher than that of any other tissue in all pigs, and FTO protein expression in longissimus dorsi muscle was second highest (Figure 3).

FTO protein expression in tissues of high-fat pigs was extremely significantly ($p < 0.01$) higher than that of low-fat pigs, and the expression difference between high-fat pigs and low-fat pigs was consistent with the real-time PCR results (Table 3). Considering that longissimus dorsi muscle area and lean meat percentage of high-fat pigs were both significantly ($p < 0.05$) lower than that of low-fat pigs (Table 1), the results showed that a higher FTO expression might conducive to a lower longissimus dorsi muscle area and lean meat percentage, which was opposite to fat content. Consequently, FTO might be involved in regulating the swine fat deposition.

cSNP Identification

The coding sequence (CDS, 15 to 1,532 bp long) of swine FTO was cloned from Suzhong pigs (GenBank: JX873956), and there were 10 SNPs discovered, including G52A, A227G, C523T, C608G, G628A, A655G, A810G, A886G, G1002A and A1344G. Among these SNPs, the A227G (at position 227 bp of cDNA sequence, an A/G transition) within Exon 3 (609-bp long) of swine FTO was polymorphic in 51 Suzhong pigs. At A227G locus, three PCR products of each genotype were random selected and sequenced by automated sequencing to validate the results. If position 227 is A, then it was designated as the A allele, G for the G allele (Figures 4 and 5).

Association of FTO polymorphism with meat quality traits

All genotypes of FTO occurred in the Suzhong pigs, which were found to be in Hardy-Weinberg equilibrium for

Table 3. Change in protein expression intensity of fat mass and obesity associated gene (FTO) in different tissues (Mean \pm SE)

	Item	FTO/ β -actin
Tissue	Heart	0.24 \pm 0.03 ^{Bc}
	Liver	0.24 \pm 0.04 ^{Bc}
	Kidney	0.49 \pm 0.05 ^{Aa}
	Longissimus dorsi muscle	0.34 \pm 0.03 ^{Bb}
	Backfat	0.25 \pm 0.03 ^{Bc}
Item	Low-fat pig	0.22 \pm 0.02 ^A
	High-fat pig	0.40 \pm 0.03 ^B

Values with different superscripts show significant levels within columns (^{a,b} $p < 0.05$, ^{A,B} $p < 0.01$).




Figure 4. The PCR-Sequencing results for the FTO gene polymorphism, (A) comparison results using DNAMAN software, (B) Peak value figure.

```

AGATTAGCAGCAGCATGAAGCGAACCCCAACCGCCGAGGAACGAGAGCGCGAAGCTAAGAAACTGAGGCTTCTTGAAGAGCTGGAAGACACTTGGC
TTCCTTATCTGACCCCCAAAGATGATGAATTCATCAGCAGTGGCAGCTGAAATACCTAAGCTAATTCCTCGAGAAGCAGGCAGCGTCCCTGAGGGAC
TCCAAAAGAGGTTTCAAGAAGCCTTCTCGCACTGCACAAGCATGGCTGCTTATTTTCGGGACCTGGTCAGGATCCAAGGCAAGATTGTCTACGCCA
GTATCTCGCTCCTCATTGGTAAACCCCGGCTGCACCTACAAGTACCTGAACACCAGGCTCTTACGGTCCCTGGCCAGTGAAGGGCTCTGATGCAAA
GTACAATGAGGCCGAGATAGGCGCCGCTGCCAGACCTTCTCAAGCTCAACGACTACCTGCAGATTGAGACCATCCAGGCGCTGGAGGAACTCGCT
GCCAAGGAGAAAGCCAATATCGACACCGTCCCGGTGTGTATAGTCCAGATTTCCCCAGGGTCGGCATGGGGTCATCCTTTGACGGCCATGACGAGGT
GGACAGGAAGAGCAGAGCCGCTACAACCTAACTTTGTTGAATTCATGGATCCCCAGAAAATGCCGTGCCTGAAAAGAGGAGCCCTACTTTGGCATG
GGGAAGATGGCTGTGAGCTGGCATCAGATGAAAATCTGGTGGCAGGTCAGCGGTGGCAGTGTACAATTATAGCTGTGAAGGCCCTGAAGAGGAAA
GCGAGGATGATCCCCAGCTCGAAGGCAGAGATCCCGATGTGTGGCATGTTGGCTTTAAGATCTCATGGGACATAGAGACCCCTGGTTTGGCGATACCC
TTCACCGAGGAGACTGCTACTTTATGCTGGATGATCTCAATGCCACCCACCAACTGTGTTTGGCTGGTTACACCCCGGTTTAGTTCCACCCACC
GAGTGGCCGAGTGCACGGGAACCTTGGATTACATCTTACAGCGCTGCCAGTTGGCCCTGCAGAATGCTCCGTGATGAGGCGGACAGTGGTGAAGT
CTCTTTGAAATCCTTGGAGCCTGCGGTTTTGAAACAAGGAGAAGAAATCCACAACGAGGTCGAGTTTGAGTGGCTGAGACAGTTTTGGTTTCAAGGC
AATCGATACAAAAGTGCACCGATTGGTGGTGTCAACCCATGACTCAGCTGGAAGAGCTTTGGAAGAAGATGGAAGGTGCGACCCATGCTGTGCTTC
GTGAAGTTAGAAGAGAGGGGGCCCTGTGGAACAGAGCAGTGACATCCTGACTGCCATCTAGCCGTGCTACCCGCTCGCCAGAACCTGAGGAGGG
AGTGGCATGCCAGGTGCCAGTCCCGAATTGCCCGAACTCTGCCGTGTGGACCAGAAGCCAGAATGCCGGCCGTATTGGGAAAAGGATGATCCCTCCATG
CCTCTGCCGTTTGTATCTCACAGACACTGTGGCTGAACTCAGAGGTCTGCTTCTGGAAGCCAAACCCTAG
    
```

Figure 5. The coding sequence (CDS, 15-1,532 bp) of fat mass and obesity associated gene (FTO) in Suzhong pigs by cloning and sequencing, and a new GenBank accession number was obtained (GenBank: JX873956), sequence with a gray color background was the cloning sequence in Suzhong pigs.

the genotyped locus through the chi-square test ($p > 0.05$). The allele and genotype frequencies are shown in Table 4.

The data of meat quality traits were analyzed by genotype, and these traits included longissimus dorsi muscle area/weight, lean meat percentage, PH value, luminosity, “a”, “b”, meat color score, marbling score and water holding capacity. For longissimus dorsi muscle area/weight, lean meat percentage, PH value, meat color score and water holding capacity, those of Genotype AA were significantly ($p < 0.05$) higher than those of Genotype GG. In contrast, for luminosity and marbling score, those of Genotype AA were lower than those of Genotype GG significantly ($p < 0.05$). These results suggested that G allele seemed to have advantageous effects on fat deposition (Table 5).

DISCUSSION

Fat deposition could influence meat-quality traits, such as backfat thickness, longissimus dorsi muscle area, intramuscular fat (IMF) content and lean meat percentage, etc (Wood et al., 2008). These meat quality traits were all important breeding goals for pig producers, because they are closely related to pork quality, which is positively correlated with meat tenderness, juiciness, and taste (DeVol et al., 1988; Cui et al., 2011). Recently, there have been some genes reported to be associated with fat deposition, such as Apolipoprotein M (APOM) (Pan et al., 2010), diacylglycerol acyltransferase (DGAT) (Cui et al., 2011), nudix (nucleoside diphosphate linked moiety X)-type motif 6 (Nudt6) (Sun et al., 2012) as well as others. Fat mass and obesity associated gene (FTO) played an important role in appetite control and energy consumption in human and mice (Klötting et al., 2008; Fischer et al., 2009; Olszewski et al., 2009), therefore the present research studied the FTO mRNA expression in 16 tissues and protein expression in 5 tissues, examined FTO expression difference between low-fat pigs and high-fat pigs, and analyzed association of FTO

Table 4. Allele and genotype frequencies of FTO in Suzhong pigs

NO. of pigs	Genotype distribution			Allele frequencies		χ^2
	AA	GA	GG	C	T	
51	30	18	3	0.76	0.24	0.01

χ^2 (df = 2)^{0.01} = 9.21. χ^2 (df = 2)^{0.05} = 5.99.

Table 5. Effects of FTO genotype on meat quality traits (LS Means±SE)

Meat quality traits	Genotype			Additive effect	Dominant effect
	AA	GA	GG		
Longissimus dorsi muscle area/weight	0.57±0.01 ^{Aa}	0.55±0.02 ^{Aa}	0.37±0.04 ^{Bb}	1.00±0.02	0.08±0.03
Lean meat percentage	55.78±0.98 ^A	51.34±1.25 ^B	40.54±3.06 ^C	7.62±1.61	3.18±2.03
PH value	5.83±0.05 ^{ab}	5.91±0.06 ^a	5.57±0.16 ^b	0.13±0.08	0.21±0.11
Luminosity (L)	41.09±1.04 ^{Aa}	41.54±1.32 ^{Aa}	46.50±3.22 ^{Ab}	-2.71±1.70	-2.26±2.14
a (range from magenta to green)	6.35±0.46	5.29±0.58	6.15±1.42	0.09±0.75	-0.97±0.94
b (range from yellow to blue)	3.50±0.29	3.07±0.37	3.38±0.90	0.06±0.48	-0.37±0.60
Meat color score	3.55±0.11 ^{Aa}	3.54±0.14 ^{Aa}	2.69±0.35 ^{Ab}	0.43±0.18	0.42±0.23
Marbling score	4.46±0.22 ^{Aa}	4.91±0.28 ^{Aab}	6.03±0.69 ^{Ab}	-0.78±0.37	-0.34±0.46
Water holding capacity	16.69±0.77 ^{Aa}	13.60±0.98 ^{Ab}	14.11±2.38 ^{Aab}	1.29±1.26	-1.80±1.58

Longissimus dorsi muscle area/weight was values of longissimus dorsi muscle area divided by values of weight in pigs. Luminosity (L), a and b were three parameters of meat color measured using chroma meter cr400 (Konica Minolta, Japan). Among these parameters, values of L ranged from 0 to 100 (color range from black to white), values of a ranged from 127 to -128 (color range from magenta to green) and values of b ranged from 127 to -128 (color range from yellow to blue). Values of meat color score ranged from 1.0 to 6.0 (color range from white to red), and values of marbling score ranged from 1.0 to 10.0, both of them were measured using American shade guide. Values with different superscripts show significant levels within rows (^{A,C} $p < 0.01$, ^{a,b} $p < 0.05$).

polymorphism with meat quality traits in a local pig breed of China.

At the RNA level, there're 16 tissues of FTO mRNA expression in Suzhong pigs with significant differences ($p < 0.05$) by RT-PCR, ranking by mRNA expression intensity from highest to lowest, it was backfat, heart, kidney, ovary, lung, oviduct, endometrium, longissimus dorsi muscle, large intestine, uterus, cervix, spleen, bladder, stomach, liver and small intestine in turn. In Göttingen minipigs, FTO mRNA expression was detected in muscle, brain, cerebellum, hippocampus, liver, kidney and heart with significantly higher levels in brain tissues ($p < 0.05$) (Madsen et al., 2009). In sheep, FTO mRNA expression was detected in hypothalamus, pancreas, heart, kidney, adipose tissues and liver and skeletal muscle (Sébert et al., 2010). At protein level, western blot results indicated that FTO protein was detected in all five tissues of Suzhong pigs with a significantly higher expression in backfat than that of any other tissue ($p < 0.05$). Differences between mRNA and protein expression might be due to different regulation of these genes on translation level (Fu et al., 2012a).

Both FTO mRNA and protein expression in tissues of high-fat pigs was higher than that of low-fat pigs at significant levels ($p < 0.05$), and lean meat percentage of high-fat pigs was significantly lower than that of low-fat pigs ($p < 0.05$), this might suggest that FTO expression was inversely proportional to lean meat percentage, and lean meat percentage was inversely proportional to fat content, so FTO expression had advantageous effects on fat deposition. Related studies have reported that obese animals exhibited a significant upregulation of FTO mRNA abundance (Sébert et al., 2010), while in humans, allelic distributions of FTO SNPs were similarly related to alterations of energy intake in childhood (Timpson et al.,

2008; Wardle et al., 2009).

At the A227G locus, the meat quality traits of longissimus dorsi muscle area/weight, lean meat percentage, PH value, meat color score and water holding capacity were significantly higher in pigs with Genotype AA than those of genotype GG ($p < 0.05$). In contrast, for meat quality traits of luminosity and marbling score, were significantly ($p < 0.05$) lower in Genotype AA than those of Genotype GG, suggesting G allele seemed to have advantageous effects on fat deposition. In Italian Duroc pigs, FTO polymorphism was associated with fat deposition traits (Fontanesi et al., 2009). In an Italian Duroc population and a Berkshire×Yorkshire population, two SNPs were detected in Exon 3 and Intron 4 of swine FTO, and the mutation effects on obesity-related traits were also confirmed (Fontanesi et al., 2010). The cSNP in Exon 3 of FTO did not directly alter the amino acid residue, which indicates that it was possible the polymorphism indirectly affects reproductive traits by being in linkage disequilibrium with another polymorphism that directly influences the analyzed quantitative traits (Niu et al., 2009; Fu et al., 2012b).

In conclusion, the study showed both FTO mRNA and protein expression in tissues of high-fat pigs was significantly higher than that of low-fat pigs, and at A227G locus, the G allele seemed to have advantageous effects on fat deposition. This suggested that FTO might be involved in fat deposition mechanisms, and would be one of the major genes affecting meat quality traits.

ACKNOWLEDGEMENTS

This research was supported by National Natural Science Foundation of China (No. 31201767), Postdoctoral Research Funding Project of Jiangsu Province (No.

026056511104), Jiangsu Agricultural Sciences and Technology Autonomy Innovation Fund Project (NO. CX(13)5042) and Nanjing Station Foundation of National Modern Swine Industrial-Technology System (No. nycytx-009). These authors greatly appreciate Hongze Xinxiang Swine Breeding Company (Jiangsu, China) for assistance in collecting experimental materials.

REFERENCES

- Cui, J., Y. Zeng, H. Wang, W. Chen, J. Du, Q. Chen, Y. Hu, and L. Yang. 2011. The effects of DGAT1 and DGAT2 mRNA expression on fat deposition in fatty and lean breeds of pig. *Livest. Sci.* 140:292-296.
- DeVol, D., F. McKeith, P. J. Bechtel, J. Novakofski, R. Shanks, and T. Carr. 1988. Variation in composition and palatability traits and relationships between muscle characteristics and palatability in a random sample of pork carcasses. *J. Anim. Sci.* 66:385-395.
- Dina, C., D. Meyre, S. Gallina, E. Durand, A. Körner, P. Jacobson, L. M. S. Carlsson, W. Kiess, V. Vatin, and C. Lecoeur. 2007. Variation in FTO contributes to childhood obesity and severe adult obesity. *Nat. Genet.* 39:724-726.
- Du, Z. Q., B. Fan, X. Zhao, R. Amoako, and M. F. Rothschild. 2008. Association analyses between type 2 diabetes genes and obesity traits in pigs. *Obesity* 17:323-329.
- Fischer, J., L. Koch, C. Emmerling, J. Vierkotten, T. Peters, J. C. Brüning, and U. Ruther. 2009. Inactivation of the Fto gene protects from obesity. *Nature* 458:894-898.
- Fontanesi, L., E. Scotti, L. Buttazzoni, S. Dall'Olio, A. Bagnato, D. P. Lo Fiego, R. Davoli, and V. Russo. 2010. Confirmed association between a single nucleotide polymorphism in the FTO gene and obesity-related traits in heavy pigs. *Mol. Biol. Rep.* 37:461-466.
- Fontanesi, L., E. Scotti, L. Buttazzoni, R. Davoli, and V. Russo. 2009. The porcine fat mass and obesity associated (FTO) gene is associated with fat deposition in Italian Duroc pigs. *Anim. Genet.* 40:90-93.
- Frayling, T. M., N. J. Timpson, M. N. Weedon, E. Zeggini, R. M. Freathy, C. M. Lindgren, J. R. B. Perry, K. S. Elliott, H. Lango, and N. W. Rayner. 2007. A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* 316:889-894.
- Fu, Y., J. Fu, Q. Ren, X. Chen, and A. Wang. 2012a. Expression of Eph A molecules during swine embryo implantation. *Mol. Biol. Rep.* 39:2179-2185.
- Fu, Y., J. Fu, and A. Wang. 2012b. Association of EphA4 polymorphism with swine reproductive traits and mRNA expression of EphA4 during embryo implantation. *Mol. Biol. Rep.* 39:2689-2696.
- Gerbens, F., D. De Koning, F. Harders, T. Meuwissen, L. Janss, M. Groenen, J. Veerkamp, J. Van Arendonk, and M. Te Pas. 2000. The effect of adipocyte and heart fatty acid-binding protein genes on intramuscular fat and backfat content in Meishan crossbred pigs. *J. Anim. Sci.* 78:552-559.
- Hausman, G. and S. Poulos. 2004. Recruitment and differentiation of intramuscular preadipocytes in stromal-vascular cell cultures derived from neonatal pig semitendinosus muscles. *J. Anim. Sci.* 82:429-437.
- Hu, Z. L. and J. M. Reecy. 2007. Animal QTLdb: beyond a repository. *Mamm. Genome* 18:1-4.
- Klötting, N., D. Schleinitz, K. Ruschke, J. Berndt, M. Fasshauer, A. Tönjes, M. Schön, P. Kovacs, M. Stumvoll, and M. Blüher. 2008. Inverse relationship between obesity and FTO gene expression in visceral adipose tissue in humans. *Diabetologia* 51:641-647.
- Kouba, M., M. Bonneau, and J. Noblet. 1999. Relative development of subcutaneous, intermuscular, and kidney fat in growing pigs with different body compositions. *J. Anim. Sci.* 77:622-629.
- Livak, K. and T. Schmittgen. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2- $[\Delta\Delta]CT$ method. *Methods* 25:402-408.
- Lord, E., B. Murphy, J. Desmarais, S. Ledoux, D. Beaudry, and M. Palin. 2006. Modulation of peroxisome proliferator-activated receptor δ and γ transcripts in swine endometrial tissue during early gestation. *Reproduction* 131:929-942.
- Madsen, M. B., M. M. Birck, M. Fredholm, and S. Cirera. 2009. Expression studies of the obesity candidate gene FTO in pig. *Anim. Biotechnol.* 21:51-63.
- Mandard, S., M. Müller, and S. Kersten. 2004. Peroxisome proliferator-activated receptor a target genes. *Cell. Mol. Life Sci.* 61:393-416.
- Niu, B., L. Ye, F. Li, C. Deng, S. Jiang, M. Lei, and Y. Xiong. 2009. Identification of polymorphism and association analysis with reproductive traits in the porcine RNF4 gene. *Anim. Reprod. Sci.* 110:283-292.
- Olszewski, P., R. Fredriksson, A. Olszewska, O. Stephansson, J. Alsiö, K. Radomska, A. Levine, and H. Schiöth. 2009. Hypothalamic FTO is associated with the regulation of energy intake not feeding reward. *BMC Neurosci.* 10:129-140.
- Pan, G., Y. Fu, B. Zuo, Z. Ren, D. Xu, M. Lei, R. Zheng, and Y. Z. Xiong. 2010. Molecular characterization, expression profile and association analysis with fat deposition traits of the porcine APOM gene. *Mol. Biol. Rep.* 37:1363-1371.
- Sébert, S., M. Hyatt, L. Chan, M. Yiallourides, H. Fainberg, N. Patel, D. Sharkey, T. Stephenson, S. Rhind, and R. Bell. 2010. Influence of prenatal nutrition and obesity on tissue specific fat mass and obesity-associated (FTO) gene expression. *Reproduction* 139:265-274.
- Sambrook, J. and D. W. Russell. 2001. *Molecular cloning: a laboratory manual*. Cold spring harbor laboratory press.
- Schmittgen, T. and K. Livak. 2008. Analyzing real-time PCR data by the comparative CT method. *Nat. Protoc.* 3:1101-1108.
- Shen, Y., A. Lookene, L. Zhang, and G. Olivecrona. 2010. Site-directed mutagenesis of apolipoprotein CII to probe the role of its secondary structure for activation of lipoprotein lipase. *J. Biol. Chem.* 285:7484-7492.
- Sun, L., S. Yu, H. Wang, B. Fan, and B. Liu. 2012. NUDT6, the FGF-2's antisense gene, showed associations with fat deposition related traits in pigs. *Mol. Biol. Rep.* 39:4119-4126.
- Tang, S., D. Sun, J. Ou, Y. Zhang, and G. Xu. 2010. Evaluation of the IGFs (IGF1 and IGF2) genes as candidates for growth, body measurement, carcass, and reproduction traits in Beijing You and Silkie chickens. *Anim. Biotechnol.* 21:104-113.
- Timpson, N. J., P. M. Emmett, T. M. Frayling, I. Rogers, A. T. Hattersley, M. I. McCarthy, and G. Davey Smith. 2008. The fat

- mass-and obesity-associated locus and dietary intake in children. *Am. J. Clin. Nutr.* 88:971-978.
- Van Wijk, H., D. Arts, J. Matthews, M. Webster, B. Ducro, and E. Knol. 2005. Genetic parameters for carcass composition and pork quality estimated in a commercial production chain. *J. Anim. Sci.* 83:324-333.
- Van Wijk, H., B. Dibbitts, E. Baron, A. Brings, B. Harlizius, M. Groenen, E. Knol, and H. Bovenhuis. 2006. Identification of quantitative trait loci for carcass composition and pork quality traits in a commercial finishing cross. *J. Anim. Sci.* 84:789-799.
- Wardle, J., C. Llewellyn, S. Sanderson, and R. Plomin. 2009. The FTO gene and measured food intake in children. *Int. J. Obes.* 33:42-45.
- Wong, M., and J. Medrano. 2005. Real-time PCR for mRNA quantitation. *Biotechniques* 39:75-85.
- Wood, J., M. Enser, A. Fisher, G. Nute, P. Sheard, R. Richardson, S. Hughes, and F. Whittington. 2008. Fat deposition, fatty acid composition and meat quality: A review. *Meat Sci.* 78:343-358.