

A SOLUTION OF EINSTEIN'S FIELD EQUATIONS FOR THE THIRD CLASS IN X_4

JONG WOO LEE*

ABSTRACT. The main goal in the present paper is to obtain a solution of Einstein's unified field equations for the third class in X_4 .

1. Introduction

Einstein([1]) proposed a new unified field theory that would include both gravitation and electromagnetism. Hlavatý([5]) gave the mathematical foundation of the Einstein's unified field theory in a 4-dimensional generalized Riemannian space X_4 (i.e., space-time) for the first time. Since then this theory had been generalized in a generalized Riemannian manifold X_n , the so-called *Einstein's n-dimensional unified field theory*, and many consequences of this theory has been obtained by a number of mathematicians. However, it has been unable yet to represent a general n-dimensional Einstein's connection in a surveyable tensorial form, probably due to the complexity of the higher dimensions. The purpose of the present paper is to obtain a necessary and sufficient condition for a connection with a new torsion tensor be an Einstein's connection in X_n . In the next, we obtain a solution of Einstein's field equations for the third class in X_4 . The obtained results and discussions in the present paper will be useful for the 4-dimensional considerations of the unified field theory.

2. Preliminary

Let X_n be an n-dimensional generalized Riemannian manifold covered by a system of real coordinate neighborhoods $\{U; x^\nu\}$, where, here and

Received October 31, 2012; Accepted January 11, 2013.

2010 Mathematics Subject Classification: Primary 53A45, 53B50, 53C25.

Key words and phrases: Einstein's connection, Einstein's field equations, Einstein's vector, the third class, torsion tensor, torsion vector, unified field theory.

in the sequel, Greek indices run over the range $\{1, 2, \dots, n\}$ and follow the summation convention. The algebraic structure on X_n is imposed by a basic real non-symmetric tensor $g_{\lambda\mu}$, which may be split into its symmetric part $h_{\lambda\mu}$ and skew-symmetric part $k_{\lambda\mu}$:

$$(2.1) \quad g_{\lambda\mu} = h_{\lambda\mu} + k_{\lambda\mu},$$

where we assume that

$$(2.2) \quad (a) G = \det((g_{\lambda\mu})) \neq 0, \quad (b) H = \det((h_{\lambda\mu})) \neq 0.$$

Since $\det((h_{\lambda\mu})) \neq 0$, we may define a unique tensor $h^{\lambda\nu} (= h^{\nu\lambda})$ by

$$(2.3) \quad h_{\lambda\mu} h^{\lambda\nu} = \delta_{\mu}^{\nu}.$$

We use the tensors $h^{\lambda\nu}$ and $h_{\lambda\mu}$ as tensors for raising and/or lowering indices for all tensors defined on X_n in the usual manner. Then we may define new tensors by

$$(2.4) \quad (a) k^{\alpha}_{\mu} = k_{\lambda\mu} h^{\lambda\alpha}, \quad (b) k_{\lambda}^{\alpha} = k_{\lambda\mu} h^{\mu\alpha}, \quad (c) k^{\alpha\beta} = k_{\lambda\mu} h^{\lambda\alpha} h^{\mu\beta}.$$

The manifold X_n is assumed to be connected by a general real connection $\Gamma_{\lambda\mu}^{\nu}$ which may also be split into its symmetric part $\Lambda_{\lambda\mu}^{\nu}$ and skew-symmetric part $S_{\lambda\mu}^{\nu}$, called the *torsion tensor* of $\Gamma_{\lambda\mu}^{\nu}$:

$$(2.5) \quad (a) \Lambda_{\lambda\mu}^{\nu} = \Gamma_{(\lambda\mu)}^{\nu} = \frac{1}{2}(\Gamma_{\lambda\mu}^{\nu} + \Gamma_{\mu\lambda}^{\nu}),$$

$$(b) S_{\lambda\mu}^{\nu} = \Gamma_{[\lambda\mu]}^{\nu} = \frac{1}{2}(\Gamma_{\lambda\mu}^{\nu} - \Gamma_{\mu\lambda}^{\nu}).$$

The *Einstein's n-dimensional unified field theory in X_n* is governed by the following set of equations:

$$(2.6) \quad \partial_{\omega} g_{\lambda\mu} - g_{\alpha\mu} \Gamma_{\lambda\omega}^{\alpha} - g_{\lambda\alpha} \Gamma_{\omega\mu}^{\alpha} = 0 \quad (\partial_{\nu} = \frac{\partial}{\partial x^{\nu}}),$$

and

$$(2.7) \quad (a) S_{\lambda} = S_{\lambda\alpha}^{\alpha} = 0, \quad (b) R_{[\lambda\mu]} = \partial_{[\lambda} P_{\mu]}, \quad (c) R_{(\lambda\mu)} = 0,$$

where P_{μ} is an arbitrary vector, called the *Einstein's vector*, and $R_{\lambda\mu}$ is the contracted curvature tensor $R_{\lambda\mu\alpha}^{\alpha}$ of the curvature tensor $R_{\lambda\mu\nu}^{\omega}$:

$$(2.8) \quad R_{\lambda\mu\nu}^{\omega} = \partial_{\mu} \Gamma_{\lambda\nu}^{\omega} - \partial_{\nu} \Gamma_{\lambda\mu}^{\omega} + \Gamma_{\lambda\nu}^{\alpha} \Gamma_{\alpha\mu}^{\omega} - \Gamma_{\lambda\mu}^{\alpha} \Gamma_{\alpha\nu}^{\omega}.$$

The equation (2.6) is called the *Einstein's equation*, and the solution $\Gamma_{\lambda\mu}^{\nu}$ of the Einstein's equation is called an *Einstein's connection*. And the vector S_{λ} , defined by (2.7)(a), is called the *torsion vector*.

3. An Einstein's connection in X_n

The following theorem was proved by Hlavatý([5]).

THEOREM 3.1. *In X_n , if the Einstein's equation (2.6) admits a solution $\Gamma_{\lambda\mu}^\nu$, then this solution must be of the form*

$$(3.1) \quad \Gamma_{\lambda\mu}^\nu = \{\lambda^\nu{}_\mu\} + 2h^{\nu\alpha}S_{\alpha(\lambda}{}^\beta k_{\mu)\beta} + S_{\lambda\mu}{}^\nu,$$

where $\{\lambda^\nu{}_\mu\}$ are the Christoffel symbols defined by $h_{\lambda\mu}$.

REMARK 3.2. In virtue of Theorem 3.1, the equation (3.1) reduces the investigation of $\Gamma_{\lambda\mu}^\nu$ to the study of its torsion tensor $S_{\lambda\mu}{}^\nu$. Hence in order to know an Einstein's connection $\Gamma_{\lambda\mu}^\nu$, it is necessary and sufficient to know its torsion tensor $S_{\lambda\mu}{}^\nu$. For this, we introduce a new torsion tensor $S_{\lambda\mu}{}^\nu$ given by

$$(3.2) \quad S_{\lambda\mu}{}^\nu = 2\delta_{[\lambda}^\nu k_{\mu]\alpha} Y^\alpha + k_{\lambda\mu} Y^\nu,$$

for some nonzero vector Y_λ .

THEOREM 3.3. *In X_n , if the connection (3.1) is a connection such that its torsion tensor is of the form (3.2) for some nonzero vector Y_λ , then the connection is given by*

$$(3.3) \quad \Gamma_{\lambda\mu}^\nu = \{\lambda^\nu{}_\mu\} + 2\delta_{[\lambda}^\nu k_{\mu]\alpha} Y^\alpha + k_{\lambda\mu} Y^\nu.$$

Proof. Since the torsion tensor of the connection (3.1) is of the form (3.2), we obtain

$$(3.4) \quad 2h^{\nu\alpha}S_{\alpha(\lambda}{}^\beta k_{\mu)\beta} = 0$$

by a straightforward computation. Substituting (3.2) and (3.4) into (3.1), we obtain (3.3). \square

THEOREM 3.4. *In X_n , the connection (3.3) is an Einstein's connection if and only if the vector Y_λ defining (3.3) satisfies the following condition*

$$(3.5) \quad \nabla_\nu k_{\lambda\mu} = 2h_{\nu[\lambda} k_{\mu]\alpha} Y^\alpha - 2k_{\nu[\lambda} Y_{\mu]},$$

where ∇_ω is the symbolic vector of the covariant derivative with respect to $\{\lambda^\nu{}_\mu\}$.

Proof. The connection (3.3) is an Einstein's connection if and only if the connection (3.3) satisfies the Einstein's equation (2.6). Substituting (2.1) and (3.3) into (2.6), and making use of $\nabla_\nu h_{\lambda\mu} = 0$, we obtain

$$(3.6) \quad \nabla_\nu k_{\lambda\mu} - 2h_{\nu[\lambda} k_{\mu]\alpha} Y^\alpha + 2k_{\nu[\lambda} Y_{\mu]} = 0$$

by a straightforward computation. Hence the connection (3.3) is an Einstein's connection if and only if the vector Y_λ defining (3.3) satisfies the condition (3.5). \square

THEOREM 3.5. *In X_n , if the connection (3.3) is an Einstein's connection, then its torsion vector satisfies the following relation :*

$$(3.7) \quad S_\lambda = \nabla_\alpha k_\lambda^\alpha$$

Proof. Contracting for (3.2) for μ and ν , we obtain

$$(3.8) \quad S_\lambda = S_{\lambda\alpha}{}^\alpha = (2-n)k_{\lambda\alpha}Y^\alpha.$$

Next, multiplying $h^{\mu\alpha}$ on both sides of (3.5), and contracting for ν and α , we obtain

$$(3.9) \quad \nabla_\alpha k_\lambda^\alpha = (2-n)k_{\lambda\alpha}Y^\alpha.$$

The results (3.8) and (3.9) imply the relation (3.7). \square

4. A solution of field equations for the third class in X_4

In this section we shall display a solution for the third class of (2.6) and (2.7) in X_4 . Assume $h_{\lambda\mu}$ to be of the form

$$(4.1) \quad ((h_{\lambda\mu})) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}.$$

Define two vectors by

$$(4.2) \quad (a) A_\lambda : (0, 0, 1, -1), \quad (b) B_\lambda : (\phi, \psi, 0, 0),$$

where $\phi = \phi(x_1, x_2, x_3, x_4)$ and $\psi = \psi(x_1, x_2, x_3, x_4)$ are nonzero real-valued functions to be determined. Now, we define a basic tensor $g_{\lambda\mu}$ in X_4 by

$$(4.3) \quad g_{\lambda\mu} = h_{\lambda\mu} + k_{\lambda\mu},$$

where $h_{\lambda\mu}$ is defined by (4.1), and $k_{\lambda\mu}$ is defined by

$$(4.4) \quad k_{\lambda\mu} = 2A_{[\lambda}B_{\mu]},$$

that is,

$$(4.5) \quad ((k_{\lambda\mu})) = \begin{pmatrix} 0 & 0 & -\phi & \phi \\ 0 & 0 & -\psi & \psi \\ \phi & \psi & 0 & 0 \\ -\phi & -\psi & 0 & 0 \end{pmatrix},$$

which is obviously of the third class :

$$(4.6) \quad (a) \text{Det}(k_{\lambda\mu}) = 0, \quad (b) K(= \frac{1}{4}k_{\alpha\beta}k^{\alpha\beta}) = 0.$$

Then all the Christoffel symbols $\{\lambda^\nu{}_\mu\}$ vanish. Hence the components of the first covariant derivatives with respect to $\{\lambda^\nu{}_\mu\}$ are ordinary derivatives, and $H_{\lambda\mu\nu}^\omega = 0$. Furthermore,

$$(4.7) \quad (a) A^\lambda(= h^{\lambda\nu}A_\nu) : (0, 0, 1, 1), \quad (b) B^\lambda(= h^{\lambda\nu}B_\nu) : (\phi, \psi, 0, 0)$$

$$(4.8) \quad (a) A_\alpha A^\alpha = A_\alpha B^\alpha = 0, \quad (b) k_{\lambda\alpha}A^\alpha = 0, \quad (c) \partial_\lambda A_\mu = 0.$$

The following theorem is immediate consequences of Theorem 3.3 and Theorem 3.4, in virtue of (2.7)(a), (3.8), and (4.3).

THEOREM 4.1. *In X_4 , for the basic tensor $g_{\lambda\mu}$ given by (4.3), the connection (3.3) is given by*

$$(4.9) \quad \Gamma_{\lambda\mu}^\nu = 2\delta_{[\lambda}^\nu k_{\mu]\alpha} Y^\alpha + k_{\lambda\mu} Y^\nu,$$

and this connection (4.9) is a solution of (2.6) and (2.7)(a) if and only if the vector Y^ν defining (4.9) satisfies the following conditions

$$(4.10) \quad (a) k_{\mu\alpha} Y^\alpha = 0, \quad (b) \partial_\nu k_{\lambda\mu} = -2k_{\nu[\lambda} Y_{\mu]}.$$

If these conditions (4.10) are satisfied, then the connection (4.9) is given by

$$(4.11) \quad \Gamma_{\lambda\mu}^\nu = k_{\lambda\mu} Y^\nu,$$

which is an Einstein's connection with zero torsion vector.

REMARK 4.2. In X_4 , since the tensor $k_{\lambda\mu} \neq 0$ is skew-symmetric, we know from elementary algebra that the rank of the matrix $((k_{\lambda\mu}))$ can be either four or two. In virtue of (4.6)(a), in our case the rank must be two. Therefore, the homogeneous equations (4.10)(a) have at least two distinct solutions $Y_1^\nu : (0, 0, 1, 1)$ and $Y_2^\nu : (\psi, -\phi, 0, 0)$. Every linear combination

$$(4.12) \quad Y^\nu = \rho Y_1^\nu + \eta Y_2^\nu : (\eta\psi, -\eta\phi, \rho, \rho)$$

with scalars ρ, η is also a solution of (4.10)(a). On the other hand, if (4.12) is a solution of the condition (4.10)(b), then, in virtue of $k_{12} = 0$ and $Y_\lambda = h_{\lambda\nu} Y^\nu : (\eta\psi, -\eta\phi, \rho, -\rho)$, we obtain

$$(4.13) \quad 0 = \partial_3 k_{12} = -2k_{3[1} Y_{2]} = -\phi(-\eta\phi) + \psi(\eta\psi) = \eta(\phi^2 + \psi^2),$$

which implies that $\eta = 0$. Therefore the solutions of the conditions (4.10) are of the form :

$$(4.14) \quad Y^\nu = \rho Y_1^\nu = \rho A^\nu,$$

for some nonzero real-valued function $\rho = \rho(x_1, x_2, x_3, x_4)$ to be determined.

THEOREM 4.3. *In X_4 , the vector $Y^\nu = \rho A^\nu$ given by (4.14) is a solution of the conditions (4.10) if and only if the vector B_λ given by (4.2)(b) satisfies the following condition*

$$(4.15) \quad \partial_\omega B_\mu = \rho A_\omega B_\mu.$$

Proof. Suppose that (4.15) is satisfied. Differentiating both sides of (4.4), and making use of (4.8)(c) and (4.15), we obtain

$$(4.16) \quad \partial_\omega k_{\lambda\mu} = A_\lambda(\rho A_\omega B_\mu) - A_\mu(\rho A_\omega B_\lambda) = -k_{\omega\lambda}(\rho A_\mu) + k_{\omega\mu}(\rho A_\lambda).$$

Hence, in virtue of Remark 4.2 and the relation (4.16), the vector $Y^\nu = \rho A^\nu$ is a solution of (4.10). Conversely, suppose that the vector $Y^\nu = \rho A^\nu$ is a solution of (4.10). Since $B_\mu = k_{3\mu}$, we obtain

$$(4.17) \quad \begin{aligned} \partial_\omega B_\mu &= \partial_\omega k_{3\mu} = -k_{\omega 3} Y_\mu + k_{\omega\mu} Y_3 \\ &= -(A_\omega B_3 - A_3 B_\omega)(\rho A_\mu) + (A_\omega B_\mu - A_\mu B_\omega)(\rho A_3) = \rho A_\omega B_\mu, \end{aligned}$$

in virtue of (4.4), (4.10)(b), and (4.2). Hence the condition (4.15) is satisfied. \square

THEOREM 4.4. *In X_4 , the condition (4.15) is satisfied if and only if the functions ρ , ϕ , and ψ , given in (4.15), satisfy the following conditions, respectively,*

$$(4.18) \quad \partial\phi/\partial x^1 = 0, \quad \partial\phi/\partial x^2 = 0, \quad \partial\phi/\partial x^3 = \rho\phi, \quad \partial\phi/\partial x^4 = -\rho\phi,$$

$$(4.19) \quad \partial\psi/\partial x^1 = 0, \quad \partial\psi/\partial x^2 = 0, \quad \partial\psi/\partial x^3 = \rho\psi, \quad \partial\psi/\partial x^4 = -\rho\psi$$

$$(4.20) \quad \partial\rho/\partial x^1 = 0, \quad \partial\rho/\partial x^2 = 0, \quad \partial\rho/\partial x^3 + \partial\rho/\partial x^4 = 0.$$

Proof. In virtue of (4.15), we obtain

$$(4.21) \quad \partial\phi/\partial x^\omega = \partial_\omega B_1 = \rho A_\omega B_1 = \rho\phi A_\omega,$$

which imply (4.18), in virtue of (4.2)(a). Similarly, we obtain (4.19). Next, differentiating both sides of (4.21), we obtain

$$(4.22) \quad \frac{\partial^2\phi}{\partial x^\nu \partial x^\omega} = \frac{\partial\rho}{\partial x^\nu} \phi A_\omega + \rho^2 \phi A_\nu A_\omega,$$

in virtue of (4.8)(c) and (4.21), and we also obtain

$$(4.23) \quad \frac{\partial^2 \phi}{\partial x^\omega \partial x^\nu} = \frac{\partial \rho}{\partial x^\omega} \phi A_\nu + \rho^2 \phi A_\omega A_\nu.$$

Hence we obtain

$$(4.24) \quad 0 = \frac{\partial^2 \phi}{\partial x^\nu \partial x^\omega} - \frac{\partial^2 \phi}{\partial x^\omega \partial x^\nu} = \left(\frac{\partial \rho}{\partial x^\nu} A_\omega - \frac{\partial \rho}{\partial x^\omega} A_\nu \right) \phi,$$

which implies that, since $\phi \neq 0$,

$$(4.25) \quad \frac{\partial \rho}{\partial x^\nu} A_\omega - \frac{\partial \rho}{\partial x^\omega} A_\nu = 0.$$

From the above condition (4.25), we obtain that if $\nu = 1$ and $\omega = 3$, then $\partial \rho / \partial x^1 = 0$, if $\nu = 2$ and $\omega = 3$, then $\partial \rho / \partial x^2 = 0$, and if $\nu = 3$ and $\omega = 4$, then $\partial \rho / \partial x^3 + \partial \rho / \partial x^4 = 0$. Hence we obtain (4.20). Obviously, the converse is true. \square

THEOREM 4.5. *In X_4 , for the basic tensor $g_{\lambda\mu}$ given by (4.3), the connection (4.11) which is a solution of (2.6) and (2.7)(a) is given by*

$$(4.26) \quad \Gamma_{\lambda\mu}^\nu = 2\rho A_{[\lambda} B_{\mu]} A^\nu,$$

where ρ satisfies the condition (4.20). And the curvature tensor $R_{\lambda\mu\alpha}^\alpha$ with respect to this connection (4.26) is given by

$$(4.27) \quad R_{\lambda\mu\nu}^\omega = 2\{(\partial_{[\mu}\rho)B_{\nu]}A_\lambda - (\partial_{[\mu}\rho)A_{\nu]}B_\lambda\}A^\omega + 2\rho^2 A_{[\mu} B_{\nu]} A_\lambda A^\omega,$$

and its contracted curvature tensor $R_{\lambda\mu}$ satisfies

$$(4.28) \quad R_{\lambda\mu} = 0.$$

Proof. Substituting (4.4) and (4.14) into (4.11), we obtain (4.26), in virtue of Remark 4.2, Theorem 4.3, and Theorem 4.4. Substituting (4.26) into (2.8), we obtain (4.27) by a straightforward computation. In the next, Contracting for (4.27) for ω and ν , we obtain

$$(4.29) \quad R_{\lambda\mu} = -2(\partial_\alpha \rho) A^\alpha A_{[\lambda} B_{\mu]}.$$

On the other hand, in virtue of (4.7)(a) and (4.20), we obtain

$$(4.30) \quad (\partial_\alpha \rho) A^\alpha = \partial \rho / \partial x^3 + \partial \rho / \partial x^4 = 0$$

Hence we obtain (4.28). \square

REMARK 4.6. The set of the functions ϕ satisfying (4.18) is not empty. For example, when $\rho = \text{constant}$, the function

$$(4.31) \quad \phi(x_1, x_2, x_3, x_4) = e^{\rho(x^3 - x^4)}$$

satisfies (4.18). Similarly, we can define the function ψ satisfying (4.19).

Conclusion. In virtue of Theorem 4.5, if X_4 is endowed with a non-symmetric tensor $g_{\lambda\mu} = h_{\lambda\mu} + k_{\lambda\mu}$ such that (4.1) and (4.5), where ϕ and ψ satisfy the conditions (4.18) and (4.19), respectively. Then a solution $\Gamma_{\lambda\mu}^\nu$ of (2.6) and (2.7)(a) is given by (4.26), where ρ satisfies the condition (4.20). In the next, since the contracted curvature tensor $R_{\lambda\mu}$ with respect to the connection (4.26) satisfies $R_{\lambda\mu} = 0$, the field equation (2.7)(c) is satisfied automatically, and the field equation (2.7)(b) is equivalent to $\partial_{[\lambda} P_{\mu]} = 0$. Since the field equation (2.7)(b) is satisfied by a vector $P_\mu = \partial_\mu P$, the vector $P_\mu = \partial_\mu P$ is an Einstein's vector.

References

- [1] A. Einstein, *The meaning of relativity*, Princeton University Press, Princeton, New Jersey, 1950.
- [2] J. W. Lee, *Field equations of SE(k)-manifold X_n* , International Journal of Theoretical Physics **30** (1991), 1343–1353.
- [3] J. W. Lee, *An Einstein's connection with zero torsion vector in even-dimensional UFT X_n* , Jour. of the Chungcheong Math. Soc. **24** (2011), no. 4, 869–881.
- [4] J. W. Lee and K. T. Chung, *A solution of Einstein's unified field equations*, Comm. Korean Math. Soc. **11** (1996), no. 4, 1047–1053.
- [5] V. Hlavatý, *Geometry of Einstein's unified field theory*, P. Noordhoff Ltd. New York, 1957.

*

Department of Mathematics
 Yonsei University
 Wonju 220-710, Republic of Korea
E-mail: jwlee806@yonsei.ac.kr