

**PERIODIC SOLUTIONS OF STOCHASTIC DELAY
DIFFERENTIAL EQUATIONS AND APPLICATIONS TO
LOGISTIC EQUATION AND NEURAL NETWORKS**

DINGSHI LI AND DAOYI XU

ABSTRACT. In this paper, we consider a class of periodic Itô stochastic delay differential equations by using the properties of periodic Markov processes, and some sufficient conditions for the existence of periodic solution of the delay equations are given. These existence theorems improve the results obtained by Itô et al. [6], Bainov et al. [1] and Xu et al. [15]. As applications, we study the existence of periodic solution of periodic stochastic logistic equation and periodic stochastic neural networks with infinite delays, respectively. The theorem for the existence of periodic solution of periodic stochastic logistic equation improve the result obtained by Jiang et al. [7].

1. Introduction

Since Itô introduced his stochastic calculus about 50 years ago, the theory of stochastic differential equations has been developed very quickly [1–15, 17]. It is now being recognized to be not only richer than the corresponding theory of differential equations without stochastic perturbation but also represent a more natural framework for mathematical modeling of many real-world phenomena. Now there also exists a well-developed qualitative theory of stochastic differential equations [6, 10, 12]. However, not so much has been developed in the direction of the periodically stochastic differential equations. Till now only a few papers have been published on this topic [1, 3, 4, 15, 17]. In papers [3, 6], the authors got the conditions for the existence of periodic solution of differential equations with random right sides. Hasminskii in [4] gave some basic results on the existence of periodic solution of stochastic differential equations without delays. But, the above results can not be used to check the existence of periodic solution of general stochastic delay differential equations. In [15], Xu et

Received March 24, 2011; Revised May 6, 2013.

2010 *Mathematics Subject Classification.* 34A37, 34D23.

Key words and phrases. stochastic differential equations, periodic solution, infinite delay, logistic equation, neural networks.

The work is supported by National Natural Science Foundation of China under Grant 11271270 and 11201320.

al. showed that stochastic differential equations with finite delay (SFDE) has a periodic solution if its solutions are uniformly bounded and point dissipativity. Motivated by the above discussions, we will generalize the existence theorem of the periodic solution for SFDE to stochastic differential equations with infinite delay (ISFDE) at phase space $BC((-\infty, 0]; R^n)$. The obtained results improve the results obtained by Itô et al. [6], Bainov et al. [1] and Xu et al. [15]. As applications, we study the existence of periodic solution of periodic stochastic logistic equation and periodic stochastic neural networks with infinite delays, respectively. The theorem for the existence of periodic solution of periodic stochastic logistic equation improve the result obtained by Jiang et al. [7]

2. Preliminaries

For convenience, we introduce several notations and recall some basic definitions.

$C[X, Y]$ denotes the space of continuous mappings from the topological space X to the topological space Y . Let $R_+ = (0, +\infty)$ and $\overline{R}_+ = [0, +\infty)$. Especially, let $BC \triangleq C([- \tau, 0], R^n)$ is the space of all bounded continuous R^n -value functions ϕ defined on $[-\tau, 0]$ with the norm $\|\phi\| = \sup_{-\tau \leq s \leq 0} |\phi(s)|$, where $|\cdot|$ is any norm in R^n and τ is a fixed number or $\tau = \infty$. When $\tau = \infty$ we mean, of course, that $BC \triangleq C((-\infty, 0], R^n)$. Let $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$ be a complete probability space with a filtration $\{\mathcal{F}_t\}_{t \geq 0}$ satisfying the usual conditions (i.e., it is right continuous and \mathcal{F}_0 contains all P -null sets). If $x(t)$ is an R^n -valued stochastic process on $t \in [-\tau, \infty)$, we let $x_t = x(t+s) : -\tau \leq s \leq 0$, which is regarded as a BC -valued stochastic process for $t \geq 0$. Denote by $BC_{\mathcal{F}_0}^b([- \tau, 0], R^n)$ the family of all bounded \mathcal{F}_0 -measurable, BC -valued random variables ϕ , satisfying $E[\|\phi\|] < \infty$, where $E[f]$ mean the mathematical expectation of f .

Definition 2.1. A stochastic process $x_t(\varpi)$ with values in Banach space BC , defined for $t \geq 0$ on a probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$ is called a Markov process if, for all $A \in \mathcal{B}$, $0 \leq v < t$,

$$P\{x_t(\varpi) \in A | \mathcal{F}_v\} = P\{x_t(\varpi) \in A | x_v(\varpi)\},$$

where \mathcal{F}_v is the σ -algebra of events generated by all events of the form $\{x_u(\varpi) \in A, u \leq v\}$ and \mathcal{B} denotes the σ -algebra of Borel sets in BC .

Definition 2.2. A stochastic process $x_t(\varpi)$ is said to be periodic with period ω if its finite dimensional distributions are periodic with period ω , i.e., for any positive integer m and any moments of time t_1, \dots, t_m , the joint distributions of the random variables $x_{t_1+k\omega}(\varpi), \dots, x_{t_m+k\omega}(\varpi)$ are independent of k ($k = \pm 1, \pm 2, \dots$).

Remark 2.1. By the definition of periodicity, if $x(t)$ is a ω -periodic stochastic process, then its mathematic expectation and variance are ω -periodic [4, p. 49].

The transition function of a Markov process, $p(v, x_v, t, A) = P(x_t \in A | x_v)$, a.s., is called periodic if $p(v, x_v, t + v, A)$ is periodic in v .

Later on we shall often denote a family of Markov processes by $x_t^{(t_0, \phi)}(\varpi)$ for all $t_0 \in R_+$ and $x_{t_0} = x(t_0 + s) = \phi(s) \in BC_{\mathcal{F}_0}^b([-\tau, 0], R^n)$.

Definition 2.3. The Markov families $x_t^{(t_0, \phi)}(\varpi)$ are said to be uniformly bounded, if for each $\alpha > 0$, $t_0 \in R_+$, there exists a positive constant $\theta = \theta(\alpha)$ which is independent of t_0 such that $E \|x_{t_0}\| \leq \alpha$ implies $E [\|x_t(t_0, x_{t_0})\|] \leq \theta$, $t \geq t_0$. In a general way, the Markov families $x_t^{(t_0, \phi)}(\varpi)$ are said to be p -uniformly bounded if $E [\|\cdot\|]$ is replaced by $E [\|\cdot\|^p]$.

Denote $U_r = \{\phi \in BC : \|\phi\| < r\}$ by $\overline{U}_r = \{\phi \in BC : \|\phi\| \geq r\}$.

Lemma 2.1. *A sufficient condition for the existence of an ω -periodic Markov process with a given ω -periodic transition function $p(v, x_v, t, A)$ is that for some $t_0, \phi, x_t^{(t_0, \phi)}(\varpi)$ are uniformly stochastically continuous and*

$$(1) \quad \lim_{r \rightarrow \infty} \liminf_{T \rightarrow \infty} \frac{1}{T} \int_{t_0}^{t_0+T} p(t_0, \phi, t, \overline{U}_r) dt = 0,$$

provided the transition function $p(v, x_v, t, A)$ satisfies the following not very restrictive assumption that

$$(2) \quad \alpha(r) = \sup_{\phi \in U_{\beta(r)}, 0 < t_0, t - t_0 < \omega} p(t_0, \phi, t, \overline{U}_r) \rightarrow 0 \quad \text{as } r \rightarrow \infty,$$

for some function $\beta(r)$ which tends to infinity as $r \rightarrow \infty$.

The proof of Lemma 2.1 is essentially the same as that of Lemma 2.3 in [15], Theorem 2.1 including Remark 2.1 in [1] and [9, Theorem 2.1, p. 491] except one chooses $\phi \in BC_{\mathcal{F}_0}^b([-\tau, 0], R^n)$. But, in fact, the conditions of Lemma 2.1 are of little use for stochastic differential equations, since the properties of transition functions of such processes are usually not expressible in terms of the coefficients of the equation. So, in the following, we will give some new useful sufficient conditions.

Lemma 2.2. *If Markov families $x_t^{(t_0, \phi)}(\varpi)$ with ω -periodic transition functions are uniformly bounded and uniformly stochastically continuous, then there is an ω -periodic Markov process.*

Proof. Using Markov inequality, Xu et al. [15, p. 1009] gave that

$$p(t_0, \phi, t, \overline{U}_r) \leq \frac{1}{rP(x_{t_0} = \phi)} E [\|x_t\|].$$

From the definition of uniform boundedness, for each $\alpha > 0$, there exists a positive constant $\theta = \theta(\alpha)$ such that $E \|\phi\| \leq \alpha$ implies $E [\|x_t(t_0, x_{t_0})\|] \leq \theta$, $t \geq t_0$.

So, we get

$$\begin{aligned} & \lim_{r \rightarrow \infty} \varliminf_{T \rightarrow \infty} \frac{1}{T} \int_{t_0}^{t_0+T} p(t_0, \phi, t, \bar{U}_r) dt \\ & \leq \lim_{r \rightarrow \infty} \frac{1}{rP(\phi)} \varliminf_{T \rightarrow \infty} \frac{1}{T} \int_{t_0}^{t_0+T} E[\|x_t\|] dt \\ & \leq \lim_{r \rightarrow \infty} \frac{\theta(\alpha)}{rP(\phi)} = 0, \end{aligned}$$

that is, (1) is true. The proof of (2) is similar to the remainder of the proof of Theorem 2.4 in [15], so we omit it. \square

Lemma 2.3 ([16]). *If*

- (A₁) $u : R_+ \rightarrow R^n$ is uniformly continuous,
- (A₂) $g : R^n \rightarrow R$ is continuous and $g(x) = 0$ if and only if $x = 0$,
- (A₃) $h : R_+ \rightarrow R_+$ satisfies $\inf_{t \geq 0} \int_t^{t+\delta} h(s) ds > 0$ for any $\delta > 0$,
- (A₄) $\lim_{t \rightarrow \infty} \int_0^t h(s) g(u(s)) ds$ exists and is finite.

Then $\lim_{t \rightarrow \infty} u(t) = 0$.

3. Periodic solution of ISFDE

In this section, we consider the following periodic ISFDE

$$(3) \quad \begin{cases} dx(t) = f(t, x_t) dt + g(t, x_t) dW(t), & t \geq t_0 \geq 0, \\ x_{t_0} = x(t_0 + s) = \phi(s), & s \in [-\tau, 0], \end{cases}$$

on the probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$. The equation (3) has a Borel measurable ω -periodic drift coefficient function $f : [t_0, \infty) \times BC \rightarrow R^n$ and a Borel measurable ω -periodic diffusion coefficient function $g : [t_0, \infty) \times BC \rightarrow R^{n \times m}$ driven by m -dimensional Brownian motion W . We assume that \mathcal{F}_t is the completion of the σ -algebra $\sigma\{W(u) : t_0 \leq u \leq t\}$ for each $t \geq t_0$. The initial condition $\phi(s) \in BC_{\mathcal{F}_0}^b([-\tau, 0], R^n)$ is independent of $W(t)$, $t \geq t_0$, and $E[\|\phi\|] < \infty$.

In this section, we always assume that system (3) has a unique global solution (see [13, 14]).

Theorem 3.1. *Assume that the solutions of periodic system (3) are p -uniformly bounded for $p > 2$ and $f(t, x_t)$ and $g(t, x_t)$ satisfy*

$$(4) \quad |f(t, x_t)|^p + |g(t, x_t)|^p \leq \varphi(\|x_t\|^p), \quad p > 2,$$

where φ is a concave non-decreasing function, then there is an ω -periodic Markov process.

Proof. As the proof of Lemma 3.2 in [15], we have that the unique solution x_t of (3) is a Markov process with its transition function $p(v, x_v, t, A) = P(x_t \in A | x_v)$. Moreover, the transition function $p(v, x_v, t, A)$ of (3) is ω -periodic since the coefficients of (3) are ω -periodic in t .

Since φ is a concave non-decreasing function, we get

$$E\varphi(\|x_t\|^p) \leq \varphi(E\|x_t\|^p).$$

From the p -boundedness of x_t and the condition (4), we can get that there exists a constant $\mu > 0$ such that

$$E|f(t, x_t)|^p + E|g(t, x_t)|^p \leq \mu, \quad p > 2.$$

By Lemma 3.4 in [15], the solutions of periodic system (3) are uniformly stochastically continuous.

In the view of above, all conditions of Lemma 2.2 are satisfied. The proof is complete. \square

Suppose (3) is the autonomous system, that is, $f(t, x_t)$ and $g(t, x_t)$ do not depend on t . We have the following corollary:

Corollary 3.1. *Assume that the solutions of autonomous system (3) are p -uniformly bounded for $p > 2$ and $f(x_t)$ and $g(x_t)$ satisfy Condition (4), then there is a stationary Markov process.*

Remark 3.1. Itô et al. [6, Theorem 2, p. 19] (Bainov et al. [1, Theorem 5.3, p. 23]) gave that there is a stationary (periodic) Markov process of autonomous (periodic) system (3) if the uniformly bounded in 4-th moment and

$$(5) \quad \begin{cases} |f(t, x_t)|^4 \leq q_1 + \int_{-\infty}^0 |x(s)|^4 dk_1(s), \\ |g(t, x_t)|^4 \leq q_2 + \int_{-\infty}^0 |x(s)|^4 dk_2(s), \end{cases}$$

where q_1, q_2 are some positive constants and $k_i(s), i = 1, 2$, are scalar non-decreasing bounded functions on $(-\infty, 0]$. Clearly, Conditions (5) is a special case of Conditions (4) by taking $p = 4$ and φ to be linear. Therefore, Corollary 3.1 and Theorem 3.1 is a generation of Th.2 in [6] and Th.5.3 in [1], respectively.

Let $V \in C^{1,2}([-\tau, +\infty) \times BC, \overline{R}_+)$ denote the family of all non-negative real-value functions $V(t, x_t)$ on $[-\tau, +\infty) \times BC$ which are continuously twice differentiable in the first term and once in the second term. If $V \in C^{1,2}([-\tau, +\infty) \times BC, \overline{R}_+)$, define an operator LV from $[-\tau, +\infty) \times BC$ to R by

$$LV(t, x_t) = V_t(t, x_t) + V_x(t, x_t) f(t, x_t) + \frac{1}{2} \text{trace} [g^T(t, x_t) V_{xx}(t, x_t) g(t, x_t)],$$

where $V_t(t, x_t) = \frac{\partial V(t, x_t)}{\partial t}$, $V_x(t, x_t) = \left(\frac{\partial V(t, x_t)}{\partial x_1}, \dots, \frac{\partial V(t, x_t)}{\partial x_n} \right)$, $V_{xx}(t, x_t) = \left(\frac{\partial^2 V(t, x_t)}{\partial x_i \partial x_j} \right)_{n \times n}$.

Theorem 3.2. *Assume that the solutions of (3) are uniformly stochastically continuous. Let $V \in C^{1,2}([-\tau, +\infty) \times BC, \overline{R}_+)$ and satisfy that there exists a constant M such that*

$$(6) \quad LV(t, x_t) \leq 0, \quad \|x_t\| \geq M,$$

$$(7) \quad \inf_{\|x_t\| \in \overline{U}_R} V(t, x_t) \rightarrow \infty \quad \text{as } R \rightarrow \infty.$$

Then Eq.(3) has a ω -periodic solution.

Proof. Without loss of generality, fix the initial value $x_{t_0} \in \overline{U}_M$ arbitrarily. We write $x_t = x_t(t_0, x_{t_0})$ and set $V_r = \inf_{x_t \in \overline{U}_r} V(t, x_t)$, where $r > M$. Let τ be the first enter time of x_t into U_M , that is

$$\tau = \inf \{t \geq t_0 : x_t \in U_M\}.$$

By Itô's formula, for any $t \geq t_0$,

$$\begin{aligned} V(\tau \wedge t, x_{\tau \wedge t}) &= V(t_0, x_{t_0}) + \int_{t_0}^{\tau \wedge t} LV(s, x_s) ds \\ &\quad + \int_{t_0}^{\tau \wedge t} V_x(s, x_s) g(s, x_s) dW(s). \end{aligned}$$

Taking the expectation on both sides and making use of the condition (6), we obtain that

$$(8) \quad EV(\tau \wedge t, x_{\tau \wedge t}) \leq V(t_0, x_{t_0}).$$

Using this and Čebyšev's inequality, we get

$$\begin{aligned} P(\|x_t\| \geq r) &\leq \frac{EV(\tau \wedge t, x_{\tau \wedge t})}{V_r} \\ &\leq \frac{V(t_0, x_{t_0})}{V_r}. \end{aligned}$$

This yields

$$\begin{aligned} p(t_0, \phi, t, \overline{U}_r) &= P(x_t \in \overline{U}_r | x_{t_0} = \phi) \\ &= \frac{P(\{\|x_t\| \geq r\} \cap (x_{t_0} = \phi))}{P(x_{t_0} = \phi)} \\ &\leq \frac{P(\{\|x_t\| \geq r\})}{P(x_{t_0} = \phi)} \rightarrow 0 \quad \text{as } r \rightarrow \infty. \end{aligned}$$

Clearly, the condition (1) of Lemma (2.1) holds.

Further, it follows from the condition (6) that $LV(t, x_t) \leq K, x_t \in BC$, where K is a sufficiently large constant. Using this inequality and Itô's formula, we easily obtain that

$$EV(t, x_t) \leq V(t_0, x_{t_0}) + K(t - t_0).$$

Together with Čebyšev's inequality, this implies

$$p(t_0, x_{t_0}, t, \overline{U}_R) \leq \frac{K(t - t_0) + V(t_0, x_{t_0})}{\inf_{\|x_t\| \in \overline{U}_R} V(t, x_t)}.$$

Thus the condition (2) of Lemma 2.1 will hold if $\beta(R)$ is chosen so that

$$\frac{\sup_{\|x_t\| \in U_{\beta(R)}} V(t, x_t)}{\inf_{\|x_t\| \in \bar{U}_R} V(t, x_t)} \rightarrow 0 \quad \text{as } R \rightarrow \infty.$$

This is possible because (7) holds. The proof is completed. □

Remark 3.2. The uniform stochastic continuity is required to guarantee the compactness of stochastic sequences defined in an infinite dimensional space BC . For the stochastic ODE defined in the finite dimensional space R^n , the uniform stochastic continuity is not required. Therefore, we have the following corollary.

Corollary 3.2. *Assume that system (3) without delays exists the global solutions. Let $V \in C^{1,2}(\bar{R}_+ \times R^n, \bar{R}_+)$ and satisfy that there exists a constant M such that*

$$(9) \quad LV(t, x) \leq 0, \quad |x| \geq M,$$

$$(10) \quad \inf_{|x| > R} V(t, x) \rightarrow \infty \quad \text{as } R \rightarrow \infty.$$

Then Eq.(3) without delays has a ω -periodic solution.

4. Periodic solution of nonautonomous stochastic logistic equation

In this section, we consider the following nonautonomous stochastic logistic equation [7]

$$(11) \quad dN(t) = N(t) [a(t) - b(t)N(t)] dt + \sigma(t) N(t) dB(t),$$

on $t \geq 0$ with initial value $N(0) = N_0 > 0$, and $a(t)$, $b(t)$ and $\sigma(t)$ are continuous functions. $B(t)$ is a Brownian motion defined on $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$.

As $N(t)$ in Eq.(11) is population size at time t , it should be nonnegative. Moreover, in order for a stochastic differential equation to have a unique global (i.e., no explosion in a finite time) solution for any given initial data, the coefficients of the equation are generally required to satisfy the linear growth condition and local Lipschitz condition. However, the coefficients of Eq.(11) do not satisfy the linear growth condition, though they are locally Lipschitz continuous, so the solution of Eq.(11) may explode at a finite time. In this section we shall show that under simple hypothesis the solution of Eq.(11) is not only positive but will also not explode at any finite time by using the global existence theorem in [13, 14].

Theorem 4.1. *Assume*

(H1) *There exist constants $\sigma_1, \sigma_2, a_1, a_2, b_1 > 0, b_2 > 0$ and continuous bounded function $h(t) \geq 0$ such that*

$$\sigma_1 h(t) \leq \sigma^2(t) \leq \sigma_2 h(t), \quad a_1 h(t) \leq a(t) \leq a_2 h(t), \quad b_1 h(t) \leq b(t) \leq b_2 h(t).$$

Then for any initial conditions $N_0 > 0$, there is a unique solution $x(t)$ to Eq.(11) on $t \geq 0$, and the solution will remain in R_+ almost surely, i.e., $x(t) \in R_+$ for any $t \geq 0$ with probability 1.

Proof. First consider the equation

$$(12) \quad du(t) = \left(a(t) - b(t) e^{u(t)} - \frac{1}{2} \sigma^2(t) \right) dt + \sigma(t) dB(t)$$

on $t \geq 0$ with initial value $u(0) = \ln N(0)$. Obviously, the coefficients of Eq.(12) satisfy the local Lipschitz condition, then there is a unique non-continuable solution $u(t)$ on $t \in [0, \tau_e)$, where τ_e is the right endpoint of the maximum existing interval of $u(t)$ (see [13, 14]). Therefore, by Itô's formula, it is easy to see $N(t) = e^{u(t)}$, is the unique positive solution to Eq.(11) with initial value $N(0) \in R_+$.

Next, we will show this non-continuable solution is global, i.e., $\tau_e = \infty$, a.s. To show this statement, let us define a $C^{1,2}$ -function $V : \overline{R}_+ \rightarrow \overline{R}_+$ by

$$V(N) = \left[\sqrt{N} - 1 - 0.5 \log(N) \right].$$

The nonnegativity of this function can be seen from

$$\sqrt{u} - 1 - 0.5 \log(u) \geq 0, \quad u > 0.$$

Note that

$$(13) \quad \lim_{N \rightarrow \infty} V(N) = \infty.$$

Moreover, we estimate operator LV . Since

$$\begin{aligned} LV(N) = & -0.5b(t)N^{1.5} + 0.5b(t)N + 0.5a(t)N^{0.5} - 0.5a(t) \\ & + 0.25\sigma^2(t) - 0.125\sigma^2(t)N^{0.5}, \end{aligned}$$

by hypothesis (H1), we obtain

$$\begin{aligned} LV(N) & \leq (-0.5b_1N^{1.5} + 0.5b_2N + 0.5a_2N^{0.5} - 0.5a_1 - 0.125\sigma_1^2N^{0.5} + 0.25\sigma_2^2) h(t). \end{aligned}$$

Since the coefficient of highest-degree term of the right side is negative and $h(t)$ is bounded, there is a constant $K > 0$ such that

$$ELV(x) \leq K.$$

It follows from Theorem 5.1 in [14] that $\tau_e = \infty$, a.s. The proof is complete. \square

Theorem 4.2. Assume that $a(t)$, $b(t)$ and $\sigma(t)$ are continuous T -periodic functions, (H1) and

$$(H2) \quad B = \int_0^T [a(s) - \frac{1}{2}\sigma^2(s)] ds > 0$$

hold. Then Eq.(11) has a positive T -periodic solution. Moreover, if

$$(H3) \quad \inf_{t \geq 0} \int_t^{t+\delta} h(s) ds > 0, \text{ for any } \delta > 0$$

hold. Then Eq.(11) has a unique positive T -periodic solution, which attracts all other positive solutions of Eq.(11).

Proof. From Theorem 4.1, we know that for any initial conditions $N_0 \in R_+$, there is a unique positive solution $N(t)$ to Eq.(11) on $t \geq 0$ under Condition (H1). Taking the transformation of the form

$$u = \ln N.$$

By Itô's formula, we have

$$(14) \quad du(t) = \left(a(t) - b(t)e^{u(t)} - \frac{1}{2}\sigma^2(t) \right) dt + \sigma(t) dB(t)$$

on $t \geq 0$ with initial value $u(0) = \ln N(0)$.

For any initial value $u(0) \in R$, we denote by $u(t) = u(t, 0, u(0))$ the solution of (14). Define

$$p(t) = - \int_0^t \left(a(s) - \frac{1}{2}\sigma^2(s) - Bh(s) \right) ds,$$

where B is defined in (H2). It is to see that $p(t)$ is T -periodic continuous functions. Set

$$v(t) = u(t) + p(t).$$

By Itô's formula, we have

$$(15) \quad dv(t) = \left(Bh(t) - b(t)e^{v(t)-p(t)} \right) + \sigma(t) dB(t).$$

For (15), we define a Lyapunov function

$$V(v) = |v|,$$

which satisfies (10).

Then we have

$$LV(v) = \operatorname{sgn} v \left\{ Bh(t) - b(t)e^{v(t)-p(t)} \right\}$$

Condition (H1) implies that

$$(16) \quad LV(v) \leq h(t)(B - b_1e^{v(t)-p(t)}) \leq 0 \quad \text{as } v \rightarrow \infty.$$

We consider the case $v \rightarrow -\infty$. For the above $B > 0$, there exists $l > 0$ such that

$$(17) \quad b_2e^{v(t)-p(t)} < B, \quad v < -l.$$

Thus,

$$(18) \quad \begin{aligned} LV(v) &= \left\{ -Bh(t) + b(t)e^{v(t)-p(t)} \right\} \\ &\leq h(t)(-B + b_2e^{v(t)-p(t)}) \leq 0. \end{aligned}$$

From (16) and (18), we get that there exists a positive constant M such that

$$(19) \quad LV \leq 0, \quad |v| \geq M.$$

Thus, the condition (9) of Corollary 3.2 is satisfied, and there is a periodic solution of (14) or (15). From $N(t) = e^{u(t)}$, Eq.(11) has a positive periodic solution.

In order to prove the uniqueness of periodic solution $N^*(t)$ of Eq.(11), we will prove the global attractivity of $N^*(t)$ by Lemma 2.3. To this end, we need that sample path of the solution $N(t)$ of Eq.(11) is uniformly continuous. This conclusion has been proved by Lemma 3.2 in [8] under $b(t) > 0 \Rightarrow EN^p(t) < \infty$. So, we only need prove that $b(t) \geq 0$ with (H1) $\Rightarrow EN^p(t) < \infty$. Define

$$V(N(t)) = e^{\int_0^t h(s)ds} N^p(t),$$

where $p > 0$. By Itô's formula, we have that

$$\begin{aligned} dV(N(t)) &= e^{\int_0^t h(s)ds} N^p(t) \left[h(t) + pa(t) + \frac{1}{2}p(p-1)\sigma^2(t) - pb(t)N(t) \right] dt \\ (20) \quad &+ pe^{\int_0^t h(s)ds} N^p(t) dB(t). \end{aligned}$$

Integrating (20) from 0 to t and taking expectation on both sides, we obtain from (H1) that

$$\begin{aligned} &e^{\int_0^t h(s)ds} EN^p(t) \\ &= N^p(0) + E \int_0^t e^{\int_0^s h(u)du} N^p(s) \left[h(s) + pa(s) + \frac{1}{2}p(p-1)\sigma^2(s) - pb(s)N(s) \right] ds \\ &\leq N^p(0) + E \int_0^t e^{\int_0^s h(u)du} N^p(s) h(s) \left[1 + pa_2 + \frac{1}{2}p|p-1|\sigma_2 - pb_1N(s) \right] ds \\ &\leq N^p(0) + K \int_0^t e^{\int_0^s h(u)du} h(s) ds \\ &= N^p(0) + K \left(e^{\int_0^t h(s)ds} - 1 \right), \end{aligned}$$

where K is the maximum of the function $N^p [1 + pa_2 + \frac{1}{2}p|p-1|\sigma_2 - pb_1N]$. Then we have

$$EN^p(t) \leq N^p(0) + K < \infty.$$

Following Lemma 3.2 in [8], this yields that almost every sample path of the solution $N(t)$ of Eq.(11) is uniformly continuous on $t \geq 0$.

In order to apply Lemma 2.3, we consider a Lyapunov function $V(t)$ defined by

$$V(t) = |\log N(t) - \log N^*(t)|, \quad t \geq 0.$$

By Itô's formula, we have that

$$d(\log N(t) - \log N^*(t)) = -b(t)(N(t) - N^*(t)) dt.$$

Thus, a direct calculation of the right differential $d^+V(t)$ of $V(t)$ along the solutions leads to

$$\begin{aligned} (21) \quad d^+V(t) &= \text{sgn}(N(t) - N^*(t)) d(N(t) - N^*(t)) \\ &= -\text{sgn}(N(t) - N^*(t)) [b(t)(N(t) - N^*(t))] dt \end{aligned}$$

$$= -b(t) |N(t) - N^*(t)| dt.$$

Integrating (21) from 0 to t , we have

$$V(t) + \int_0^t b(s) |N(s) - N^*(s)| ds \leq V(0) < \infty.$$

which leads to

$$(22) \quad \int_0^\infty b_1 h(s) |N(s) - N^*(s)| ds \leq V(0).$$

Therefore from Lemma 2.3 and (H3), we obtain

$$\lim_{t \rightarrow +\infty} |N(s) - N^*(s)| = 0 \quad \text{for almost all } \omega \in \Omega,$$

which shows that $N^*(t)$ attracts all other positive solutions of Eq.(11). This implies that Eq.(11) has a unique positive T -periodic solution. \square

Remark 4.1. In [7], the authors showed that $E[1/N(t)]$ has a unique positive T -periodic solution $E[1/N_p(t)]$ provided the condition (H2), $a(t) > 0$ and $b(t) > 0$. From Theorem 4.2, Eq.(11) has a unique positive T -periodic solution. So is $1/N_p(t)$ or $E[1/N_p(t)]$ with periodic T by Remark 2.1. But, under the condition (H2) we need not $a(t) > 0$ and admit $b(t) \geq 0$ with (H1) and (H3). The improvement is in effect since the usual periodic functions $\sin t, \cos t$ etc. are admitted. However, a benefit of the results in [7] is to get the representation of global solution.

5. Application in the stochastic neural networks with infinite delays

For convenience, we introduce several notations and recall some basic definitions. Let I denote an $n \times n$ unit matrix. For $A, B \in R^{m \times n}$ or $A, B \in R^n$, the notation $A \geq B$ ($A > B$) means that each pair of corresponding elements of A and B satisfies the inequality " \geq " (" $>$ "). Especially, $A \in R^{m \times n}$ is called a nonnegative matrix if $A \geq 0$, and $z \in R^n$ is called a positive vector if $z > 0$.

$$\wp = \left\{ \psi(t) : R_+ \rightarrow R \mid \psi(t) \text{ is continuous and } \int_0^\infty \psi(s) ds < \infty \right\}.$$

For $\varphi \in C[J \subset R, R^n]$, we define

$$[\varphi(t)]_\tau = \{[\varphi_1(t)]_\tau, \dots, [\varphi_n(t)]_\tau\}, \quad [\varphi_i(t)]_\tau = \sup_{-\tau \leq \theta \leq 0} \{\varphi_i(t + \theta)\}.$$

In this section, the periodic stochastic neural networks with infinite delays is described by the following model

$$(23) \quad \begin{cases} dx_i(t) = \left[-a_i(t)x_i(t) + \sum_{j=1}^n a_{ij}(t)f_j(x_j(t)) + \sum_{j=1}^n b_{ij}(t) \int_{-\infty}^t p_{ij}(t-s)g_j(x_j(s))ds + I_i(t) \right] dt \\ \quad + \sum_{j=1}^n \sigma_{ij}(t, x_i(t)) dw_j(t), \quad t \geq t_0 \geq 0, \\ x_i(t_0 + s) = \phi_i(s), \quad s \in (-\infty, 0], \quad i = 1, 2, \dots, n, \end{cases}$$

where n corresponds to the number of units in a neural network; $x_i(t)$ corresponds to the state of the i th unit at time t ; $f_j(x_j(t))$ and $g_j(x_j(t))$ denote the activation functions of the j th unit at time t ; $a_i(t) \geq 0$ represents the rate with which the i th unit will reset its potential to the resting state in isolation when disconnected from the network and external inputs; $(a_{ij}(t))_{n \times n}$ and $(b_{ij}(t))_{n \times n}$ are connection matrices; the delay kernel $p_{ij}(t) \in \wp$; $I_i(t)$ is the external bias on the i th unit; We assume that functions $a_i(t), a_{ij}(t), b_{ij}(t)$ and $I_i(t)$ are periodic continuous functions with periodic $\omega > 0$ for $t \geq t_0, i, j = 1, \dots, n$; $\sigma(\cdot, \cdot) = (\sigma_1(\cdot, \cdot), \dots, \sigma_n(\cdot, \cdot)) : [t_0, \infty) \times R^n \rightarrow R^{n \times n}$ is the ω -periodic diffusion coefficient matrix; $w(t) = (w_1(t), \dots, w_n(t))^T$ is an n -dimensional Brownian motion defined on $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$. The initial condition $\phi(s) \in BC_{\mathcal{F}_0}^b((-\infty, 0], R^n)$.

For convenience, we will introduce the following assumptions.

(A₁) All f_j and g_j satisfy the Lipschitz condition, that is, there exist $l_j > 0$ and $k_j > 0$ such that

$$|f_j(u) - f_j(v)| \leq l_j |u - v|, \quad |g_j(u) - g_j(v)| \leq k_j |u - v|, \quad \forall u, v \in R, \quad j = 1, \dots, n.$$

(A₂) There exist constants $c_i \geq 0, d_i \geq 0$ such that for $u_1, u_2 \in R, i = 1, \dots, n$,

$$\left| (\sigma_i(t, u_1) - \sigma_i(t, u_2)) (\sigma_i(t, u_1) - \sigma_i(t, u_2))^T \right| \leq c_i h(t) |u_1 - u_2|^2.$$

where $h(t) \geq 0$ and $\sup_{0 \leq t < \infty} h(t) < \infty$.

(A₃) There exist constants $a_i, a_{ij} \geq 0, b_{ij} \geq 0, I_i > 0$ such that

$$\begin{aligned} a_i(t) &\geq a_i h(t), \quad |a_{ij}(t)| \leq a_{ij} h(t), \quad |b_{ij}(t)| \leq b_{ij} h(t), \\ |I_i(t)| &\leq I_i h(t), \quad \forall i, j = 1, 2, \dots, n. \end{aligned}$$

(A₄) There exists an integral $p > 2$ such that $S = -(P + Q)$ is an M -matrix, where $P = (p_{ij})_{n \times n}, Q = (q_{ij})_{n \times n}$,

$$\begin{aligned} p_{ii} &= -pa_i + \sum_{j=1}^n a_{ij}l_j(p-1) + \sum_{j=1}^n b_{ij}k_j(p-1) \int_0^\infty |p_{ij}(s)| ds + (p-1) \\ &\quad + \frac{1}{2}c_i p(p-1) + a_{ii}l_i, \\ p_{ij} &= a_{ij}l_j, \quad i \neq j, \quad q_{ij} = b_{ij}k_j \int_0^\infty |p_{ij}(s)| ds. \end{aligned}$$

For the sake of simplicity, we use $x(t)$ to denote the solutions $x(t_0, \phi)(t)$ of (23). In order to obtain the boundedness of the stochastic system (23), we first show the following theorem.

Theorem 5.1. *Let $P = (p_{ij})_{n \times n}$ with $p_{ij} \geq 0 (i \neq j)$ and $Q(t) = (q_{ij}(t))_{n \times n}$ with $0 \leq q_{ij}(t) \in \wp$. Denote $Q = (q_{ij})_{n \times n} \triangleq (\int_0^\infty q_{ij}(t) dt)_{n \times n}$ and $I = (I_1, \dots, I_n) > 0$. Let $D = -(P + Q)$ be an M -matrix, and $u(t) = (u_1(t), u_2(t), \dots, u_n(t))^T$ a nonnegative continuous vector function satisfies the following differential inequality*

$$(24) \quad D^+u(t) \leq h(t)[Pu(t) + \int_0^\infty Q(s)u(t-s) ds + I], \quad t \geq t_0,$$

with the initial condition $u(t_0 + s) = \phi(s) \in BC^+, s \in (-\infty, 0]$, where $BC^+ = \{\phi \in BC \mid \phi \geq 0\}$ and $h(t) \geq 0$, then all solutions of the inequality (24) are uniformly bounded.

Proof. Since $D = -(P + Q)$ is an M -matrix, from the properties of M -matrix [16, Lemma 2.1], $D^{-1}I > 0$. For any given initial function $\phi \in BC^+$, there is a $d \geq 0$ such that $\phi \leq -d(P + Q)^{-1}I$. We will prove that

$$(25) \quad u(t) \leq -d(P + Q)^{-1}I, \quad t \geq t_0,$$

that is, all solutions of the inequality (24) are uniformly bounded.

We set that $-(P + Q)^{-1}I = N$. So, we have $(P + Q)N + I = 0$, or,

$$(26) \quad \sum_{j=1}^n p_{ij}N_j + \sum_{j=1}^n q_{ij}N_j + I_i = 0, \quad i = 1, 2, \dots, n.$$

If (25) is not true, then there must exist a positive constant $t_1 > t_0$ and some integer m such that

$$(27) \quad u_m(t_1) = dN_m, \quad D^+u_m(t_1) > 0,$$

$$(28) \quad u_i(t) \leq dN_i, \quad t \in (-\infty, t_1], \quad i = 1, 2, \dots, n.$$

By using (24), (25), (26), (28) and $p_{ij} \geq 0 (i \neq j)$, $q_{ij} \geq 0$, noting $d \geq 1$, we have

$$\begin{aligned} D^+u_m(t_1) &\leq \left[\sum_{j=1}^n p_{mj}u_j(t_1) + \sum_{j=1}^n q_{mj}[u_j(t_1)]_\infty + I_i \right] h(t) \\ &\leq \left[\sum_{j=1}^n (p_{mj} + q_{mj}) dN_j + I_i \right] h(t) \\ &= (1 - d) I_m h(t) \leq 0, \end{aligned}$$

which contradicts the inequality in (25). That implies that $u(t)$ is uniformly bounded. □

Theorem 5.2. *Suppose that (A_1) - (A_4) hold, then the system (23) must have a ω -periodic Markov process.*

Proof. Let $V_i(x(t)) = |x_i(t)|^p$, $i = 1, \dots, n$, where $x(t) = (x_1(t), \dots, x_n(t))^T$ is the solution of Eq.(23). By the conditions (A_1) - (A_4) , we obtain

$$\begin{aligned}
LV_i(x(t)) &\leq p|x_i(t)|^{p-1} \left\{ -a_i(t)|x_i(t)| + \sum_{j=1}^n |a_{ij}(t)| |f_j(x_j(t))| \right. \\
&\quad \left. + \sum_{j=1}^n |b_{ij}(t)| \int_{-\infty}^t |p_{ij}(t-s)| |g_j(x_j(s))| ds + |I_i(t)| \right\} \\
&\quad + \frac{1}{2}p(p-1)|x_i(t)|^{p-2} \sigma_i^T(t, x_i(t)) \sigma_i(t, x_i(t)) \\
&\leq \left(-pa_i|x_i(t)|^p + \sum_{j=1}^n a_{ij}l_j p|x_i(t)|^{p-1}|x_j(t)| \right. \\
&\quad \left. + \sum_{j=1}^n b_{ij}k_j p|x_i(t)|^{p-1} \int_0^\infty |p_{ij}(s)| |x_j(t-s)| ds + I_i p|x_i(t)|^{p-1} \right. \\
&\quad \left. + \frac{1}{2}p(p-1)|x_i(t)|^{p-2} c_i x_i^2 \right) h(t) \\
&\leq \left(-pa_i|x_i(t)|^p + \sum_{j=1}^n a_{ij}l_j (|x_j(t)|^p + (p-1)|x_i(t)|^p) \right. \\
&\quad \left. + \sum_{j=1}^n b_{ij}k_j \int_0^\infty |p_{ij}(s)| (|x_j(t-s)|^p + (p-1)|x_i(t)|^p) ds \right. \\
&\quad \left. + (p-1)|x_i(t)|^p + I_i^p + \frac{1}{2}c_i p(p-1)|x_i(t)|^p \right) h(t) \\
&= \left(\left\{ -pa_i + \sum_{j=1}^n a_{ij}l_j(p-1) + \sum_{j=1}^n b_{ij}k_j(p-1) \int_0^\infty |p_{ij}(s)| ds \right. \right. \\
&\quad \left. \left. + (p-1) + \frac{1}{2}c_i p(p-1) \right\} |x_i(t)|^p + \sum_{j=1}^n a_{ij}l_j |x_j(t)|^p \right. \\
&\quad \left. + \sum_{j=1}^n b_{ij}k_j \int_0^\infty |p_{ij}(s)| |x_j(t-s)|^p ds + I_i^p \right) h(t) \\
(29) \quad &= \left(\sum_{j=1}^n p_{ij} V_j(x) + \sum_{j=1}^n \int_0^\infty q_{ij}(s) V_j(x(t-s)) ds + I_i^p \right) h(t), \quad t \geq t_0.
\end{aligned}$$

By Itô's formula, we obtain

$$V_i(x(t)) = V_i(x(t_0)) + \int_{t_0}^t LV_i(x(s))ds + \int_{t_0}^t \frac{\partial V_i(x(s))}{\partial x} \sigma_{ij}(s, x(s)) dw(s).$$

Then we have

$$(30) \quad EV_i(x(t)) = V_i(x(t_0)) + \int_{t_0}^t ELV_i(x(s))ds,$$

and for small enough $\Delta t > 0$,

$$(31) \quad EV_i(x(t + \Delta t)) = V_i(x(t_0)) + \int_{t_0}^{t+\Delta t} ELV_i(x(s))ds,$$

Thus, from (29), (30) and (31), we have

$$(32) \quad \begin{aligned} & EV_i(x(t + \Delta t)) - EV_i(x(t)) \\ &= \int_t^{t+\Delta t} ELV_i(x(s))ds \\ &\leq \int_t^{t+\Delta t} \left(\sum_{j=1}^n p_{ij} EV_j(x(s)) + \sum_{j=1}^n \int_0^\infty q_{ij}(u) EV_j(x(s-u)) du + I_i^p \right) h(t)ds, \end{aligned}$$

Then from (32), we obtain that

$$D^+ EV_i(x(t)) \leq \left(\sum_{j=1}^n p_{ij} EV_j(x(t)) + \sum_{j=1}^n \int_0^\infty q_{ij}(s) EV_j(x(t-s)) ds + I_i^p \right) h(t).$$

Since $-(P + Q)$ is an M -matrix, then from Theorem 5.1, the solutions of (23) are uniformly bounded. By simply computing, Condition (4) is satisfied by Condition (A_1) - (A_3) . From Theorem 3.1, then there must exist an ω -periodic Markov process. The proof is complete. □

6. Conclusion

In this paper, we discuss a class of periodic Itô stochastic delay differential equations by using the properties of periodic Markov processes, and some sufficient conditions for the existence of periodic solution of the delay equations are given. As applications, we study the existence of periodic solution of periodic stochastic logistic equation and periodic stochastic neural networks with infinite delays, respectively. In our following papers, we will apply the existence theorem of periodic solution obtained in this paper to study the existence of periodic solution of periodic stochastic population system.

Acknowledgments. The authors would like to thank the referees for their detailed comments and valuable suggestions which considerably improved the presentation of the paper.

References

- [1] D. D. Bainov and V. B. Kolmanovskii, *Periodic solution of stochastic functional differential equations*, Math. J. Toyama Univ. **14** (1991), 1–39.
- [2] L. E. Bertram and P. E. Sarachik, *Stability of Circuits with randomly time-varying parameters*, IRE. Trans. Circuit Theory, CT-6, Special supplement, 1959, 260–270.
- [3] R. Z. Has'minskii, *On the dissipativity of random processes defined by differential equations*, Problemy Peredači Informacii **1** (1965), no. 1, 88–104.
- [4] ———, *Stochastic Stability of Differential Equations*, Sijthoff and Noordhoff, Maryland, 1980.
- [5] K. Itô, *On stochastic differential equations*, Mem. Amer. Math. Soc. (1951), no. 4, 51 pp.
- [6] K. Itô and M. Nisio, *On stationary solutions of a stochastic differential equation*, J. Math. Kyoto Univ. **4** (1964), no. 1, 1–75.
- [7] D. Q. Jiang and N. Z. Shi, *A note on nonautonomous logistic equation with random perturbation*, J. Math. Anal. Appl. **303** (2005), no. 1, 164–172.
- [8] D. Q. Jiang, N. Z. Shi, and X. Y. Li, *Global stability and stochastic permanence of a non-autonomous logistic equation with random perturbation*, J. Math. Anal. Appl. **340** (2008), no. 1, 588–597.
- [9] V. B. Kolmanovskii and A. Myshkis, *Introduction to the Theory and Application of Functional Differential Equations*, London, 1999.
- [10] X. X. Liao and X. R. Mao, *Exponential stability and instability of stochastic neural networks*, Stochastic Anal. Appl. **14** (1996), no. 2, 165–185.
- [11] K. N. Lu and B. Schmalfuss, *Invariant manifolds for stochastic wave equations*, J. Differential Equations **236** (2007), no. 2, 460–492.
- [12] X. R. Mao, *Exponential Stability of Stochastic Differential Equations*, Monographs and Textbooks in Pure and Applied Mathematics, 182. Marcel Dekker, Inc., New York, 1994.
- [13] ———, *Stochastic Differential Equations and Applications*, Horwood, Chichester, 1997.
- [14] L. Y. Teng, L. Xiang, and D. Y. Xu, *Existence-uniqueness of the solution for neutral stochastic functional differential equations*, Rocky Mountain Journal of Mathematics (in press).
- [15] D. Y. Xu, Y. M. Huang, and Z. G. Yang, *Existence theorems for periodic Markov process and stochastic functional differential equations*, Discrete Contin. Dyn. Syst. **24** (2009), no. 3, 1005–1023.
- [16] D. Y. Xu and L. G. Xu, *New results for studying a certain class of nonlinear delay differential systems*, IEEE Trans. Automat. Control **55** (2010), no. 7, 1641–1645.
- [17] B. G. Zhang, *On the periodic solution of n -dimensional stochastic population models*, Stochastic Anal. Appl. **18** (2000), no. 2, 323–331.

DINGSHI LI
 SCHOOL OF MATHEMATICS
 SOUTHWEST JIAOTONG UNIVERSITY
 CHENGDU, 610031, P. R. CHINA
E-mail address: lidingshi2006@163.com

DAOYI XU
YANGTZE CENTER OF MATHEMATICS
SICHUAN UNIVERSITY
CHENGDU, 610064, P. R. CHINA
E-mail address: `dyxu@scu.edu.cn`