

Surgical extraction of mandibular third molar in pterygomandibular space: a case report

Young-Kyu Lee¹, Sung-Soo Park^{1,2}, Hoon Myoung^{1,3}

Department of Oral and Maxillofacial Surgery, ¹School of Dentistry, Seoul National University, Seoul, ²Korean Armed Forces Capital Hospital, Seongnam, ³Dental Research Institute, Seoul National University, Seoul, Korea

Abstract (J Korean Assoc Oral Maxillofac Surg 2013;39:242-245)

Impacted mandibular third molars are located between the second mandibular molar and mandibular ramus. However, ectopic mandibular third molars with heterotopic positions are reported in the subcondylar or pterygomandibular space. The usual cause of malposition is a cyst or tumor, and malposition without a pathology is rare. This case report described an impacted mandibular third molar in the pterygomandibular space without any associated pathology.

Key words: Third molar, Pterygomandibular space

[paper submitted 2013. 8. 9 / revised 2013. 9. 9 / accepted 2013. 9. 11]

I. Introduction

The surgical extraction of an impacted mandibular third molar is one of the most common procedures performed by oral and maxillofacial surgeons. Most mandibular third molars are impacted in the mandibular ramus area near the second molar, and the level of difficulty of extraction is classified according to the degree of impaction, position in the mandibular ramus, and angulation of the long axis of teeth. Usually, a third molar impacted far away from its original site is affected by a cyst or a tumor. Only a few cases of ectopic mandibular third molar in the region of pterygomandibular space without association of cystic lesion--such as odontogenic keratocyst and dentigerous cyst--have been reported^{1,2}. We report a case of mandibular third molar located in the pterygomandibular space that seems to have been displaced by neither cyst nor tumor.

II. Case Report

A 46-year-old male patient visited the Department of Oral and Maxillofacial Surgery, Seoul National University Dental Hospital, complaining of swelling and pain in the right preauricular region. He was previously told by a general dentist at a local clinic that he had a malposed tooth in the right mandible and was advised not to have the tooth extracted until symptoms appear. A panoramic radiograph showed a third molar located high in the ascending ramus of the right mandible.(Fig. 1) To identify the exact location of the tooth, computed tomography (CT) was taken, showing the third molar in the pterygomandibular space associated with a radiolucent lesion.(Fig. 2) The radiolucent lesion was evaluated as cystic lesion such as odontogenic keratocyst or dentigerous cyst or secondary inflammation accompanied by soft tissue involvement.

The surgery was performed under general anesthesia via the intraoral approach. An incision was made over the right external oblique ridge and extended from the second molar to the posterosuperior mandibular ascending ramus. Subperiosteal dissection was done superiorly, exposing the anterior border of the ramus from the retromolar area almost to the tip of the coronoid process. Medial subperiosteal dissection proceeded posteriorly, exposing the lingula and inferior alveolar neurovascular bundle up to the condyle neck. Osteotomy was

Hoon Myoung

Department of Oral and Maxillofacial Surgery, Dental Research Institute, Seoul National University, 101, Daehak-ro, Jongno-gu, Seoul 110-749, Korea
TEL: +82-2-2072-3059 FAX: +82-2-766-4948
E-mail: myoungh@snu.ac.kr

©This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2013 The Korean Association of Oral and Maxillofacial Surgeons. All rights reserved.

performed with a round bur on the medial side of the mandible. The tooth was exposed and carefully removed.(Fig. 3) The tooth was easily removed from the area between the lingula and the sigmoid notch. The postoperative panoramic radiograph showed the removal of ectopic mandibular third molar.(Fig. 4) Sharp areas were smoothed, with the site curetted and cleaned with sterile saline solution. The surgical



Fig. 1. Panoramic view of ectopic mandibular third molar in the right ascending ramus of the mandible.
Young-Kyu Lee et al: Surgical extraction of mandibular third molar in pterygomandibular space: a case report. J Korean Assoc Oral Maxillofac Surg 2013

site was primarily sutured, with 4/0 vicryl applied. The postoperative wound healing was uneventful, with no nerve damage symptom. A small follicular space enveloping the crown of the tooth was also identified, suggesting inflamed granulation tissue.(Fig. 5) A connection to the periodontal space of the mandibular second molar was detected. In view of the sclerotic change of the underlying mandible (Fig. 2) and dental caries (Fig. 3B), we assume that there had been prolonged communication with the oral cavity. This ultimately led to the infectious process.

III. Discussion

Several studies have reported ectopic mandibular third molar in the mandibular ramus^{3,4}, mandibular condyle⁵⁻⁷, and coronoid process⁸.

The true incidence and etiology of ectopic mandibular third molar remain unknown⁹. An aberrant eruption pattern has been suggested to occur when the tooth has been displaced by a lesion, usually an odontogenic cyst^{3,10,11}. Dentigerous cyst is the most common benign lesion related to




Fig. 2. Axial image (A) and coronal image (B) of computed tomography (CT). CT scans show that the mandibular third molar is located in the pterygomandibular space. The sclerotic change of the mandible around the mandibular third molar is consistent with chronic infection.

Young-Kyu Lee et al: Surgical extraction of mandibular third molar in pterygomandibular space: a case report. J Korean Assoc Oral Maxillofac Surg 2013



Fig. 3. A. Intraoperative view of the osteotomy site in the medial aspect of the ascending ramus of the mandible. B. Removed ectopic mandibular third molar whose crown was blackened.

Young-Kyu Lee et al: Surgical extraction of mandibular third molar in pterygomandibular space: a case report. J Korean Assoc Oral Maxillofac Surg 2013



Fig. 4. Postoperative panoramic radiograph showing the removal of the ectopic third molar in the pterygomandibular space.
Young-Kyu Lee et al: Surgical extraction of mandibular third molar in pterygomandibular space: a case report. J Korean Assoc Oral Maxillofac Surg 2013




Fig. 5. Histopathologic imaging (H&E, $\times 200$).
Young-Kyu Lee et al: Surgical extraction of mandibular third molar in pterygomandibular space: a case report. J Korean Assoc Oral Maxillofac Surg 2013

impacted mandibular third molar¹². Over time, the pressure exerted by the intracystic fluid on the occlusal aspect of the third molar may cause its displacement, sometimes from its original location^{3,4,13}.

In the present case, the mandibular third molar was not displaced by a cystic lesion but by an uncertain cause. The development of the tooth germ in an aberrant position or aberrant tooth germ eruption pattern may be the most likely etiology. Otherwise, primary and total dislocation of tooth base may be the cause⁸. In the process of mandibular skeletal growth, bone is typically added along the posterior ramus border and resorbed along the anterior border; mandibular condyle develops as a result of bone apposition in the posterior ramus¹⁴. In this case, the presence of dental caries implies that tooth dislocation occurred after its exposure to the oral cavity.

Keros and Susić⁸ reported the ectopic mandibular third molar in the coronoid process and assumed that the bone forming the mandibular base in childhood may shift to the region beneath the coronoid process in adulthood, with the ectopic mandibular third molar embedded. Following the normal growth pattern, the third molar crown moved upward, eventually reaching the coronoid process of the mandible in non-inverted state.

Nonetheless, the etiology of ectopic impaction and migration of tooth is still unclear. Peck¹⁵ reported that the intraosseous migration of impacted mandibular tooth is related to genetic determinants. According to Marks and Schroeder¹⁶, regional disturbance in the dental follicle might lead to local defective osteoclastic function, with an abnormal eruption pathway being formed. Sutton¹⁷ believed that an abnormally strong eruption force or a change affecting the crypt of the tooth germ might lead to erroneous eruption.

Surgical extraction is mostly performed by an intra-oral approach. Extra-oral approach is done in extremely displaced impacted tooth cases. When teeth are located near the mandibular condyle, the preauricular approach can be used. The approach has the advantage of good exposure of the surgical site but may result in complications such as extraoral scar, damage to temporomandibular joint, and facial nerve injury¹¹. The intraoral approach may avoid these problems, but access to and view of the severely displaced tooth may be limited; thus making the surgery difficult. In this case, the impacted tooth was located on the lingual side of the pterygomandibular space, and the surgery was performed using the intraoral approach. During the surgery, the inferior alveolar nerve should be protected. Moreover, excessive grinding of the coronoid process or mandibular condyle should not be done to prevent fracture.

Nowadays, most third molar extractions are performed when the patients are in their twenties, so the dislocation becomes rarer. Moreover, there may be patients with ectopic tooth without clinical symptoms, not knowing that they have a dislocated tooth. Therefore, annual panoramic radiograph taking from childhood is recommended for the early detection of ectopic third molar and its pathologic changes such as cyst formation and infection. Impacted teeth diagnosed upon routine radiographic examination--and which are not associated with any pathology--usually do not require treatment, but they should be removed to prevent cyst formation, infection, and weakening of the bone predisposing to fracture⁷. The surgical extraction of the ectopic third molar should be carefully planned and performed to minimize complications induced by surgery.

References

1. Alling CC 3rd, Catone GA. Management of impacted teeth. *J Oral Maxillofac Surg* 1993;51(1 Suppl 1):3-6.
2. Kupferman SB, Schwartz HC. Malposed teeth in the pterygomandibular space: report of 2 cases. *J Oral Maxillofac Surg* 2008;66:167-9.
3. Bux P, Lisco V. Ectopic third molar associated with a dentigerous cyst in the subcondylar region: report of case. *J Oral Maxillofac Surg* 1994;52:630-2.
4. Tümer C, Eset AE, Atabek A. Ectopic impacted mandibular third molar in the subcondylar region associated with a dentigerous cyst: a case report. *Quintessence Int* 2002;33:231-3.
5. Medici A, Raho MT, Anghinoni M. Ectopic third molar in the condylar process: case report. *Acta Biomed Ateneo Parmense* 2001;72:115-8.
6. Wassouf A, Eyrych G, Lebeda R, Grätz KW. Surgical removal of a dislocated lower third molar from the condyle region: case report. *Schweiz Monatsschr Zahnmed* 2003;113:416-20.
7. Gadre KS, Waknis P. Intra-oral removal of ectopic third molar in the mandibular condyle. *Int J Oral Maxillofac Surg* 2010;39:294-6.
8. Keros J, Susić M. Heterotopia of the mandibular third molar: a case report. *Quintessence Int* 1997;28:753-4.
9. Bortoluzzi MC, Manfro R. Treatment for ectopic third molar in the subcondylar region planned with cone beam computed tomography: a case report. *J Oral Maxillofac Surg* 2010;68:870-2.
10. Salmerón JI, del Amo A, Plasencia J, Pujol R, Vila CN. Ectopic third molar in condylar region. *Int J Oral Maxillofac Surg* 2008;37:398-400.
11. Wang CC, Kok SH, Hou LT, Yang PJ, Lee JJ, Cheng SJ, et al. Ectopic mandibular third molar in the ramus region: report of a case and literature review. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;105:155-61.
12. Suarez-Cunheiro MM, Schoen R, Schramm A, Gellrich NC, Schmelzeisen R. Endoscopic approach to removal of an ectopic mandibular third molar. *Br J Oral Maxillofac Surg* 2003;41:340-2.
13. Szerlip L. Displaced third molar with dentigerous cyst--an unusual case. *J Oral Surg* 1978;36:551-2.
14. Buschang PH, Gandini Júnior LG. Mandibular skeletal growth and modelling between 10 and 15 years of age. *Eur J Orthod* 2002;24:69-79.
15. Peck S. On the phenomenon of intraosseous migration of nonerupting teeth. *Am J Orthod Dentofacial Orthop* 1998;113:515-7.
16. Marks SC Jr, Schroeder HE. Tooth eruption: theories and facts. *Anat Rec* 1996;245:374-93.
17. Sutton PR. Migration and eruption of non-erupted teeth: a suggested mechanism. *Aust Dent J* 1969;14:269-70.