

## ON THE STABILITY OF A GENERAL ADDITIVE FUNCTIONAL INEQUALITY IN BANACH SPACES

SANG-CHO CHUNG\*

ABSTRACT. In this paper, we prove the generalized Hyers-Ulam stability of the additive functional inequality

$$\|f(2x_1) + f(2x_2) + \cdots + f(2x_n)\| \leq \|tf(x_1 + x_2 + \cdots + x_n)\|$$

in Banach spaces where a positive integer  $n \geq 3$  and a real number  $t$  such that  $2 \leq t < n$ .

### 1. Introduction

In 1940, S. M. Ulam [4] suggested the stability problem of functional equations concerning the stability of group homomorphisms.

In the next year, D. H. Hyers [1] gave a first (partial) affirmative answer to the question of Ulam for Banach spaces as follows: *If  $\delta > 0$  and if  $f : \mathcal{X} \rightarrow \mathcal{Y}$  is a mapping between Banach spaces  $\mathcal{X}$  and  $\mathcal{Y}$  satisfying*

$$\|f(x + y) - f(x) - f(y)\| \leq \delta$$

*for all  $x, y \in \mathcal{X}$ , then there is a unique additive mapping  $A : \mathcal{X} \rightarrow \mathcal{Y}$  such that  $\|f(x) - A(x)\| \leq \delta$  for all  $x, y \in \mathcal{X}$ .*

This type is called the Hyers-Ulam stability.

Throughout this paper, let  $\mathcal{X}$  be a normed linear space and  $\mathcal{Y}$  a Banach space. Let  $f : \mathcal{X} \rightarrow \mathcal{Y}$  be a mapping. In 2007, C. Park, Y. S. Cho and M. H. Han [3] proved the generalized Hyers-Ulam stability of the additive functional inequality

$$\|f(x) + f(y) + f(z)\| \leq \|f(x + y + z)\|$$

in Banach spaces. In 2011, J. R. Lee, C. Park and D. Y. Shin [2] studied the generalized Hyers-Ulam stability of the additive functional inequality

$$\|f(2x) + f(2y) + 2f(z)\| \leq \|2f(x + y + z)\|$$

---

Received September 31, 2013; Accepted October 30, 2013.

2010 Mathematics Subject Classification: Primary 39B82; Secondary 46B06.

Key words and phrases: additive functional inequality, Hyers-Ulam stability, Banach space.

in Banach spaces.

In this paper, we give some generalized Hyers-Ulam stability of the additive functional inequality

$$\|f(2x_1) + f(2x_2) + \cdots + f(2x_n)\| \leq \|tf(x_1 + x_2 + \cdots + x_n)\|$$

in Banach spaces where  $3 \leq n$  and  $2 \leq t < n$  ( $n \in \mathbb{Z}$  and  $t \in \mathbb{R}$ ).

### 2. Hyers-Ulam stability in Banach spaces

To obtain our main result, we need the following lemma.

LEMMA 2.1. *Let  $f : \mathcal{X} \rightarrow \mathcal{Y}$  be a mapping and let  $3 \leq n$  and  $2 \leq t < n$  where  $n$  is an integer and  $t$  is a real number. Then  $f$  is additive if and only if it satisfies*

$$(2.1) \quad \|f(2x_1) + f(2x_2) + \cdots + f(2x_n)\| \leq \|tf(x_1 + x_2 + \cdots + x_n)\|$$

for all  $x_1, x_2, \dots, x_n \in \mathcal{X}$ .

*Proof.* If  $f$  is additive, then clearly

$$\|f(2x_1) + \cdots + f(2x_n)\| = \|2f(x_1 + \cdots + x_n)\| \leq \|tf(x_1 + \cdots + x_n)\|$$

for all  $x_i \in \mathcal{X}$ .

Conversely, assume that  $f$  satisfies (2.1). Letting  $x_i = 0$  in (2.1), we gain  $\|nf(0)\| \leq \|tf(0)\|$  and so  $f(0) = 0$  by the assumption. Putting  $x_i = 0$  for all  $i = 3, \dots, n$ , and replacing  $x_1, x_2$  by  $x, -x$  in (2.1), we get  $\|f(-x) + f(x)\| \leq \|tf(0)\| = 0$  and so  $f(-x) = -f(x)$  for all  $x \in \mathcal{X}$ . Setting  $x_1 = \frac{x+y}{2}, x_2 = \frac{-x}{2}, x_3 = \frac{-y}{2}, x_i = 0 (4 \leq i \leq n)$  in (2.1), we have

$$\|f(x+y) + f(-x) + f(-y)\| \leq \|tf(0)\| = 0$$

for all  $x, y \in \mathcal{X}$ . Thus we obtain  $f(x+y) = f(x) + f(y)$  for all  $x, y \in \mathcal{X}$ . □

THEOREM 2.2. *Let  $f : \mathcal{X} \rightarrow \mathcal{Y}$  be a mapping with  $f(0) = 0$  and let  $3 \leq n$  and  $2 \leq t < n$ . If there is a function  $\varphi : \mathcal{X}^n \rightarrow [0, \infty)$  satisfying*

$$(2.2) \quad \|f(2x_1) + \cdots + f(2x_n)\| \leq \|tf(x_1 + \cdots + x_n)\| + \varphi(x_1, \dots, x_n)$$

and

$$(2.3)$$

$$\tilde{\varphi}(x_1, \dots, x_n) := \sum_{j=0}^{\infty} \frac{1}{2^j} \varphi((-2)^j x_1, (-2)^j x_2, (-2)^j x_3, x_4, \dots, x_n) < \infty$$

for all  $x_1, \dots, x_n \in \mathcal{X}$ , then there exists a unique additive mapping  $A : \mathcal{X} \rightarrow \mathcal{Y}$  such that

$$(2.4) \quad \|f(x) - A(x)\| \leq \frac{1}{2} \tilde{\varphi} \left( x, -\frac{x}{2}, -\frac{x}{2}, 0, \dots, 0 \right)$$

for all  $x \in \mathcal{X}$ .

*Proof.* Replacing  $x_1, x_2, x_3, x_i (4 \leq i)$  by  $(-2)^{n+1} \frac{x}{2}, (-2)^n \frac{x}{2}, (-2)^n \frac{x}{2}, 0$ , respectively, and dividing by  $2^{n+1}$  in (2.2), since  $f(0) = 0$ , we get

$$\begin{aligned} & \left\| \frac{f((-2)^{n+1}x)}{(-2)^{n+1}} - \frac{f((-2)^n x)}{(-2)^n} \right\| \\ & \leq \frac{1}{2^{n+1}} \varphi \left( (-2)^{n+1} \frac{x}{2}, (-2)^n \frac{x}{2}, (-2)^n \frac{x}{2}, 0, \dots, 0 \right) \end{aligned}$$

for all  $x \in \mathcal{X}$  and all nonnegative integers  $n$ . From the above inequality, we have

$$(2.5) \quad \begin{aligned} & \left\| \frac{f((-2)^n x)}{(-2)^n} - \frac{f((-2)^m x)}{(-2)^m} \right\| \\ & \leq \sum_{j=m}^{n-1} \left\| \frac{f((-2)^{j+1} x)}{(-2)^{j+1}} - \frac{f((-2)^j x)}{(-2)^j} \right\| \\ & \leq \sum_{j=m}^{n-1} \frac{1}{2^{j+1}} \varphi \left( (-2)^j x, (-2)^{j-1} x, (-2)^{j-1} x, 0, \dots, 0 \right) \end{aligned}$$

for all  $x \in \mathcal{X}$  and all nonnegative integers  $m, n$  with  $m < n$ . By the condition (2.3), the sequence  $\left\{ \frac{f((-2)^n x)}{(-2)^n} \right\}$  is a Cauchy sequence for all  $x \in \mathcal{X}$ . Since  $\mathcal{Y}$  is complete, the sequence  $\left\{ \frac{f((-2)^n x)}{(-2)^n} \right\}$  converges for all  $x \in \mathcal{X}$ . So we can define a mapping  $A : \mathcal{X} \rightarrow \mathcal{Y}$  by

$$A(x) := \lim_{n \rightarrow \infty} \frac{f((-2)^n x)}{(-2)^n}$$

for all  $x \in \mathcal{X}$ .

In order to prove that  $A$  satisfies (2.4), taking  $m = 0$  and letting  $n$  tend to  $\infty$  in (2.5), then we have the following inequality (2.4).

$$\begin{aligned} \left\| A(x) - f(x) \right\| & \leq \sum_{j=0}^{\infty} \frac{1}{2^{j+1}} \varphi \left( (-2)^j x, (-2)^{j-1} x, (-2)^{j-1} x, 0, \dots, 0 \right) \\ & = \frac{1}{2} \tilde{\varphi} \left( x, -\frac{x}{2}, -\frac{x}{2}, 0, \dots, 0 \right). \end{aligned}$$

Next we show that  $A$  is additive. Replacing  $x_i$  by  $(-2)^n x_i$  for all  $i = 1, 2, \dots, n$ , and dividing by  $2^n$  in (2.2), we obtain

$$\begin{aligned} & \left\| \frac{f((-2)^n 2x_1)}{(-2)^n} + \frac{f((-2)^n 2x_2)}{(-2)^n} + \dots + \frac{f((-2)^n 2x_n)}{(-2)^n} \right\| \\ & \leq \left\| t \frac{f((-2)^n(x_1 + x_2 + \dots + x_n))}{(-2)^n} \right\| \\ & \quad + \frac{1}{2^n} \varphi((-2)^n x_1, (-2)^n x_2, \dots, (-2)^n x_n) \end{aligned}$$

for all  $x_1, x_2, \dots, x_n \in \mathcal{X}$  and all nonnegative integers  $n$ . Since (2.3) gives that

$$\lim_{n \rightarrow \infty} \frac{1}{2^n} \varphi((-2)^n x_1, (-2)^n x_2, \dots, (-2)^n x_n) = 0$$

for all  $x_1, x_2, \dots, x_n \in \mathcal{X}$ , letting  $n$  tend to  $\infty$  in the above inequality, we have

$$\left\| A(2x_1) + A(2x_2) + \dots + A(2x_n) \right\| \leq \left\| tA(x_1 + x_2 + \dots + x_n) \right\|$$

so  $A$  is additive by Lemma 2.1.

Let  $A' : \mathcal{X} \rightarrow \mathcal{Y}$  be another additive mapping satisfying (2.4). Since both  $A$  and  $A'$  are additive, we have, for all positive integer  $n$

$$\begin{aligned} & \|A(x) - A'(x)\| \\ & = \frac{1}{2^n} \|A((-2)^n x) - A'((-2)^n x)\| \\ & \leq \frac{1}{2^n} (\|A((-2)^n x) - f((-2)^n x)\| + \|f((-2)^n x) - A'((-2)^n x)\|) \\ & \leq \frac{1}{2^n} \tilde{\varphi}((-2)^n x, (-2)^{n-1} x, (-2)^{n-1} x, 0, \dots, 0) \\ & = \sum_{j=n}^{\infty} \frac{1}{2^j} \varphi((-2)^{j-n} x, (-2)^{j-1-n} x, (-2)^{j-1-n} x, 0, \dots, 0) \end{aligned}$$

which goes to zero as  $n \rightarrow \infty$  for all  $x \in \mathcal{X}$  by (2.3). Therefore,  $A$  is a unique additive mapping satisfying (2.4), as desired.  $\square$

**THEOREM 2.3.** *Let  $f : \mathcal{X} \rightarrow \mathcal{Y}$  be a mapping and let  $3 \leq n$  and  $2 \leq t < n$ . If there is a function  $\varphi : \mathcal{X}^n \rightarrow [0, \infty)$  satisfying*

$$(2.6) \quad \|f(2x_1) + \dots + f(2x_n)\| \leq \|tf(x_1 + \dots + x_n)\| + \varphi(x_1, \dots, x_n).$$

where

$$(2.7) \quad \begin{aligned} & \tilde{\varphi}(x_1, x_2, \dots, x_n) \\ & := \sum_{j=1}^{\infty} 2^j \varphi \left( \frac{x_1}{(-2)^j}, \frac{x_2}{(-2)^j}, \frac{x_3}{(-2)^j}, x_4, \dots, x_n \right) < \infty \end{aligned}$$

for all  $x_1, x_2, \dots, x_n \in \mathcal{X}$ , then there exists a unique additive mapping  $A : \mathcal{X} \rightarrow \mathcal{Y}$  such that

$$(2.8) \quad \|f(x) - A(x)\| \leq \frac{1}{2} \tilde{\varphi} \left( x, -\frac{x}{2}, -\frac{x}{2}, 0, \dots, 0 \right)$$

for all  $x \in \mathcal{X}$ .

*Proof.* We have  $\varphi(0, \dots, 0) = 0$  by (2.7), and so  $f(0) = 0$  by (2.6). Replacing  $x_1, x_2, x_3, x_i (4 \leq i)$  by  $\frac{x}{(-2)^n}, \frac{x}{(-2)^{n+1}}, \frac{x}{(-2)^{n+1}}, 0$ , respectively, and multiplying by  $2^{n-1}$  in (2.6), since  $f(0) = 0$ , we get

$$\begin{aligned} & \left\| (-2)^{n-1} f \left( \frac{x}{(-2)^{n-1}} \right) - (-2)^n f \left( \frac{x}{(-2)^n} \right) \right\| \\ & \leq 2^{n-1} \varphi \left( \frac{x}{(-2)^n}, \frac{x}{(-2)^{n+1}}, \frac{x}{(-2)^{n+1}}, 0, \dots, 0 \right) \end{aligned}$$

for all  $x \in \mathcal{X}$  and all nonnegative integers  $n$ . From the above inequality, we have

$$(2.9) \quad \begin{aligned} & \left\| (-2)^n f \left( \frac{x}{(-2)^n} \right) - (-2)^m f \left( \frac{x}{(-2)^m} \right) \right\| \\ & \leq \sum_{j=m+1}^n \left\| (-2)^j f \left( \frac{x}{(-2)^j} \right) - (-2)^{j-1} f \left( \frac{x}{(-2)^{j-1}} \right) \right\| \\ & \leq \sum_{j=m+1}^n 2^{j-1} \varphi \left( \frac{x}{(-2)^j}, \frac{x}{(-2)^{j+1}}, \frac{x}{(-2)^{j+1}}, 0, \dots, 0 \right) \end{aligned}$$

for all  $x \in \mathcal{X}$  and all nonnegative integers  $m, n$  with  $m < n$ . By the condition (2.7), the sequence  $\left\{ (-2)^n f \left( \frac{x}{(-2)^n} \right) \right\}$  is a Cauchy sequence for all  $x \in \mathcal{X}$ . Since  $\mathcal{Y}$  is complete, the sequence  $\left\{ (-2)^n f \left( \frac{x}{(-2)^n} \right) \right\}$  converges for all  $x \in \mathcal{X}$ . So we can define a mapping  $A : \mathcal{X} \rightarrow \mathcal{Y}$  by

$$A(x) := \lim_{n \rightarrow \infty} \left\{ (-2)^n f \left( \frac{x}{(-2)^n} \right) \right\}$$

for all  $x \in \mathcal{X}$ .

In order to prove that  $A$  satisfies (2.8), taking  $m = 0$  and letting  $n$  tend to  $\infty$  in (2.9), then we have the following inequality (2.8).

$$\begin{aligned} \left\| A(x) - f(x) \right\| &\leq \sum_{j=0}^{\infty} 2^{j-1} \varphi \left( \frac{x}{(-2)^j}, \frac{x}{(-2)^{j+1}}, \frac{x}{(-2)^{j+1}}, 0, \dots, 0 \right) \\ &= \frac{1}{2} \tilde{\varphi} \left( x, -\frac{x}{2}, -\frac{x}{2}, 0, \dots, 0 \right). \end{aligned}$$

Next we show that  $A$  is additive. Replacing  $x_i$  by  $\frac{x_i}{(-2)^n}$  for all  $i = 1, 2, \dots, n$ , and multiplying by  $2^n$  in (2.6), we obtain

$$\begin{aligned} &\left\| (-2)^n f \left( \frac{2x_1}{(-2)^n} \right) + (-2)^n f \left( \frac{2x_2}{(-2)^n} \right) + \dots + (-2)^n f \left( \frac{2x_n}{(-2)^n} \right) \right\| \\ &\leq \left\| t(-2)^n f \left( \frac{(x_1 + x_2 + \dots + x_n)}{(-2)^n} \right) \right\| \\ &\quad + 2^n \varphi \left( \frac{x_1}{(-2)^n}, \frac{x_2}{(-2)^n}, \dots, \frac{x_n}{(-2)^n} \right) \end{aligned}$$

for all  $x_1, x_2, \dots, x_n \in \mathcal{X}$  and all nonnegative integers  $n$ . Since (2.7) gives that

$$\lim_{n \rightarrow \infty} 2^n \varphi \left( \frac{x_1}{(-2)^n}, \frac{x_2}{(-2)^n}, \dots, \frac{x_n}{(-2)^n} \right) = 0$$

for all  $x_1, x_2, \dots, x_n \in \mathcal{X}$ , letting  $n$  tend to  $\infty$  in the above inequality, we have

$$\left\| A(2x_1) + A(2x_2) + \dots + A(2x_n) \right\| \leq \left\| tA(x_1 + x_2 + \dots + x_n) \right\|$$

so  $A$  is additive by Lemma 2.1.

Let  $A' : \mathcal{X} \rightarrow \mathcal{Y}$  be another additive mapping satisfying (2.8). Since both  $A$  and  $A'$  are additive, we have, for all positive integer  $n$

$$\begin{aligned} &\|A(x) - A'(x)\| \\ &= 2^n \left\| A \left( \frac{x}{(-2)^n} \right) - A' \left( \frac{x}{(-2)^n} \right) \right\| \\ &\leq 2^n \left( \left\| A \left( \frac{x}{(-2)^n} \right) - f \left( \frac{x}{(-2)^n} \right) \right\| + \left\| f \left( \frac{x}{(-2)^n} \right) - A' \left( \frac{x}{(-2)^n} \right) \right\| \right) \\ &\leq 2^n \tilde{\varphi} \left( \frac{x_1}{(-2)^n}, \frac{-x_2}{(-2)^{n+1}}, \frac{-x_3}{(-2)^{n+1}}, 0, \dots, 0 \right) \end{aligned}$$

$$= \sum_{j=n+1}^{\infty} 2^j \varphi \left( \frac{x_1}{(-2)^{j-n}}, \frac{-x_2}{(-2)^{j+1-n}}, \frac{-x_3}{(-2)^{j+1-n}}, 0, \dots, 0 \right)$$

which goes to zero as  $n \rightarrow \infty$  for all  $x \in \mathcal{X}$  by (2.7). Therefore,  $A$  is a unique additive mapping satisfying (2.8), as desired.  $\square$

### References

- [1] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. **27** (1941), 222-224.
- [2] J. R. Lee, C. Park, and D. Y. Shin, *Stability of an additive functional inequality in proper  $CQ^*$ -algebras*, Bull. Korean Math. Soc. **48** (2011), 853-871.
- [3] C. Park, Y. S. Cho, and M. H. Han, *Functional inequalities associated with Jordan-von Neumann-type additive functional equations*, J. Inequal. Appl. **2007** (2007), Article ID 41820, 13 pages.
- [4] S. M. Ulam, *A Collection of the Mathematical Problems*, Interscience Publ., New York, 1960.

\*

Department of Mathematics Education  
 Mokwon University  
 Daejeon 302-729, Republic of Korea  
*E-mail:* math888@naver.com