

Analysis of Gene Expression in Cyclooxygenase-2-Overexpressed Human Osteosarcoma Cell Lines

Jeong A. Han¹, Ji-Yeon Kim², Jong-Il Kim^{3,4,5*}

¹Department of Biochemistry and Molecular Biology, Kangwon National University School of Medicine, Chuncheon 200-701, Korea,

²Department of Internal Medicine, Seoul National University Hospital, Seoul 110-744, Korea,

³Department of Biochemistry and Molecular Biology, Seoul National University College of Medicine, Seoul 110-799, Korea,

⁴Department of Biomedical Sciences, Seoul National University Graduate School, Seoul 110-799, Korea,

⁵Genomic Medicine Institute, Medical Research Center, Seoul National University, Seoul 110-799, Korea

Osteosarcoma is the most common primary bone tumor, generally affecting young people. While the etiology of osteosarcoma has been largely unknown, recent studies have suggested that cyclooxygenase-2 (COX-2) plays a critical role in the proliferation, migration, and invasion of osteosarcoma cells. To understand the mechanism of action of COX-2 in the pathogenesis of osteosarcoma, we compared gene expression patterns between three stable COX-2-overexpressing cell lines and three control cell lines derived from U2OS human osteosarcoma cells. The data showed that 56 genes were upregulated, whereas 20 genes were downregulated, in COX-2-overexpressed cell lines, with an average fold-change > 1.5. Among the upregulated genes, *COL1A1*, *COL5A2*, *FBN1*, *HOXD10*, *RUNX2*, and *TRAPPC2* are involved in bone and skeletal system development, while *DDR2*, *RAC2*, *RUNX2*, and *TSPAN31* are involved in the positive regulation of cell proliferation. Among the downregulated genes, *HIST1H1D*, *HIST1H2AI*, *HIST1H3H*, and *HIST1H4C* are involved in nucleosome assembly and DNA packaging. These results may provide useful information to elucidate the molecular mechanism of the COX-2-mediated malignant phenotype in osteosarcoma.

Keywords: cell proliferation, cyclooxygenase 2, invasion, osteosarcoma, overexpression, migration

Introduction

Osteosarcoma is the most common primary bone tumor, representing over 56% of all bone tumors. It generally affects young people with the age of 15 to 19 and usually occurs in long bones near the metaphyseal growth plates. It has a high tendency to metastasize, and about 10%–20% of patients have macroscopic evidence of metastasis at the time of diagnosis, while 80%–90% of patients are assumed to have micrometastasis. Although it has been postulated that rapid bone growth, exposure to radiation, and genetic predispositions, such as *RB* or *p53* mutations, are thought to be risk factors for developing osteosarcoma, the etiology has not been fully understood [1-3].

Prostaglandin endoperoxide synthase 2 (PTGS2), also

called as cyclooxygenase-2 (COX-2), catalyzes the conversion of arachidonic acid to prostaglandin H₂, from which various prostanoids, including prostaglandin E₂, are produced [4].

Accumulating evidence indicates that COX-2 is involved in osteosarcoma development and progression. Several studies have reported that high levels of COX-2 expression is associated with advanced clinical stage and metastasis [5, 6], as well as with lower overall survival rates and disease-free survival rates [7-9]. In addition, COX-2 inhibition by using RNAi or antisense oligonucleotide inhibits cell proliferation and invasion in human osteosarcoma cells [10, 11]. Also, selective COX-2 inhibitors reduce not only osteosarcoma cell proliferation and invasion *in vitro* but also tumor growth and metastasis *in vivo* [12, 13]. Moreover, we have previously reported that COX-2 overexpression promotes cell

Received October 20, 2014; Revised November 9, 2014; Accepted November 10, 2014

*Corresponding author: Tel: +82-2-740-8251, Fax: +82-2-744-4534, E-mail: jongil@snu.ac.kr

Copyright © 2014 by the Korea Genome Organization

© It is identical to the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>).

proliferation, migration, and invasion in U2OS human osteosarcoma cells [14]. These studies strongly suggest that COX-2 might be a causal factor for the development and progression of osteosarcoma. However, the exact mechanisms of action of COX-2 in osteosarcoma are largely unknown.

In an attempt to figure out the mechanism of action of COX-2 in osteosarcoma, we analyzed the gene expression profiles in three COX-2-overexpressed U2OS stable cell lines and three control stable cell lines.

Methods

Establishment and maintenance of stable cell lines

Human COX-2 cDNA was subcloned into the pcDNA3 vector containing neo^r. U2OS cells were transfected with COX-2 or pcDNA3 DNA using Lipofectamine2000 (Life Technologies, Grand Island, NY, USA). Transfectants were selected in the presence of geneticin, and individual clones were maintained in Dulbecco's modified Eagle's medium, containing fetal bovine serum (10%), penicillin (100 units/mL), streptomycin (100 units/mL), and geneticin (700 μ g/mL), as reported previously [14].

RNA isolation

Total RNA was extracted from cells with Trizol (Life Technologies), purified with the addition of chloroform, and precipitated with the addition of isopropanol. The RNA concentration was determined by a spectrophotometer, and the quality of RNA was evaluated by the OD 260/280 ratio and gel electrophoresis.

Hybridization to expression arrays

The following procedures were carried out by Macrogen Co. (Seoul, Korea). First, total RNA was amplified and purified using the Ambion Illumina RNA amplification kit to yield biotinylated cRNA (Ambion, Austin, TX, USA). Briefly, 550 ng of total RNA was reverse-transcribed to cDNA using a T7 oligo(dT) primer. Second-strand cDNA was synthesized, in vitro-transcribed, and labeled with biotin-NTP.

After purification, 750 ng of labeled cRNA was hybridized to the humanHT-12 expression v.4 bead array (Illumina, San Diego, CA, USA) for 16-18 h at 58°C. The array signal was detected using Amersham fluorolink streptavidin-Cy3 (GE Healthcare Bio-Sciences, Little Chalfont, UK). Arrays were scanned with an Illumina bead array reader/confocal scanner. Array data were filtered by a detection p-value < 0.05 (similar to signal to noise). Selected gene signal values were log-transformed and normalized by the quantile method.

Statistical analysis

Basic statistical analyses were performed using Microsoft Excel. Hierarchical cluster analysis was conducted with normalized log₂-gene expression values using Cluster 3.0, and the results were visualized using Java Treeview [15, 16]. An unrooted tree was drawn with R package. Biological function analysis was performed with official gene names using DAVID (<http://david.abcc.ncifcrf.gov/>).

Results

Stable cell lines

We have previously established stable cell lines over-expressing human COX-2 in U2OS human osteosarcoma cells. To avoid clonal variations, we established three stable COX-2-overexpressing cell lines (U2OS-COX-2 #1, #2, and #3) and three control cell lines (U2OS-pcDNA3 #1, #2, and #3). High levels of COX-2 expression were observed by western blot analysis in U2OS-COX-2 cells, whereas the expression was barely detectable in U2OS-pcDNA3 cells. When we cultured the stable cell lines, we observed that cell proliferation, migration, and invasion rates were increased significantly in U2OS-COX-2 cells as compared with U2OS-pcDNA3 cells [14] (Table 1).

Gene expression

We first performed unsupervised cluster analysis comparing COX-2-overexpressing cell lines and control cell lines. Among 24,371 probe sets, we selected and analyzed 567

Table 1. Summary of stable cell line characteristics

Cell line	Expression level of COX-2	Proliferation rate	Migration rate (trans-well)	Invasion rate (Matrigel)
U2OS-COX-2 #1	+++	+++	+++	+++
U2OS-COX-2 #2	+++	+++	+++	+++
U2OS-COX-2 #3	+++	+++	+++	+++
U2OS-pcDNA3 #1	±	+	+	+
U2OS-pcDNA3 #2	±	+	+	+
U2OS-pcDNA3 #3	±	+	+	+

COX-2, cyclooxygenase-2.




Fig. 1. Segregation between the group of U2OS-COX-2 and the group of U2OS-pcDNA3. Unsupervised cluster analysis was done between six stable cell lines using 567 probe sets with a standard deviation > 0.5 . Normalized \log_2 -gene expression values were used for hierarchical clustering, and an unrooted tree was drawn by R package. COX-2, cyclooxygenase-2.

probe sets with a standard deviation > 0.5 . The group of U2OS-COX-2 (U2OS-COX-2 #1, #2, and #3) and the group of U2OS-pcDNA3 (U2OS-pcDNA3 #1, #2, and #3) were well segregated in the unsupervised cluster analysis, confirming that our experiment was validated (Fig. 1).

To identify significantly differentially expressed genes (DEGs) in COX-2-overexpressing osteosarcoma cells, we first performed the Wilcoxon rank sum test between the group of U2OS-COX-2 and the group of U2OS-pcDNA3. The 2,144 probe sets passed the criteria. Then, to identify DEGs, we selected 76 probe sets with an average fold-change > 1.5 .

Among the DEGs, 56 genes were upregulated while 20 genes were downregulated in the group of U2OS-COX-2 as compared to the group of U2OS-pcDNA3 (Tables 2 and 3). The heat map of DEGs is provided in Fig. 2. The gene encoding COX-2, *PTGS2*, was upregulated by 10.41-fold in the COX-2-overexpressing cell lines as compared to the control cell lines, confirming again that our experiment is

Table 2. Upregulated DEGs in COX-2-overexpressed cell lines

Symbol	Description	Fold change	Accession No.
<i>PTGS2</i>	Prostaglandin-endoperoxide synthase 2	10.41	NM_000963
<i>PAGE2B</i>	P antigen family, member 2B	4.61	NM_001015038
<i>COL1A1</i>	Collagen, type I, alpha 1	3.21	NM_000088
<i>FAP</i>	Fibroblast activation protein, alpha	2.76	NM_004460
<i>COL5A2</i>	Collagen, type V, alpha 2	2.35	NM_000393
<i>LINC00312</i>	Long intergenic non-protein coding RNA 312	2.34	NR_024065
<i>RAC2</i>	Ras-related C3 botulinum toxin substrate 2 (rho family, small GTP-binding protein Rac2)	2.33	NM_002872
<i>HOXD10</i>	Homeobox D10	2.32	NM_002148
<i>RCN3</i>	Reticulocalbin 3, EF-hand calcium-binding domain	2.07	NM_020650
<i>FBN1</i>	Fibrillin 1	2.02	NM_000138
<i>CREB3L1</i>	cAMP responsive element-binding protein 3-like 1	2.02	NM_052854
<i>C1orf85</i>	Chromosome 1 open reading frame 85	2.01	NM_144580
<i>TSPAN31</i>	Tetraspanin 31	2.01	NM_005981
<i>DPYSL5</i>	Dihydropyrimidinase-like 5	1.96	NM_020134
<i>PSMB9</i>	Proteasome (prosome, macropain) subunit, beta type, 9, transcript variant 1	1.89	NM_002800
<i>PSMB9</i>	Proteasome (prosome, macropain) subunit, beta type, 9, transcript variant 2	1.89	NM_148954
<i>GPAM</i>	Glycerol-3-phosphate acyltransferase, mitochondrial	1.89	NM_020918
<i>NLRP1</i>	NLR family, pyrin domain-containing 1, transcript variant 1	1.83	NM_033004
<i>LTBP1</i>	Latent transforming growth factor beta-binding protein 1, transcript variant 1	1.82	NM_206943
<i>UNC13A</i>	Unc-13 homolog A (<i>Caenorhabditis elegans</i>)	1.79	NM_001080421
<i>VAT1L</i>	Vesicle amine transport 1-like	1.78	NM_020927
<i>MEIS1</i>	Meis homeobox 1	1.77	NM_002398
<i>DDR2</i>	Discoidin domain receptor family, member 2, transcript variant 1	1.74	NM_001014796
<i>DYSF</i>	Dysferlin, limb girdle muscular dystrophy 2B (autosomal recessive)	1.74	NM_003494
<i>ACBD7</i>	Acyl-coenzyme A-binding domain-containing 7	1.69	NM_001039844
<i>FMN1</i>	Formin 1	1.68	NM_001103184
<i>SCN3A</i>	Sodium channel, voltage-gated, type III, alpha subunit, transcript variant 1	1.68	NM_006922
<i>NLRP11</i>	NLR family, pyrin domain-containing 11	1.68	NM_145007
<i>MBOAT2</i>	Membrane bound O-acyltransferase domain-containing 2	1.68	NM_138799

DEG, differentially expressed gene; COX-2, cyclooxygenase-2.

Table 2. Continued

Symbol	Description	Fold change	Accession No.
<i>MIR196A1</i>	MicroRNA 196a-1	1.67	NR_029582
<i>ASTN1</i>	Astrotactin 1, transcript variant 1	1.65	NM_004319
<i>TSHZ2</i>	Teashirt family zinc finger 2	1.64	NM_173485
<i>MFAP2</i>	Microfibrillar-associated protein 2, transcript variant 2	1.64	NM_002403
<i>SH3PXD2A</i>	SH3 and PX domains 2A	1.62	NM_014631
<i>SNORD116-22</i>	Small nucleolar RNA, C/D box 116-22	1.60	NR_003336
<i>TSHZ2</i>	Teashirt zinc finger homeobox 2	1.59	NM_001193421
<i>EHD3</i>	EH domain-containing 3	1.59	NM_014600
<i>LBH</i>	Limb bud and heart development homolog (mouse)	1.59	NM_030915
<i>PTPN20B</i>	Protein tyrosine phosphatase, non-receptor type 20B	1.59	NM_001042357
<i>C10orf11</i>	Chromosome 10 open reading frame 11	1.59	NM_032024
<i>SNORD116-2</i>	Small nucleolar RNA, C/D box 116-2 on chromosome 15	1.59	NR_003317
<i>EFEMP1</i>	EGF-containing fibulin-like extracellular matrix protein 1, transcript variant 1	1.59	NM_004105
<i>NAG</i>	Neuroblastoma-amplified protein	1.58	NM_015909
<i>HIST4H4</i>	Histone cluster 4, H4	1.58	NM_175054
<i>KIF5A</i>	Kinesin family member 5A	1.57	NM_004984
<i>TMEPAI</i>	Transmembrane, prostate androgen-induced RNA, transcript variant 1	1.56	NM_020182
<i>UACA</i>	Vveal autoantigen with coiled-coil domains and ankyrin repeats, transcript variant 1	1.55	NM_018003
<i>RASSF2</i>	Ras association (RalGDS/AF-6) domain family 2, transcript variant 1	1.54	NM_014737
<i>GPATCH1</i>	G patch domain containing 1	1.54	NM_018025
<i>LPAR1</i>	Lysophosphatidic acid receptor 1	1.54	NM_057159
<i>ARSI</i>	Arylsulfatase family, member I	1.53	NM_001012301
<i>TRAPP2</i>	Trafficking protein particle complex 2, transcript variant 1	1.53	NM_001011658
<i>CDR1</i>	Cerebellar degeneration-related protein 1, 34 kDa	1.53	NM_004065
<i>WDPCP</i>	WD repeat-containing and planar cell polarity effector	1.53	NM_015910
<i>CCDC186</i>	Coiled-coil domain-containing protein 186	1.52	NM_018017
<i>RUNX2</i>	Runt-related transcription factor 2, transcript variant 1	1.51	NM_001024630

Table 3. Downregulated DEGs in COX-2-overexpressed cell lines

Symbol	Description	Fold change	Accession No.
<i>HIST1H4C</i>	Histone cluster 1, H4c	4.63	NM_003542
<i>HIST1H3H</i>	Histone cluster 1, H3h	2.05	NM_003536
<i>HIST1H2AI</i>	Histone cluster 1, H2ai	1.94	NM_003509
<i>CABYR</i>	Calcium binding tyrosine-(Y)-phosphorylation regulated, transcript variant 1	1.91	NM_012189
<i>FLJ16171</i>	FLJ16171 protein	1.90	NR_046113
<i>AKR1C2</i>	Aldo-keto reductase family 1, member C2, transcript variant 2	1.89	NM_205845
<i>AURKC</i>	Aurora kinase C, transcript variant 1	1.77	NM_001015878
<i>HIST1H1D</i>	Histone cluster 1, H1d	1.72	NM_005320
<i>CRLF2</i>	Cytokine receptor-like factor 2, transcript variant 1	1.70	NM_001012288
<i>CTAG2</i>	Cancer/testis antigen 2, transcript variant 2	1.69	NM_020994
<i>SERPINB5</i>	Serpin peptidase inhibitor, clade B (ovalbumin), member 5	1.65	NM_002639
<i>RARB</i>	Retinoic acid receptor, beta, transcript variant 1	1.65	NM_000965
<i>FAT4</i>	FAT tumor suppressor homolog 4 (<i>Drosophila</i>)	1.64	NM_024582
<i>TNFRSF11B</i>	Tumor necrosis factor receptor superfamily, member 11b	1.57	NM_002546
<i>NOTUM</i>	Notum pectinacylesterase homolog (<i>Drosophila</i>)	1.57	NM_178493
<i>FGF2</i>	Fibroblast growth factor 2 (basic)	1.56	NM_002006
<i>STX11</i>	Syntaxin 11	1.53	NM_003764
<i>VEPH1</i>	Ventricular zone-expressed PH domain homolog 1 (zebrafish)	1.53	NM_024621
<i>SPATA31C1</i>	SPATA31 subfamily C, member 1	1.53	NM_001145124
<i>ADAM21</i>	ADAM metallopeptidase domain 21	1.52	NM_003813

DEG, differentially expressed gene; COX-2, cyclooxygenase-2.

validated (Table 2).

Functional analysis

The DEGs were concentrated in several important biological functions

The functions that were most significantly upregulated by




Fig. 2. Gene expression patterns of 76 differentially expressed genes (DEGs). Hierarchical cluster analysis was done between six stable cell lines using 76 DEGs. Normalized \log_2 -gene expression values were analyzed by Cluster 3.0, and the results were visualized using Java Treeview, with red being high and green indicating low.

COX-2 overexpression in U2OS cells were skeletal system development and morphogenesis, bone development and morphogenesis, ossification, and positive regulation of cell proliferation ($p < 0.05$) (Table 4). The genes of *COL1A1*, *COL5A2*, *FBN1*, *HOXD10*, *RUNX2*, and *TRAPPC2* are involved in skeletal system development and morphogenesis. The genes of *COL1A1*, *COL5A2*, and *RUNX2* are involved in bone development, morphogenesis and ossification. The genes of *DDR2*, *RAC2*, *RUNX2*, and *TSPAN31* are involved in the positive regulation of cell proliferation (Table 4).

The functions that were most significantly downregulated by COX-2 overexpression in U2OS cells were nucleosome assembly, chromatin assembly, and DNA packaging ($p < 0.05$). The genes of *HIST1H1D*, *HIST1H2AI*, *HIST1H3H*, and *HIST1H4C* are involved in these processes (Table 5).

Discussion

COX-2 has been reported to promote cell proliferation, migration, and invasion in osteosarcoma cells. COX-2 gene knockdown by RNAi significantly reduced the cell proliferation, migration, and invasion in SaOS2 cells [11], whereas stable expression of COX-2 increased the ability in U2OS cells [14] (Table 1). These studies demonstrated that abnormal COX-2 overexpression is an important determinant for the malignant phenotype of osteosarcoma cells. By using bioinformatics tools, we here provided useful information to understand the underlying mechanism by which COX-2 promotes the malignant phenotype in osteosarcoma cells.

According to our data, the genes encoding extracellular matrix proteins are highly upregulated by COX-2 (Tables 2 and 4). The genes *COL1A1*, *COL5A2*, and *FBN1* encode $\alpha 1$ chain of type I collagen, $\alpha 2$ chain of type V collagen, and fibrillin 1, respectively. It has been known that the type I and type V collagens are major extracellular matrix proteins in

Table 4. Biological functions of upregulated DEGs

Functional category	Genes	p-value
GO_Biological process		
Skeletal system development	<i>PTGS2</i> , <i>FBN1</i> , <i>COL1A1</i> , <i>COL5A2</i> , <i>RUNX2</i> , <i>HOXD10</i> , <i>TRAPPC2</i>	0.00009
Ossification	<i>PTGS2</i> , <i>COL1A1</i> , <i>COL5A2</i> , <i>RUNX2</i>	0.00248
Bone development	<i>PTGS2</i> , <i>COL1A1</i> , <i>COL5A2</i> , <i>RUNX2</i>	0.00300
Positive regulation of cell proliferation	<i>PTGS2</i> , <i>RAC2</i> , <i>TSPAN31</i> , <i>DDR2</i> , <i>RUNX2</i>	0.01579
Endochondral ossification	<i>COL1A1</i> , <i>RUNX2</i>	0.02803
Skeletal system morphogenesis	<i>COL1A1</i> , <i>RUNX2</i> , <i>HOXD10</i>	0.02867
Endochondral bone morphogenesis	<i>COL1A1</i> , <i>RUNX2</i>	0.03720
Bone morphogenesis	<i>COL1A1</i> , <i>RUNX2</i>	0.04629
GO_Molecular function		
Extracellular matrix structural constituent	<i>FBN1</i> , <i>COL1A1</i> , <i>COL5A2</i>	0.01891

DEG, differentially expressed gene.

Table 5. Biological functions of downregulated DEGs

Functional category	Genes	p-value
GO_Biological process		
Nucleosome assembly	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.00005
Chromatin assembly	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.00005
Protein-DNA complex assembly	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.00006
Nucleosome organization	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.00007
DNA packaging	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.00013
Chromatin assembly or disassembly	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.00017
Cellular macromolecular complex assembly	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.00242
Cellular macromolecular complex subunit organization	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.00336
Chromatin organization	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.00395
Chromosome organization	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.00791
Macromolecular complex assembly	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.01865
Macromolecular complex subunit organization	<i>HIST1H1D, HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.02219
KEGG_Pathway		
Systemic lupus erythematosus	<i>HIST1H2AI, HIST1H4C, HIST1H3H</i>	0.01236

DEG, differentially expressed gene.

bone [17]. Although there is a report that type I collagen promotes cell proliferation and migration in gastric cancer cells [18], the causal relationship between the overexpression of those genes and osteosarcoma has not been investigated well. Therefore, further studies are necessary to elucidate whether overexpression of these genes might be involved in the COX-2-mediated malignant phenotype in osteosarcoma.

Extracellular matrix proteins not only are structural components of the connective tissue in the body but also regulate cell fates through their receptors on the cell surface. DDR2 is one of the receptors for collagen, the expression of which was highly upregulated by COX-2 (Tables 2 and 4). DDR2 is a receptor tyrosine kinase, and its overexpression promotes the migration and invasion of cancer cells derived from breast, head and neck, and prostate [19-21]. Therefore, COX-2 might promote cell proliferation and motility through upregulation of DDR2. Further studies are necessary to elucidate this possibility.

RAC2, a highly upregulated gene by COX-2, is a member of the Rho family of GTPases (Tables 2 and 4). Although there is no report on Rac2 function in osteosarcoma, there is a report that Rac2 GTPase is required for B cell proliferation and survival [22]. Therefore, COX-2 might promote cell proliferation through upregulation of Rac2 function.

Among the highly upregulated genes, *RUNX2* encodes a transcription factor governing osteoblast differentiation and skeletal morphogenesis. Recent studies on Runx2 function in cancer are controversial. Runx2 overexpression reduces cell proliferation in osteosarcoma cell lines [23] but promotes cell motility or invasion in MCF-7 breast cancer cells and PC prostate cancer cells [24, 25]. Therefore, further

studies are required to elucidate the function of Runx2 in osteosarcoma and the relation between COX-2 and Runx2.

Intriguingly, the genes encoding histone proteins are highly downregulated by COX-2 (Tables 3 and 5). The genes *HIST1H1D*, *HIST1H2AI*, *HIST1H3H*, and *HIST1H4C* encode histone H1H1, histone H2A1, histone H3A1, and histone H41, respectively. As is well known, the nucleosome is the basic unit of DNA packaging, consisting of a core particle and linker DNA. A core particle comprises 147 base pairs of DNA wrapped around a histone octamer (2 copies each of histone H2A, H2B, H3, and H4). The core particles are connected by 10–90 bp of linker DNA, which could be naked or bound to histone H1 [26]. The nucleosome itself has been known to be a general gene repressor [27, 28]. In addition, RNAi-mediated knockdown of H2A.Z reduced cell proliferation in bladder cancer cells [29]. These results are compatible with our finding that COX-2 downregulates the expression of histone genes. Therefore, it is worth studying the mechanism of action of histones in osteosarcoma and the mechanism by which COX-2 downregulates histone genes.

Osteosarcoma is a highly malignant tumor, and the 5-year survival rate is only 20%–30% in patients with macroscopic evidence of metastasis at diagnosis [3]. While the etiology of osteosarcoma has been largely unknown, recent studies have suggested that COX-2 plays a critical role in osteosarcoma development and progression. In this regard, we believe that our study will provide useful information to understand the mechanism of action of COX-2 in osteosarcoma.

Acknowledgments

This study was supported by 2014 Research Grant from

Kangwon National University (No. 120141490).

References

1. He JP, Hao Y, Wang XL, Yang XJ, Shao JF, Guo FJ, *et al.* Review of the molecular pathogenesis of osteosarcoma. *Asian Pac J Cancer Prev* 2014;15:5967-5976.
2. Luetke A, Meyers PA, Lewis I, Juergens H. Osteosarcoma treatment - where do we stand? A state of the art review. *Cancer Treat Rev* 2014;40:523-532.
3. Longhi A, Errani C, De Paolis M, Mercuri M, Bacci G. Primary bone osteosarcoma in the pediatric age: state of the art. *Cancer Treat Rev* 2006;32:423-436.
4. Smith WL, Garavito RM, DeWitt DL. Prostaglandin endoperoxide H synthases (cyclooxygenases)-1 and -2. *J Biol Chem* 1996;271:33157-33160.
5. El-Badawi ZH, Muhammad EM, Noaman HH. Role of immunohistochemical cyclo-oxygenase-2 (COX-2) and osteocalcin in differentiating between osteoblastomas and osteosarcomas. *Malays J Pathol* 2012;34:15-23.
6. Jiao G, Ren T, Lu Q, Sun Y, Lou Z, Peng X, *et al.* Prognostic significance of cyclooxygenase-2 in osteosarcoma: a meta-analysis. *Tumour Biol* 2013;34:2489-2495.
7. Boulytcheva IV, Soloviev YN, Kushlinskii NE, Mahson AN. Expression of molecular markers in the tumor and survival prognosis in osteosarcoma. *Bull Exp Biol Med* 2010;150:237-242.
8. Urakawa H, Nishida Y, Naruse T, Nakashima H, Ishiguro N. Cyclooxygenase-2 overexpression predicts poor survival in patients with high-grade extremity osteosarcoma: a pilot study. *Clin Orthop Relat Res* 2009;467:2932-2938.
9. Wang Z, He M, Xiao Z, Wu H, Wu Y. Quantitative assessment of the association of COX-2 (cyclooxygenase-2) immunoeexpression with prognosis in human osteosarcoma: a meta-analysis. *PLoS One* 2013;8:e82907.
10. Wu X, Cai M, Ji F, Lou LM. The impact of COX-2 on invasion of osteosarcoma cell and its mechanism of regulation. *Cancer Cell Int* 2014;14:27.
11. Zhao Q, Wang C, Zhu J, Wang L, Dong S, Zhang G, *et al.* RNAi-mediated knockdown of cyclooxygenase2 inhibits the growth, invasion and migration of SaOS2 human osteosarcoma cells: a case control study. *J Exp Clin Cancer Res* 2011;30:26.
12. Sabino MA, Ghilardi JR, Jongen JL, Keyser CP, Luger NM, Mach DB, *et al.* Simultaneous reduction in cancer pain, bone destruction, and tumor growth by selective inhibition of cyclooxygenase-2. *Cancer Res* 2002;62:7343-7349.
13. Naruse T, Nishida Y, Hosono K, Ishiguro N. Meloxicam inhibits osteosarcoma growth, invasiveness and metastasis by COX-2-dependent and independent routes. *Carcinogenesis* 2006;27:584-592.
14. Lee EJ, Choi EM, Kim SR, Park JH, Kim H, Ha KS, *et al.* Cyclooxygenase-2 promotes cell proliferation, migration and invasion in U2OS human osteosarcoma cells. *Exp Mol Med* 2007;39:469-476.
15. de Hoon MJ, Imoto S, Nolan J, Miyano S. Open source clustering software. *Bioinformatics* 2004;20:1453-1454.
16. Saldanha AJ. Java Treeview: extensible visualization of microarray data. *Bioinformatics* 2004;20:3246-3248.
17. Shoulders MD, Raines RT. Collagen structure and stability. *Annu Rev Biochem* 2009;78:929-958.
18. Li A, Zhou T, Guo L, Si J. Collagen type I regulates beta-catenin tyrosine phosphorylation and nuclear translocation to promote migration and proliferation of gastric carcinoma cells. *Oncol Rep* 2010;23:1247-1255.
19. Zhang K, Corsa CA, Ponik SM, Prior JL, Piwnica-Worms D, Eliceiri KW, *et al.* The collagen receptor discoidin domain receptor 2 stabilizes SNAIL1 to facilitate breast cancer metastasis. *Nat Cell Biol* 2013;15:677-687.
20. Yan Z, Jin S, Wei Z, Huilian H, Zhanhai Y, Yue T, *et al.* Discoidin domain receptor 2 facilitates prostate cancer bone metastasis via regulating parathyroid hormone-related protein. *Biochim Biophys Acta* 2014;1842:1350-1363.
21. Xu Z, Choudhary S, Voznesensky O, Mehrotra M, Woodard M, Hansen M, *et al.* Overexpression of COX-2 in human osteosarcoma cells decreases proliferation and increases apoptosis. *Cancer Res* 2006;66:6657-6664.
22. Walmsley MJ, Ooi SK, Reynolds LF, Smith SH, Ruf S, Mathiot A, *et al.* Critical roles for Rac1 and Rac2 GTPases in B cell development and signaling. *Science* 2003;302:459-462.
23. Lucero CM, Vega OA, Osorio MM, Tapia JC, Antonelli M, Stein GS, *et al.* The cancer-related transcription factor Runx2 modulates cell proliferation in human osteosarcoma cell lines. *J Cell Physiol* 2013;228:714-723.
24. Baniwal SK, Khalid O, Gabet Y, Shah RR, Purcell DJ, Mav D, *et al.* Runx2 transcriptome of prostate cancer cells: insights into invasiveness and bone metastasis. *Mol Cancer* 2010;9:258.
25. Leong DT, Lim J, Goh X, Pratap J, Pereira BP, Kwok HS, *et al.* Cancer-related ectopic expression of the bone-related transcription factor RUNX2 in non-osseous metastatic tumor cells is linked to cell proliferation and motility. *Breast Cancer Res* 2010;12:R89.
26. Richmond TJ, Davey CA. The structure of DNA in the nucleosome core. *Nature* 2003;423:145-150.
27. Han M, Grunstein M. Nucleosome loss activates yeast downstream promoters *in vivo*. *Cell* 1988;55:1137-1145.
28. Lorch Y, LaPointe JW, Kornberg RD. Nucleosomes inhibit the initiation of transcription but allow chain elongation with the displacement of histones. *Cell* 1987;49:203-210.
29. Kim K, Punj V, Choi J, Heo K, Kim JM, Laird PW, *et al.* Gene dysregulation by histone variant H2A.Z in bladder cancer. *Epigenetics Chromatin* 2013;6:34.