

Curcumin Inhibits the Activation of Immunoglobulin E-Mediated Mast Cells and Passive Systemic Anaphylaxis in Mice by Reducing Serum Eicosanoid and Histamine Levels

Xian Li¹, Yue Lu², Ye Jin¹, Jong-Keun Son¹, Seung Ho Lee^{1,*} and Hyeun Wook Chang^{1,*}

¹College of Pharmacy, Yeungnam University, Gyeongsan 712-749, Republic of Korea,

²School of Pharmacy, Shanghai University of Traditional Chinese Medicine, Shanghai 201203, China

Abstract

Curcumin is naturally occurring polyphenolic compound found in turmeric and has many pharmacological activities. The present study was undertaken to evaluate anti-allergic inflammatory activity of curcumin, and to investigate its inhibitory mechanisms in immunoglobulin E (IgE)/Ag-induced mouse bone marrow-derived mast cells (BMMCs) and in a mouse model of IgE/Ag-mediated passive systemic anaphylaxis (PSA). Curcumin inhibited cyclooxygenase-2 (COX-2) dependent prostaglandin D₂ (PGD₂) and 5-lipoxygenase (5-LO) dependent leukotriene C₄ (LTC₄) generation dose-dependently in BMMCs. To probe the mechanism involved, we assessed the effects of curcumin on the phosphorylation of Syk and its downstream signal molecules. Curcumin inhibited intracellular Ca²⁺ influx via phospholipase C_γ1 (PLC_γ1) activation and the phosphorylation of mitogen-activated protein kinases (MAPKs) and the nuclear factor-κB (NF-κB) pathway. Furthermore, the oral administration of curcumin significantly attenuated IgE/Ag-induced PSA, as determined by serum LTC₄, PGD₂, and histamine levels. Taken together, this study shows that curcumin offers a basis for drug development for the treatment of allergic inflammatory diseases.

Key Words: Curcumin, Mast cell, Prostaglandin D₂, Leukotriene C₄, Mitogen activated protein kinase, Passive systemic anaphylaxis

INTRODUCTION

Anaphylaxis is an immediate, immunoglobulin E (IgE)-mediated hypersensitivity reaction resulting from the sudden release of mast cell- and basophil-derived chemical mediators such as histamine, serotonin, newly synthesized lipid-derived mediators, such as prostaglandin D₂ (PGD₂), leukotriene B₄ (LTB₄), cysteinyl leukotrienes LTC₄, LTD₄, and LTE₄, platelet-activating factor (PAF), and various cytokines (Kemp and Lockey, 2002; Simons, 2008). Furthermore, the releases of these chemical mediators are a consequence of the binding of IgE to FcεR1 on the surfaces of mast cells. This initial interaction results in the phosphorylations of tyrosine residues in the immune receptor tyrosine based activation motif (ITAM) of the β and γ subunits of FcεR1 by Lyn and recruits the tyrosine kinase Syk, which is required for phosphorylation of linker for activation of T cells (LAT). Phosphorylated LAT binds phospholipase C_γ (PLC_γ) and the adaptors Gads and Grb2 and generates inositol-1,4,5-triphosphate (IP₃), which

can increase intracellular calcium release from endoplasmic reticulum (ER) calcium stores (Siraganian, 2003). Elevated intracellular calcium then triggers the degranulation and the translocation of cytosolic phospholipase A₂ (cPLA₂) and 5-lipoxygenase (5-LO) to the nuclear membrane. The syntheses of cyclooxygenase-2 (COX-2)-dependent PGD₂ and of 5-LO-dependent LTC₄ in mast cells are initiated by the release of free arachidonic acid (AA) from membrane phospholipid by cPLA₂ (Kudo and Murakami, 1999; Yamaguchi *et al.*, 1999). Furthermore, it is well known that COX-2 dependent PGD₂ is very important to the development of inflammation and allergic disease such as asthma (Ricciotti and FitzGerald, 2011). LTC₄ plays important roles in inflammatory and allergic diseases (Werz and Steinhilber, 2006), and increasing evidence suggests it play a role in cancer (Avis *et al.*, 2001) and cardiovascular disease (Mehrabian and Allayee, 2003). Therefore, the inhibition of eicosanoid production constitutes an important therapeutic strategy against various allergic inflammatory diseases.

Open Access <http://dx.doi.org/10.4062/biomolther.2013.092>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received Nov 1, 2013 Revised Dec 9, 2013 Accepted Dec 11, 2013

*Corresponding Author

E-mail: hwchang@yu.ac.kr (Chang HW), seungho@yumail.ac.kr (Lee SH)
Tel: +82-53-810-2811 (Chang HW), +82-53-810-2815 (Lee SH)
Fax: +82-53-810-4654 (Chang HW), +82-53-810-4654 (Lee SH)
The first two authors contributed equally to this work.

Curcumin (diferuloyl methane) is a major constituent of the rhizome of *Curcuma longa*, and is used traditionally to treat inflammation, gastrointestinal disorders, hepatic disorders, diabetic wounds, skin wounds, rheumatism, sinusitis, and other disorders (Ammon and Wahl, 1991). Furthermore, scientific studies have shown that curcumin inhibits histamine release and the secretions of tumor necrosis factor- α (TNF- α) and interleukin-4 (IL-4) from mast cells triggered by IgE, calcium ionophore A23187, or compound 48/80 (Suzuki *et al.*, 2005; Lee *et al.*, 2008; Choi *et al.*, 2010). Curcumin has also been reported to inhibit IgE-induced type I hypersensitivity and ovalbumin-induced airway hyperreactivity (Yano *et al.*, 2000; Ram *et al.*, 2003; Lee *et al.*, 2008), and to inhibit COX-2 gene expression in phorbol ester-treated human gastrointestinal epithelial cells and mouse skin (Chun *et al.*, 2003; Ricciotti and FitzGerald, 2011) and *in vitro* lipoxygenase and cyclooxygenase activities in mouse epidermis (Huang *et al.*, 1991). However, the effect of curcumin on IgE/Ag-induced COX-2 dependent PGD₂ and 5-LO dependent LTC₄ generation in mast cells and IgE-mediated systemic anaphylactic response have not been well investigated.

In this study, we evaluated the effects of curcumin on the generation of eicosanoid (PGD₂ and LTC₄) in Fc ϵ RI-induced mast cells and on passive systemic anaphylaxis (PSA) response in mice.

MATERIALS AND METHODS

Plant material

Curcumin was isolated from the ethyl acetate fraction of a methanol extract of the rhizome of *Curcuma longa*, as described previously (Kiuchi *et al.*, 1993), and produced a single TLC spot and had a HPLC determined purity of >99.5%. Curcumin was prepared by dissolving it in dimethyl sulfoxide (DMSO). The final concentration of DMSO in culture media was adjusted to 0.1% (v/v). DMSO alone was run as a control in all cases.

Chemicals and reagents

Mouse anti-dinitrophenyl (DNP) IgE and DNP-human serum albumin (HSA) were purchased from Sigma Chemical Co. (St. Louis, MO, USA). Fexofenadine-HCl (Fexo), a histamine H1 receptor antagonist, was obtained from Korea Pharma (Seoul, Korea). The rabbit polyclonal antibodies specific for phospho-I κ B, IKK α / β , ERK1/2, JNK, p38, Akt, β -actin, and total form for I κ B, ERK1/2, JNK, p38, and Akt were purchased from Cell Signaling Technology, Inc. (Danvers, MA, USA). Rabbit polyclonal antibodies for phospho-cPLA₂ α (Ser505), cPLA₂ α , 5-LO, PLC γ 1, IKK α / β lamin B and NF- κ B p65 as well as secondary goat anti-rabbit IgG-HRP and rabbit anti-goat IgG-HRP antibodies, total Syk, total LAT, and Bay 61-3606 were purchased from Santa Cruz Biotechnology (Dallas, Texas, USA) and antibodies for phosphotyrosine was purchased from Millipore (Millipore, Billerica, MA, USA). The antibody-reactive bands were visualized with an enhanced chemiluminescence (ECL) system (Pierce Biotechnology, Rockford, IL, USA). The enzyme immunoassay (EIA) kits for PGD₂, LTC₄, histamine and the antibody for COX-2 were purchased from Cayman Chemicals (Ann Arbor, MI, USA).

Culture and activation of bone marrow derived mast cells (BMMCs)

BMMCs were isolated from bone marrow of C57BL/6 mice and differentiated as described previously (Lu *et al.*, 2011). Briefly, BMMCs were cultured in RPMI 1640 medium containing 10% fetal bovine serum, 100 U/ml penicillin, 10 mM HEPES, 100 μ M MEM non-essential amino acid solution (Invitrogen, Grand Island, NY, USA) and 20% pokeweed mitogen-spleen cell conditioned medium as a source of IL-3. For stimulation, 10⁶ cells/ml were sensitized overnight with 500 ng/ml anti-DNP IgE, pretreated with indicated concentration of curcumin or Bay 61-3606, and stimulated for appropriate periods with 100 ng/ml DNP-HSA. The reactions were terminated by centrifugation of the cells at 3,000 rpm for 5 min at 4°C.

Determination of LTC₄ and PGD₂

Concentration of LTC₄ and PGD₂ were determined as described previously (Lu *et al.*, 2011). IgE sensitized BMMCs suspended in enriched medium at a cell density of 1 \times 10⁶ cells/ml were pretreated with indicated concentration of curcumin or Bay 61-3606 for 1 h and stimulated with DNP-HSA for 15 min. Supernatants were isolated for further analysis by EIA kit. The concentration of LTC₄ was determined using an EIA kit. To assess COX-2-dependent PGD₂ synthesis, BMMCs were preincubated with 1 μ g/ml of aspirin for 2 h to irreversibly inactivate preexisting COX-1. After washing, BMMCs were incubated with 100 ng/ml DNP-HSA at 37°C for 7 h in the presence of curcumin or Bay 61-3606. PGD₂ in the supernatants were quantified using PGD₂ EIA kit and cells were used for immunoblots analysis. Under the conditions employed, LTC₄ reached 4.75 ng/10⁶ cells and PGD₂ generation reached 2.12 ng/10⁶ cells. All data were the arithmetic mean of triplicate determinations.

Measurement of intracellular Ca²⁺ level

Intracellular Ca²⁺ levels were determined using FluoForte™ Calcium Assay Kit (Enzo Life Sciences, Ann Arbor, MI, USA), as described previously (Hwang *et al.*, 2013). Briefly, BMMCs (1 \times 10⁶ cells) were sensitized overnight with 500 ng/ml anti-DNP IgE. Sensitized BMMCs were preincubated with FluoForte™ Dye-Loading Solution for 1 h at room temperature. After washing the dye from cell surface with HBSS, cells (5 \times 10⁴) were seeded into 96-well microplates and pretreated with curcumin or Bay 61-3606 for 1 h before adding DNP-HSA. Fluorescence was measured using a fluorometric imaging plated reader at an excitation wavelength of 485 nm and an emission wavelength of 520 nm on a BMG Labtechnologies FLUOStar OPITIMA platereader (Offenburg, Germany).

Preparation of nuclear and cytoplasmic extracts

The nuclear and cytoplasmic extracts were prepared as described previously (Lu *et al.*, 2011). BMMCs were sensitized to DNP-specific IgE (500 ng/ml, overnight) and pretreated with curcumin or Bay 61-3606 for 1 h, and then stimulated with DNP-HSA (100 ng/ml) for 30 min. Cultured BMMCs were collected by centrifugation, washed with PBS, and lysed in a buffer containing 10 mM HEPES (pH 7.9), 10 mM KCl, 0.1 mM EDTA, 0.1 mM EGTA, 1 mM DTT, 1 mM PMSF and 0.1% NP40 on ice for 10 min. Supernatants (cytosolic fractions) were obtained by centrifugation at 1,000 *g* for 4 min. Nuclear pellets were washed and lysed in a buffer containing 20 mM HEPES (pH 7.9), 25% (v/v) glycerol, 420 mM NaCl, 1.5 mM




Fig. 1. Curcumin inhibited IgE/Ag-induced PSA reaction. In the PSA test, ICR mice were sensitized by injecting 2 μ g of IgE *i.v.* in 100 μ l saline or injected with saline alone. 24 h later mice were administered 25 or 50 mg/kg of curcumin or 50 mg/kg fexofenadine-HCl (Fexo) and 1 h later were challenged with 4 mg of DNP-HSA *i.v.* in 200 μ l saline; blood was collected by cardiac puncture 5 min after the Ag challenge. The concentration of serum LTC₄ (A), PGD₂ (B), and histamine (C) were determined using appropriate enzyme immunoassay kits (Cayman Chemicals). The values indicate the mean \pm S.D. from three independent experiments, ** p <0.01 versus IgE/Ag sensitized mice). Fexo (50 mg/kg) was used as an anti-histamine control drug.

MgCl₂, 0.2 mM EDTA, and protease inhibitor cocktail.

Immunoprecipitation (IP)

Immunoprecipitation was performed as described previously (Hwang *et al.*, 2013). Briefly, cell lysates were obtained using modified lysis buffer [0.1% Nonidet P-40, 50 mM HEPES (pH 7.0), 250 mM NaCl, 5 mM EDTA, 1 mM PMSF, and 0.5 mM dithiothreitol]. Total cell lysates (1 mg protein equivalent) were incubated with anti-Syk or anti-LAT antibodies for 2 h at 4°C and immunocomplexes were precipitated with 20 μ l of protein A-Sepharose and washed 3 times with ice-cold lysis buffer. Precipitates and total cell lysates were subjected to SDS-PAGE and immunoblotted with appropriate antibodies.

Immunoblotting

Immunoblotting was performed as described previously (Lu *et al.*, 2011). Cells were lysed in RIPA lysis buffer (50 mM Tris-HCl [pH 8.0], 150 mM NaCl, 5 mM EDTA, 1% Nonidet P-40, 1mM phenylmethanesulfonyl fluoride (PMSF), 1M dithiothreitol (DTT), 200 mM NaF, 200 mM Na₃VO₄, and protease inhibitor cocktail). Cell debris was removed by centrifugation at 14,000 g for 15 min at 4°C, and the resulting supernatant was western blotted. Protein concentration was measured using a Qubit Fluorometer (Invitrogen, USA). Samples were separated by 8% SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and then transferred to nitrocellulose membranes (Millipore, Billerica, MA, USA). Immunoreactive proteins were incubated with HRP-coupled secondary antibodies diluted at 1:3,000 for 1 h at room temperature, and then developed using an enhanced chemiluminescence (ECL) detection kit.

Passive systemic anaphylaxis (PSA) reaction in mice

PSA was induced as described previously (Lu *et al.*, 2011). Six ICR mice (Hyochoang Science, Daegu, Korea) were placed per cage within a laminar airflow cabinet and fed standard laboratory chow (Purina, Seoul, Korea) and provided with water *ad libitum*. The animals were maintained in 22 \pm 1°C, 55 \pm 10% RH environment under a 12 h/12 h (light/dark) cycle for at least 7 days prior to experiments. Mice were injected with 2 μ g of DNP-specific IgE in 100 μ l of saline or with 100 μ l of saline. 24 h later mice were orally administered curcumin (20 and 50 mg/kg) or fexofenadine-HCl (Fexo; 50 mg/kg) and 1 h

later challenged with 4 mg *i.v.* of DNP-HSA in 200 μ l of saline. Blood was collected by cardiac puncture 5 min after the Ag challenge. Serum LTC₄, PGD₂ and histamine concentrations were determined using appropriate EIA kits. All experiments using animals were approved beforehand by the Institutional Animal Care and Use Committee of Yeungnam University.

Statistical analysis

All experiments were performed three or more times. Average values are expressed as means \pm S.D. Statistical analyses were performed using SPSS 19.0 (SPSS, Chicago, IL, USA). The Student's *t*-test was used to compare pairs of independent groups. Statistical significance was accepted for p values <0.05.

RESULTS

Curcumin suppressed passive systemic anaphylaxis (PSA) in mice

Anaphylaxis is an IgE-mediated hypersensitivity reaction caused by the release of various inflammatory mediators due to binding of specific IgE to Fc ϵ R1 on the surfaces of mast cells or basophils (Siraganian, 2003). Furthermore, it has been reported that curcumin suppresses IgE or compound 48/80 induced passive cutaneous anaphylaxis (Suzuki *et al.*, 2005; Lee *et al.*, 2008; Choi *et al.*, 2010). However, no report has been issued on the effect of curcumin on PSA in mice. Therefore, we assessed the anti-allergic-inflammatory effects of curcumin using a mouse PSA model. As shown in Fig. 1, PSA was induced using an *i.v.* injection of DNP-HSA in ICR mice after the oral administration of 25 or 50 mg/kg of curcumin or 50 mg/kg of Fexo (Ciprandi *et al.*, 2003). One hour later, curcumin reduced serum LTC₄ (Fig. 1A), PGD₂ (Fig. 1B) and histamine (Fig. 1C) levels in a dose-dependent manner (n=6). The suppressive effect of 50 mg/kg of curcumin was similar to that of 50 mg/kg of Fexo, a histamine H1 receptor antagonist.

Curcumin inhibited LTC₄ generation and Ca²⁺ influx in IgE/Ag-induced BMMC

In vivo results let us to investigate the action mechanism responsible for the anti-allergic inflammatory activities of cur-




Fig. 2. Effect of Curcumin on LTC₄ generation and on Ca²⁺ mobilization in IgE/Ag-activated BMMCs. IgE-sensitized BMMCs were pre-incubated with the indicated concentration of curcumin or Bay61-3606 for 1 h and then stimulated with DNP-HSA for 15 min. LTC₄ released into the supernatant was quantified using an enzyme immunoassay kit (A). Relative intracellular Ca²⁺ levels were determined (at 5 min) (B). **p*<0.05, ***p*<0.01 and ****p*<0.001 versus the IgE/Ag-treated group. Results are the averages of three independent experiments.




Fig. 3. Effects of curcumin and Bay 61-3606 on cPLA₂ α and 5-LO translocation and MAPKs activation. IgE-sensitized BMMCs were pre-incubated for 1 h with the indicated concentrations of curcumin or Bay 61-3606 and then stimulated with DNP-HSA for 15 min. Cytosolic and nuclear fractions were immunoblotted with antibodies for phospho-cPLA₂ α (Ser505) and 5-LO (A), and cell lysates were immunoblotted for the total and phosphorylated forms of ERK1/2, JNK and p38 (B). Immunoblots of β -actin and lamin B were used as controls for cytosol and nuclear fractions, respectively.




Fig. 4. Effect of Curcumin on COX-2 dependent PGD₂ generation and on Akt-NF- κ B activation. IgE-sensitized BMMCs were pre-incubated with the indicated concentration of curcumin or Bay61-3606 for 1 h and then stimulated with DNP-HSA for 7 h. PGD₂ released into the supernatant was quantified using an enzyme immunoassay kit and the cells were used for immunoblotting of COX-2 protein (A). IgE-sensitized BMMCs were pre-incubated with curcumin or Bay 61-3606 for 1 h and stimulated with DNP-HSA for 15 min. Then cells were taken for the immunoblot detection of Akt, IKK, I κ B α cytosolic NF- κ B (p65), nuclear NF- κ B (p65), β -actin and lamin B. Nuclear extracts were used for the NF- κ B (p65) immunoblot. Results are presented as mean \pm S.D. of three independent experiments. ***p*<0.01 versus the IgE/Ag-treated group.

curcumin using mast cells, major players in PSA. LTC₄ is metabolites of arachidonic acid derived from action of 5-LO/LTC₄ synthase in mast cells, and LTC₄ have been implicated in inflammation, proliferation and allergic conditions like asthma (Murphy and Gijon, 2007). Thus, we investigated the effects of curcumin on 5-LO dependent LTC₄ generation in BMMCs. As shown in Fig. 2A, curcumin strongly inhibited LTC₄ generation in a dose-dependent manner. It is well known that Ca²⁺ is essential for arachidonic acid (AA) release from phospholipid and degranulation in IgE/Ag-induced mast cells (Kudo and Murakami, 1999; Yamaguchi *et al.*, 1999), thus we examined the effect of curcumin on intracellular Ca²⁺ influx in IgE/Ag-induced BMMCs. As shown in Fig. 2B, intracellular Ca²⁺ level in activated BMMCs was about three fold higher than in resting cells, and this increase was dose-dependently inhibited by curcumin. Bay 61-3606 (a Syk inhibitor) also strongly decreased intracellular LTC₄ generation as well as Ca²⁺ influx. Consistent with a previous report (Lee *et al.*, 2008), the release of β -Hex (a degranulation marker enzyme) was dose-dependently inhibited by curcumin (data not shown).

Curcumin inhibited cPLA₂ phosphorylation, translocation of phospho-cPLA₂ α and 5-LO and activation of MAPKs

Recently, we and other group have reported that the release of free AA from membrane phospholipid in activated mast cells requires the phosphorylation of cPLA₂ α (p-cPLA₂ α) by mitogen activated protein kinases (MAPKs) (Lin *et al.*, 1993; Lu *et al.*, 2011), and the translocation of p-cPLA₂ α is dependent on intracellular Ca²⁺ influx (Gijon and Leslie, 1999; Lu *et al.*, 2011). To determine whether curcumin inhibits the phosphorylation and translocation of cPLA₂ α , we pretreated BMMCs with different concentrations of curcumin or Bay 61-3606. As shown in Fig. 3A, curcumin dose-dependently inhibited the phosphorylation (C-p-cPLA₂ α) and translocation of C-p-cPLA₂ α to the nuclear envelope (N-p-cPLA₂ α). The synthesis of 5-LO dependent LTC₄ in IgE/Ag-induced mast cells is known to mediate the translocations of both p-cPLA₂ α and 5-LO to the nuclear envelope (Verz, 2002; Lu *et al.*, 2011; Lu *et al.*, 2012). Thus, we investigated the effect of curcumin on the translocation of 5-LO to the nuclear envelope. As was expected, curcumin or Bay 61-3606 inhibited the translocation of cytosolic 5-LO (C-5-LO) to nuclear envelope (N-5-LO). Next, to confirm that the inhibition of cPLA₂ α phosphorylation by curcumin occurred via the inhibition of MAPKs phosphorylation including extracellular regulated kinase1/2 (ERK1/2), c-jun N-terminal kinase (JNK), and p38 MAP kinase, therefore we examined the effect of curcumin or Bay 61-3606 on the phosphorylation of MAPKs. As shown in Fig. 3B, curcumin or Bay 61-3606 inhibited the phosphorylations of three MAPKs in a dose dependent manner.

Curcumin inhibits COX-2 dependent PGD₂ generation and NF- κ B activation

In mast cells, unlike 5-LO dependent LTC₄ generation, PGD₂ generation occurs in a biphasic manner. Immediate PGD₂ generation (occurring within 2 h), is associated with COX-1, and delayed PGD₂ generation (during 2-10 h) is occurred by inducible COX-2 protein (Ashraf *et al.*, 1996; Moon *et al.*, 1998). To assess COX-2-dependent delayed PGD₂ generation, BMMCs were pre-treated with aspirin to abolish preexisting COX-1 activity, followed by a brief wash, and then stimulated with Ag for 7 h with or without curcumin. As shown in Fig 4A, delayed




Fig. 5. Effect of curcumin on the Syk pathway. IgE-sensitized BMMCs were preincubated with curcumin or Bay 61-3606 for 1 h, and then stimulated with DNP-HSA for 5 min. Cell lysates were subjected to immunoprecipitation and immunoblot analysis for the phosphorylated forms of Syk, LAT and PLC γ 1. Bay 61-3606 was used as a positive control with respect to the suppression of the Syk-mediated pathway. The relative ratios of p-Syk/Syk, p-LAT/LAT and p-PLC γ 1/PLC γ 1 protein levels were determined by measuring immunoblot band intensities by scanning densitometry (** $p < 0.01$ and *** $p < 0.001$). The results shown are representative of three independent experiments.

PGD₂ generation was dose-dependently inhibited by curcumin, with a concomitant suppression of COX-2 protein expression. Yu *et al.*, reported that the Syk downstream molecules PI3K/Akt pathway affects transcription factor NF-κB activation in gastric cancer cells and mast cells (Yu *et al.*, 2010; Lu *et al.*, 2011) and NF-κB has been identified as an essential transcription factor for the induction of several inflammatory mediators including, TNF-α, COX-2, and inducible NO synthase (Reddy *et al.*, 2000; Tak and Firestein, 2000; Lu *et al.*, 2011). Thus, we examined the effect of curcumin on Akt/NF-κB axis activation. When IgE-sensitized BMMCs were pretreated with curcumin for 1 h and then stimulated with Ag for 15 min, phosphorylation of the Akt, IKK complex (p-IKKα/β) and IκBα (p-IκBα) was increased, with a concomitant decrease of total IκBα and nuclear translocation of NF-κB (C-NF-κB). As shown in Fig. 4B, both curcumin and Bay 61-3606 inhibited the phosphorylations of Akt, IKKα/β, IκBα, IκBα degradation and the translocation of cytosolic p65 to nuclear (N-NF-κB), suggesting that Syk mediated Akt/NF-κB pathway regulate the reduction of COX-2 dependent PGD₂ by curcumin.

Curcumin inhibited the Syk pathway in IgE/Ag-induced BMMCs

Previously, we and others reported that Syk plays an essential role in the initiation of FcεRI-induced mast cells activation and mediates LAT, resulting in the activation of downstream signaling, including MAPKs, phosphatidylinositol 3-kinase (PI3K), and PLCγ (Lin *et al.*, 1993; Lu *et al.*, 2011). Furthermore, it has been reported that curcumin inhibited the releases of TNF-α and IL-4 from mast cells via a Syk dependent pathway, and that the inhibitions of these secretions by curcumin is dependent on its direct inhibition of Syk kinase activity rather than Syk phosphorylation (Lee *et al.*, 2008). To determine whether curcumin affects Syk phosphorylation in IgE/Ag-induced BMMCs, we examined the effects of curcumin on the phosphorylation of Syk and its downstream signal molecules LAT and PLCγ1. As shown in Fig. 5A, curcumin did not affect Syk phosphorylation, but significantly and dose-dependently inhibited the phosphorylations of LAT and PLCγ1 (Fig. 5B, C), as previously reported (Lee *et al.*, 2008). Bay 61-3606 (the Syk inhibitor used as a positive control) completely inhibited the phosphorylations of Syk, LAT and PLCγ1. These results suggest that curcumin inhibits PGD₂, LTC₄, and degranulation by regulating the Syk signal pathway.

DISCUSSION

Curcumin has been shown to have diverse biological activities, such as, antioxidant, anti-allergic, anti-inflammatory, antiviral, antibacterial, antifungal, and anticancer activities (Ammon and Wahl, 1991). Several groups have reported that curcumin shows anti-allergic activity through the inhibition of histamine release, TNF-α and IL-4 from activated mast cells and *in vivo* type I hypersensitivity animal model (Yano *et al.*, 2000; Ram *et al.*, 2003; Lee *et al.*, 2008). In addition, curcumin also suppresses arachidonic acid metabolizing enzymes such as cPLA₂ phosphorylation, COX-2 expression and recombinant 5-LO activity in LPS-stimulated RAW 264.7 cells and A23187-stimulated HT human colon cancer cells (Hong *et al.*, 2004) and inhibits *in vitro* LOX and COX activities in TPA- and arachidonic acid-induced inflammation in mouse epidermis (Huang *et al.*, 1991). Furthermore, Curcumin modulates

the inflammatory response by down-regulating the activity of COX-2 and inducible nitric oxide synthase (iNOS) enzymes through suppression of NF-κB activation (Surh *et al.*, 2001). Even though they reported that curcumin inhibited generation of COX-dependent PGs and LOX-dependent hydroxyeicosatetraenoic acids (HETEs), previous reports were mainly examined as a part of anticancer activity of curcumin. However, the effect of curcumin on generation of COX-2 dependent PGD₂ and 5-LO dependent LTC₄ generation in IgE/Ag-induced mast cells and IgE-mediated PSA reaction have not been studied to date. Therefore, we investigated the effects of curcumin on the generation of eicosanoid (PGD₂ and LTC₄) in BMMCs and on PSA reaction in mice. It has been well established that lipid mediators, like LTC₄ and PGD₂, are closely associated with various allergic and inflammatory diseases (Werz and Steinhilber, 2006; Ricciotti and FitzGerald, 2011). Thus, the inhibition of LTC₄ and PGD₂ generation by mast cells is an important therapeutic strategy in the context of allergic-inflammatory disease. The aim of this study was to investigate the effect of curcumin on IgE/Ag-induced COX-dependent PGD₂ and 5-LO dependent LTC₄ generation in BMMCs and on IgE-induced passive systemic anaphylaxis (PSA) in mice. IgE binds to FcεRI on mast cells, the Syk-LAT axis activates PLCγ, which increases intracellular Ca²⁺ influx and leads to degranulation and eicosanoid production (Siraganian, 2003). As described above, IgE binds to FcεRI on BMMCs, it promptly elicits 5-LO dependent LTC₄ generation, which is inhibited dose-dependently by curcumin (Fig. 2A). It has been reported that the synthesis of LTC₄ in mast cells is regulated by two steps, namely, AA release from phospholipid by cPLA₂α, MAPKs-mediated phosphorylation of cPLA₂α and the conversion of free AA to LTC₄ by 5-LO (Fischer *et al.*, 2005). Translocation of both 5-LO and phospho-cPLA₂α (C-p-cPLA₂α) to the nuclear envelope are depend on the increase of cytosolic Ca²⁺ level (Werz, 2002; Lu *et al.*, 2011; Lu *et al.*, 2012). The present study showed that curcumin inhibited the translocations of both enzymes, which concurred with its observed inhibitory effect on intracellular Ca²⁺ influx (Fig. 2B). Next, to elucidate the effect of curcumin on COX-2 dependent delayed PGD₂ generation, BMMCs were pre-treated with aspirin to abolish preexisting COX-1 activity, and then stimulated with Ag for 7 h with or without curcumin. We found that curcumin also suppressed COX-2 expression and attendant PGD₂ generation (Fig. 4A). It has been previously reported that curcumin inhibited the TPA-induced up-regulation of COX-2 and MMP-9 by suppressing ERK1/2 phosphorylation and NF-κB transactivation in epithelial cells (Lee *et al.*, 2005), and that it exerts anti-inflammatory effects by inhibiting the NF-κB and MAPKs pathways (Cho *et al.*, 2007). Recently, we also reported that COX-2-dependent PGD₂ generation and COX-2 expression in BMMCs occurs via the activation of the NF-κB and MAPKs pathways (Lu *et al.*, 2011; Lu *et al.*, 2012), thus we investigated the effect of curcumin on the MAPKs and NF-κB pathways in IgE/Ag-induced BMMCs. As shown in Fig. 3B, curcumin suppressed the phosphorylations of ERK1/2, JNK, and p38 MAP kinase in a dose dependent manner, which implied that it affects the phosphorylation and translocation of cPLA₂. Since curcumin or Bay 61-3606 suppressed intracellular Ca²⁺ influx (Fig. 2B), MAPKs phosphorylation (Fig. 3B) and Akt/ NF-κB axis activation, these results suggest that Syk plays an important role in the generation of PGD₂ and LTC₄ in IgE/Ag-induced BMMCs. Therefore, we examined whether curcumin affects the phos-

phorylation of Syk in IgE/Ag-induced BMMCs. In agreement with a previous report (Lee *et al.*, 2008), curcumin was found not to directly inhibit the phosphorylation of Syk, but to inhibit the phosphorylations of LAT and PLC γ 1 which lie downstream of Syk. In view of the effect of phosphorylated PLC γ 1 on inositol phospholipid turnover and the consequent increase in Ca $^{2+}$ influx, it is likely that the observed inhibition of Ca $^{2+}$ influx by curcumin (Fig. 2B) depends on its inhibitory effect on Syk-dependent PLC γ 1 phosphorylation. In addition, to our *in vitro* results, curcumin also suppressed the IgE-dependent PSA reaction in a mast cell-dependent *in vivo* model of systemic allergic reaction (Wershil *et al.*, 1987) with a potency equivalent to that of the H1 histamine antagonist, Fexo (Fig. 1A-C). We already reported that PGD $_2$ and LTC $_4$ play important roles in mast cell-mediated anaphylactic reaction (Lu *et al.*, 2011; Hwang *et al.*, 2013). Recently, several reports have demonstrated that Syk kinase inhibitors promised for the treatment of allergic and antibody-mediated autoimmune diseases (Ruzza *et al.*, 2009) and clinical implications of Syk inhibitor showed antiallergic properties when administered orally (Mazuc *et al.*, 2008). Taken together with previous results (Lee *et al.*, 2008), the anti-allergic inflammatory activity of curcumin appear to be due to the suppressions of the secretions of TNF- α and IL-4 and histamine release and eicosanoid generation through the inhibition of Syk kinase pathway in IgE/Ag-induced mast cells.

ACKNOWLEDGMENTS

This work was supported by Yeungnam University 2013 grant.

REFERENCES

- Ammon, H. P. and Wahl, M. A. (1991) Pharmacology of Curcuma longa. *Planta Med.* **57**, 1-7.
- Ashraf, M., Murakami, M. and Kudo I. (1996) Cross-linking of the high-affinity IgE receptor induces the expression of cyclo-oxygenase 2 and attendant prostaglandin generation requiring interleukin 10 and interleukin 1 beta in mouse cultured mast cells. *Biochem. J.* **320**, 965-973.
- Avis, I., Hong, S. H., Martinez, A., Moody, T., Choi, Y. H., Trepel, J., Das, R., Jett, M. and Mulshine, J. L. (2001) Five-lipoxygenase inhibitors can mediate apoptosis in human breast cancer cell lines through complex eicosanoid interactions. *FASEB J.* **15**, 2007-2009.
- Cho, J. W., Lee, K. S. and Kim, C. W. (2007) Curcumin attenuates the expression of IL-1 β , IL-6, and TNF- α as well as cyclin E in TNF- α -treated HaCaT cells; NF- κ B and MAPKs as potential upstream targets. *Int. J. Mol. Med.* **19**, 469-474.
- Choi, Y. H., Yan, G. H., Chai, O. H. and Song, C. H. (2010) Inhibitory effects of curcumin on passive cutaneous anaphylactoid response and compound 48/80-induced mast cell activation. *Anat. Cell Biol.* **43**, 36-43.
- Chun, K. S., Keum, Y. S., Han, S. S., Song, Y. S., Kim, S. H. and Surh, Y. J. (2003) Curcumin inhibits phorbol ester-induced expression of cyclooxygenase-2 in mouse skin through suppression of extracellular signal regulated kinase activity and NF- κ B activation. *Carcinogenesis* **24**, 1515-1524.
- Ciprandi, G., Tosca, M. A., Cosentino, C., Riccio, A. M., Passalacqua, G. and Canonica, G. W. (2003) Effects of fexofenadine and other antihistamines on components of the allergic response: adhesion molecules. *J. Allergy Clin. Immunol.* **112**, S78-82.
- Fischer, L., Poekkel, D., Buerker, E., Steinhilber, D. A D Werz, O. (2005) Inhibitors of actin polymerization stimulate arachidonic acid release and 5-lipoxygenase activation by upregulation of Ca $^{2+}$ mobilisation in polymorphonuclear leukocytes involving Src family kinases. *Biochim. Biophys. Acta* **1736**, 109-119.
- Gijon, M. A. and Leslie, C. C. (1999) Regulation of arachidonic acid release and cytosolic phospholipase A2 activation. *J. Leukoc. Biol.* **65**, 330-336.
- Hong, J., Bose, M., Ju, J., Ryu, J. H., Chen, X., Sang, S., Lee, M. J. and Yang, C. S. (2004) Modulation of arachidonic acid metabolism by curcumin and related β -diketone derivatives: effects on cytosolic phospholipase A2, cyclooxygenases and 5-lipoxygenase. *Carcinogenesis* **25**, 1671-1679.
- Huang, M. T., Lysz, T., Ferraro, T., Abidi, T. F., Laskin, J. D. and Conney, A. H. (1991) Inhibitory effects of curcumin on *in vitro* lipoxygenase and cyclooxygenase activities in mouse epidermis. *Cancer Res.* **51**, 813-819.
- Hwang, S. L., Li, X., Lu, Y., Jin, Y., Jeong, Y. T., Kim, Y. D., Lee, I. K., Taketomi, Y., Sato, H., Cho, Y. S., Murakami, M. and Chang, H. W. (2013) AMP-activated protein kinase negatively regulates Fc ϵ RI-mediated mast cell signaling and anaphylaxis in mice. *J. Allergy Clin. Immunol.* **132**, 729-736.
- Kemp, S. F. and Lockey, R. F. (2002) Anaphylaxis: a review of causes and mechanisms. *J. Allergy Clin. Immunol.* **110**, 341-348.
- Kiuchi, F., Goto, Y., Sugimoto, N., Akao, N., Kondo, K. and Tsuda, Y. (1993) Nematocidal activity of turmeric: synergistic action of curcuminoids. *Chem. Pharm. Bull.* **41**, 1640-1643.
- Kudo, I. and Murakami, M. (1999) Diverse functional coupling of prostanoic biosynthetic enzymes in various cell types. *Adv. Exp. Med. Biol.* **469**, 29-35.
- Lee, J. H., Kim, J. W., Ko, N. Y., Mun, S. H., Her, E., Kim, B. K., Han, J. W., Lee, H. Y., Beaven, M. A., Kim, Y. M. and Choi, W. S. (2008) Curcumin, a constituent of curry, suppresses IgE-mediated allergic response and mast cell activation at the level of Syk. *J. Allergy Clin. Immunol.* **121**, 1225-1231.
- Lee, K. W., Kim, J. H., Lee, H. J. and Surh, Y. J. (2005) Curcumin inhibits phorbol ester-induced up-regulation of cyclooxygenase-2 and matrix metalloproteinase-9 by blocking ERK1/2 phosphorylation and NF- κ B transcriptional activity in MCF10A human breast epithelial cells. *Antioxid. Redox. Signal.* **7**, 1612-1620.
- Lin, L. L., Wartmann, M., Lin, A. Y., Knopf, J. L., Seth, A. and Davis, R. J. (1993) cPLA2 is phosphorylated and activated by MAP kinase. *Cell* **72**, 269-278.
- Lu, Y., Suh, S. J., Li, X., Hwang, S. L., Li, Y., Hwangbo, K., Park, S. J., Murakami, M., Lee, S. H., Jahng, Y., Son, J. K., Kim, C. H. and Chang, H. W. (2012) Citreorosein, a naturally occurring anthraquinone derivative isolated from *Polygoni cuspidati radix*, attenuates cyclooxygenase-2-dependent prostaglandin D2 generation by blocking Akt and JNK pathways in mouse bone marrow-derived mast cells. *Food Chem. Toxicol.* **50**, 913-919.
- Lu, Y., Yang, J. H., Li, X., Hwangbo, K., Hwang, S. L., Taketomi, Y., Murakami, M., Chang, Y. C., Kim, C. H., Son, J. K. and Chang, H. W. (2011) Emodin, a naturally occurring anthraquinone derivative, suppresses IgE-mediated anaphylactic reaction and mast cell activation. *Biochem. Pharmacol.* **82**, 1700-1708.
- Mazuc, E., Villoutreix, B. O., Malbec, O., Roumier, T., Fleury, S., Leonetti, J. P., Dombrowicz, D., Daëron, M., Martineau, P. and Dariavach, P. (2008) A novel druglike spleen tyrosine kinase binder prevents anaphylactic shock when administered orally. *J. Allergy Clin. Immunol.* **122**, 188-194.
- Mehrabian, M. and Allayee, H. (2003) 5-lipoxygenase and atherosclerosis. *Curr. Opin Lipidol.* **14**, 447-457.
- Moon, T. C., Murakami, M., Ashraf, M. D., Kudo, I. and Chang H. W. (1998) Regulation of cyclooxygenase-2 and endogenous cytokine expression by bacterial lipopolysaccharide that acts in synergy with c-kit ligand and Fc epsilon receptor I crosslinking in cultured mast cells. *Cell. Immunol.* **185**, 146-152.
- Murphy, R. C. and Gijon, M. A. (2007) Biosynthesis and metabolism of leukotrienes. *Biochem. J.* **405**, 379-395.
- Ram, A., Das, M. and Ghosh, B. (2003) Curcumin attenuates allergen-induced airway hyperresponsiveness in sensitized guinea pigs. *Biol. Pharm. Bull.* **26**, 1021-1024.
- Reddy, S. T., Wadleigh, D. J. and Herschman, H. R. (2000) Transcriptional regulation of the cyclooxygenase-2 gene in activated mast cells. *J. Biol. Chem.* **275**, 3107-3113.

- Ricciotti, E. and FitzGerald, G. A. (2011) Prostaglandins and Inflammation. *Arterioscler. Thromb. Vasc. Biol.* **31**, 986-1000.
- Ruzza, P., Biondi, B. and Calderan, A. (2009) Therapeutic prospect of Syk inhibitors. *Expert Opin. Ther. Pat.* **19**, 1361-1376.
- Simons, F. E. (2008) Anaphylaxis. *J. Allergy Clin. Immunol.* **121**, S402-407.
- Siraganian, R. P. (2003) Mast cell signal transduction from the high-affinity IgE receptor. *Curr. Opin. Immunol.* **15**, 639-646.
- Surh, Y. J., Chun, K. S., Cha, H. H., Han, S. S., Keum, Y. S., Park, K. K. and Lee, S. S. (2001). Molecular mechanisms underlying chemopreventive activities of anti-inflammatory phytochemicals: down-regulation of COX-2 and iNOS through suppression of NF-kappa B activation. *Mutat. Res.* **480-481**, 243-268.
- Suzuki, M., Nakamura, T., Iyoki, S., Fujiwara, A., Watanabe, Y., Mohri, K., Isobe, K., Ono, K. and Yano, S. (2005) Elucidation of anti-allergic activities of curcumin-related compounds with a special reference to their anti-oxidative activities. *Biol. Pharm. Bull.* **28**, 1438-1443.
- Tak, P. P., and Firestein, G. S. (2000) NF-kappaB: a key role in inflammatory diseases. *J. Clin. Invest.* **107**, 7-11.
- Wershil, B. K., Mekori, Y. A., Murakami, T. and Galli, S. J. (1987) 125I-fibrin deposition in IgE-dependent immediate hypersensitivity reactions in mouse skin. Demonstration of the role of mast cells using genetically mast cell-deficient mice locally reconstituted with cultured mast cells. *J. Immunol.* **139**, 2605-2614.
- Werz, O. (2002) 5-lipoxygenase: cellular biology and molecular pharmacology. *Curr. Drug Targets Inflamm. Allergy* **1**, 23-44.
- Werz, O. and Steinhilber, D. (2006) Therapeutic options for 5-lipoxygenase inhibitors. *Pharmacol. Ther.* **112**, 701-718.
- Yano, S., Terai, M., Shimizu, K. L., Futagami, Y., Sekine, T., Takamoto, K., Saito, K., Ueno, K. and Watanabe, K. (2000) Antiallergic activity of Curcuma longa. (II). Features of inhibitory actions on histamine release from mast cells. *Nat. Med.* **54**, 325-329.
- Yamaguchi, M., Sayama, K., Yano, K., Lantz, C. S., Noben-Trauth N., Ra C., Costa J. J. and Galli, S. J. (1999) IgE enhances Fc epsilon receptor I expression and IgE-dependent release of histamine and lipid mediators from human umbilical cord blood-derived mast cells: synergistic effect of IL-4 and IgE on human mast cell Fc epsilon receptor I expression and mediator release. *J. Immunol.* **162**, 5455-5465.
- Yu, L. L., Dai, N., Yu, H.G., Sun, L. M. and Si, J. M. (2010) Akt associates with nuclear factor kappaB and plays an important role in chemoresistance of gastric cancer cells. *Oncol. Rep.* **24**, 113-119.