

Efficient Synthesis of Novel 3-Substituted Coumarin-3-carboxamide

Enayatollah Sheikhhosseini, Saeed Balalaie^{†,*}, Mohammad Ali Bigdeli[‡],
Azizollah Habibi[‡], and Hamed Piri Moghaddam[§]

Department of Chemistry, Faculty of Science, Kerman Branch, Islamic Azad University, Kerman, Iran

[†]Peptide Chemistry Research Center, K. N. Toosi University of Technology, P. O. Box 15875-4416, Tehran, Iran

*E-mail: balalaie@kntu.ac.ir

[‡]Faculty of Chemistry, Kharazmi University, No. 49, Mofateh Ave. Tehran, Iran

[§]Department of Chemistry, Sharif University of Technology, Tehran, Iran

(Received October 13, 2013; Accepted January 22, 2014)

ABSTRACT. A series of novel pseudopeptides contained coumarin skeleton were synthesized through the Ugi-four-component reaction. The 3-substituted coumarin-3-carboxamides were formed through reaction of benzaldehyde derivatives, anilines, coumarin-3-carboxylic acid and isocyanides with high yields and high bond-forming efficiency at room temperature. These novel amidated coumarins exhibit brilliant fluorescence in range of 535–547 nm in chloroform.

Key words: 3-Substituted amidated coumarins, Ugi-4CR, Pseudopeptides, Fluorescence property

INTRODUCTION

The synthesis of coumarins and their derivatives has attracted considerable attention from organic and medicinal chemists for many years as a large number of natural products contain these heterocyclic molecules.¹ They are widely used as additives in food, perfumes, cosmetics, pharmaceutical,^{2,3} optical brighteners⁴ and dispersed fluorescent and laser dyes.⁵ Natural coumarins and their synthetic structural analogs have broad biological activities, such as antimicrobial and antimicrobial,⁶ antitumor,⁷ and antiviral activities.⁸ Meanwhile, some coumarin derivatives could affect human immunodeficiency virus integrase inhibitors.⁹ Some coumarins have inhibitory activity against some serine proteases and matrix metalloproteases (MMPs),¹⁰ and also act as a selective anti-proliferative agent.^{11–14}

It is shown that the existence of amide functional groups in the structure of coumarins could affect the biological activities of these compounds and it is related to the pattern of substitution and also its position in coumarin ring.¹⁵ According to these results, the number of amide bonds and also their position in the structure of coumarin scaffold could affect the type of biological activity.^{16–20}

Meanwhile, coumarins are the largest class of laser dyes and highly sensitive for the blue-green region, include commercially available fluorescent derivatives of extended spectral range, high emission quantum yield, photostability and good solubility in many solvents.^{21,22} Recently, Katritzky and co-workers reported synthesis of fluorescent coumarin-

aminoxy acid and coumarin-thioester which could be used for the detection of peptides.^{23,24}


Based on the diverse biological activities of coumarins and their fluorescence property, the synthesis of compounds which contained more amide bonds is an interesting area in organic synthesis.

Multi-component reactions (MCRs) have been investigated extensively in organic and diversity-oriented synthesis; primarily due to their ability to generate complex molecular functionality from simple starting materials via one-pot reactions.^{25,26} For this reason, MCRs are featured in many diversity-oriented projects, whose biological relevance has been validated by the discovery of novel biological probes and drug leads.²⁷ Due to importance of chromene scaffold, recently, Che et al. reported synthesis of chromeno [3,4-*c*]pyrrole-3,4-diones via sequential Ugi/intramolecular-Michael addition reactions.²⁸

In continuation of our research work to design novel multicomponent reactions,²⁹ we wish to report a simple one-pot four-component reaction of aromatic aldehydes (**1**), aniline (**2**), coumarin-3-carboxylic acid (**3**) and isocyanide (**4**) which leading to pseudopeptides which contained coumarin skeleton **5a-p** (Scheme 1).

EXPERIMENTAL

Commercially available materials were used without further purification. Melting points were determined on an *Electrothermal 9100* apparatus and were uncorrected.



Scheme 1. Synthesis of 3-substituted coumarin carboxamides.

IR spectra were obtained on an ABB FT-IR FTLA 2000 spectrometer. ^1H NMR and ^{13}C NMR spectra were run on Bruker DRX-300 AVANCE spectrometer at 300 MHz for ^1H NMR, and 75 MHz for ^{13}C NMR. CDCl_3 was used as solvent. HRMS was recorded on Mass-ESI-POS (Apex Qe-FT-ICR instrument) spectrometer. Fluorescence spectra for all compounds were collected at room temperature with a *Photon Technology international model MPI* steady-state fluorimeter. Fluorescence measurements were taken using *NSG Precision Cells (Farmingdale, NY)* ES quartz cuvettes (190–2000 nm).

Synthesis of Coumarin-3-carboxylic Acid

In a round-bottomed flask, salicylaldehyde (10 mmol) and Meldrum's acid (12 mmol) in water (20 ml) were heated at reflux under stirring for 10 h. The reaction mixture was cooled and filtered on Büchner funnel. Further purification was done using crystallization in methanol (Yield = 95%).

General Procedure for the Synthesis of Chromene-3-carboxamide Derivatives (5a–q)

Primary amine **2** (1 mmol) was added to a solution of aldehyde **1** (1 mmol) in methanol (5 mL) and the reaction mixture was stirred at room temperature for 1 h. Coumarin-3-carboxylic acid **3** (1 mmol) was added and stirring was continued for 15 min followed by addition of isocyanide **4** (1 mmol). The resulting solution was stirred at room temperature for 4–6 h. The reaction was complete after 4–6 h. The solvent was removed under reduced pressure and the product was precipitated by addition of water. Precipitated solid product was recrystallized from water/acetone and identified (Yields 77–93%).

N-((Cyclohexylcarbamoyl)(phenyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5a): Yield 90%. Mp = 233–236 °C. IR (KBr, cm^{-1}): 3262, 1717, 1649, 1641 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.15–2.08 (m, 10H, 5 CH_2), 3.93 (m, 1H, CH–NH), 6.35 (s, 1H, CH–N), 6.78 (br s, 1H, NH), 7.00–7.50 (14H, m, H–Ar), 7.80 (s, 1H, =CH). ^{13}C NMR (75 MHz, CDCl_3): δ : 24.8, 24.9, 25.4, 32.7, 33.0, 49.1, 65.7, 116.7, 117.9, 124.8, 125.9, 128.1, 128.2, 128.4, 128.5, 129.9, 130.5, 132.5, 134.0, 138.8, 142.3, 153.6, 158.3, 165.4, 167.8. HR-MS (ESI) calcd for $\text{C}_{30}\text{H}_{29}\text{N}_2\text{O}_4$ $[\text{M}+\text{H}]^+$

481.21249, found 481.21246; calcd for $\text{C}_{30}\text{H}_{28}\text{N}_2\text{NaO}_4$ $[\text{M}+\text{Na}]^+$ 503.19439, found 503.19442; calcd for $\text{C}_{30}\text{H}_{28}\text{KN}_2\text{O}_4$ $[\text{M}+\text{K}]^+$ 519.16828, found 519.16830.

N-((Cyclohexylcarbamoyl)(2-methoxyphenyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5b): Yield 87%. Mp = 132–136 °C. IR (KBr, cm^{-1}): 3293, 1726, 1656, 1607 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.21–2.15 (m, 10H, 5 CH_2), 3.75 (s, 3H, OMe), 3.97 (m, 1H, CH–NH), 6.60 (s, 1H, CH–N), 6.65 (d, 1H, J = 8.1 Hz, H–Ar), 6.73 (t, 1H, J = 7.5 Hz, H–Ar), 6.90–7.51 (m, 12H, NH and H–Ar), 7.83 (s, 1H, =CH). ^{13}C NMR (75 MHz, CDCl_3): δ : 24.9, 25.1, 25.5, 32.7, 33.2, 48.9, 55.1, 60.4, 110.0, 116.7, 118.1, 120.2, 122.4, 124.8, 126.3, 127.8, 128.0, 128.4, 129.6, 129.8, 131.4, 132.4, 1387.7, 142.1, 153.6, 157.4, 158.4, 165.3, 168.4. HR-MS (ESI) calcd for $\text{C}_{31}\text{H}_{31}\text{N}_2\text{O}_5$ $[\text{M}+\text{H}]^+$ 511.22292, found 511.22290; calcd for $\text{C}_{31}\text{H}_{30}\text{N}_2\text{NaO}_5$ $[\text{M}+\text{Na}]^+$ 533.20485, found 533.20484; calcd for $\text{C}_{31}\text{H}_{30}\text{KN}_2\text{O}_5$ $[\text{M}+\text{K}]^+$ 549.17867, found 549.17867.

N-((Cyclohexylcarbamoyl)(4-methoxyphenyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5c): Yield 92%. Mp = 202–205 °C. IR (KBr, cm^{-1}): 3271, 1720, 1656, 1609 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.16–2.06 (m, 10H, 5 CH_2), 3.71 (s, 3H, OMe), 3.92 (m, 1H, CH–NH), 6.29 (s, 1H, CH–N), 6.66 (br s, 1H, NH), 6.67 (d, J = 8.6 Hz, 2H, H–Ar), 7.07 (d, J = 8.6 Hz, 2H, H–Ar), 7.02–7.49 (m, 9H, H–Ar) 7.78 (s, 1H, =CH). ^{13}C NMR (75 MHz, CDCl_3): δ : 24.8, 24.9, 25.4, 32.7, 32.9, 49.0, 55.1, 65.0, 113.6, 116.6, 117.9, 124.7, 125.9, 126.0, 128.1, 128.4, 128.5, 130.0, 131.8, 132.5, 138.8, 142.1, 153.6, 158.2, 159.3, 165.3, 168.2. HR-MS (ESI) calcd for $\text{C}_{31}\text{H}_{31}\text{N}_2\text{O}_5$ $[\text{M}+\text{H}]^+$ 511.22277, found 511.22277; calcd for $\text{C}_{31}\text{H}_{30}\text{N}_2\text{NaO}_5$ $[\text{M}+\text{Na}]^+$ 533.20471, found 533.20471; calcd for $\text{C}_{31}\text{H}_{30}\text{KN}_2\text{O}_5$ $[\text{M}+\text{K}]^+$ 549.17862, found 549.17862.

N-((Cyclohexylcarbamoyl)(2,4-dimethoxyphenyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5d): Yield 93%. Mp = 189–192 °C. IR (KBr, cm^{-1}): 3274, 1718, 1654, 1607 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.19–2.08 (m, 10H, 5 CH_2), 3.69 (s, 6H, 2OMe), 3.94 (m, 1H, CH–NH), 6.20 (d, 1H, J = 2.4 Hz, H–Ar), 6.24–6.27 (dd, 1H, J = 7.8, 2.4 Hz, H–Ar), 6.50 (s, 1H, CH–N), 6.82 (br s, 1H, NH), 6.83–7.20 (7H, m, H–Ar), 7.40–7.50 (m, 2H, H–Ar), 7.79 (1H, s, =CH). ^{13}C NMR (75 MHz, CDCl_3): δ : 24.9, 25.1, 25.5, 32.7, 33.2, 48.9, 55.1, 55.2, 60.2, 97.9, 104.0, 114.8, 116.7, 118.1, 124.7, 126.3, 127.8, 128.0, 28.4, 129.7, 132.1, 132.3, 138.8, 141.9, 153.6, 158.5, 161.0, 165.3, 168.6. HR-MS (ESI) calcd for $\text{C}_{32}\text{H}_{33}\text{N}_2\text{O}_6$ $[\text{M}+\text{H}]^+$ 541.23498, found 541.23483; calcd for $\text{C}_{32}\text{H}_{32}\text{N}_2\text{NaO}_6$ $[\text{M}+\text{Na}]^+$ 563.21672, found 563.21698; calcd for $\text{C}_{32}\text{H}_{32}\text{KN}_2\text{O}_6$ $[\text{M}+\text{K}]^+$ 579.19082, found 579.19067.

N-((Cyclohexylcarbamoyl)(3,4,5-trimethoxyphenyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5e): Yield 88%. Mp = 168–170 °C. IR (KBr, cm^{-1}): 3270, 1718, 1653, 1606 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.17–2.05 (m, 10H, 5CH₂), 3.68 (s, 6H, 2OMe), 3.75 (s, 3H, OMe), 3.90 (m, 1H, CH–NH), 6.21 (s, 1H, CH–N), 6.41 (s, 2H, H–Ar), 6.62 (br s, 1H, NH), 7.06–7.50 (m, 9H, H–Ar), 7.77 (s, 1H, =CH). ^{13}C NMR (75 MHz, CDCl_3): δ : 24.8, 24.9, 25.4, 32.7, 32.9, 49.0, 56.0, 60.7, 65.8, 107.6, 116.7, 117.9, 124.8, 125.8, 128.3, 128.4, 128.6, 129.2, 129.9, 132.6, 137.8, 138.9, 142.1, 152.8, 153.6, 158.3, 165.4, 167.8. HR-MS (ESI) calcd for $\text{C}_{33}\text{H}_{35}\text{N}_2\text{O}_7$ [M+H]⁺ 571.24557, found 571.24542; calcd for $\text{C}_{33}\text{H}_{34}\text{N}_2\text{NaO}_7$ [M+Na]⁺ 593.22723, found 593.22710.

N-((Cyclohexylcarbamoyl)(p-tolyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5f): Yield 83%. Mp = 220–222 °C. IR (KBr, cm^{-1}): 3314, 1731, 1655 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.15–2.03 (m, 10H, 5CH₂), 2.24 (s, 3H, Me), 3.92 (m, 1H, CH–NH), 6.27 (s, 1H, CH–N), 6.59 (br s, 1H, NH), 6.90–7.25 (m, 11H, H–Ar), 7.41 (d, 1H, J = 7.5 Hz, H–Ar), 7.47 (t, 1H, J = 7.5 Hz, H–Ar), 7.78 (s, 1H, =CH). ^{13}C NMR (75 MHz, CDCl_3): δ : 21.1, 24.9, 25.0, 25.5, 32.8, 33.0, 49.0, 65.6, 116.7, 118.0, 124.8, 128.1, 128.2, 128.4, 128.5, 129.0, 130.0, 130.4, 130.9, 132.5, 138.1, 139.0, 142.2, 153.6, 165.8, 168.1. HR-MS (ESI) calcd for $\text{C}_{31}\text{H}_{31}\text{N}_2\text{O}_4$ [M+H]⁺ 495.22857, found 495.22850; calcd for $\text{C}_{31}\text{H}_{30}\text{N}_2\text{NaO}_4$ [M+Na]⁺ 517.21052, found 517.21045; calcd for $\text{C}_{31}\text{H}_{30}\text{KN}_2\text{O}_4$ [M+K]⁺ 533.18455, found 533.18447.

N-((Cyclohexylcarbamoyl)(4-isopropyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5g): Yield 91%. Mp = 164–166 °C. IR (KBr, cm^{-1}): 33.18, 1730, 1653 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.15 (d, J = 6.9 Hz, 6H, 2Me), 1.19–2.07 (m, 10H, 5CH₂), 2.79 (m, 1H, J = 6.90 Hz, CH(Me)₂), 3.93 (m, 1H, CH–NH), 6.27 (s, 1H, CH–N), 6.64 (br s, 1H, NH), 7.01–7.50 (m, 13H, H–Ar) 7.78 (s, 1H, =CH). ^{13}C NMR (75 MHz, CDCl_3): δ : 23.7, 24.9, 25.0, 25.5, 32.7, 32.9, 33.6, 49.0, 65.7, 116.7, 118.0, 124.7, 126.0, 126.3, 127.9, 128.4, 128.5, 129.9, 130.4, 131.2, 132.4, 139.0, 142.1, 149.0, 153.6, 158.2, 165.3, 168.1. HR-MS (ESI) calcd for $\text{C}_{33}\text{H}_{35}\text{N}_2\text{O}_4$ [M+H]⁺ 523.25906, found 523.25907; calcd for $\text{C}_{33}\text{H}_{34}\text{N}_2\text{NaO}_4$ [M+Na]⁺ 545.24092, found 545.24093; calcd for $\text{C}_{33}\text{H}_{34}\text{KN}_2\text{O}_4$ [M+K]⁺ 561.21483, found 561.21485.

N-((Cyclohexylcarbamoyl)(2,4,6-trimethylphenyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5h): Yield 90%. Mp = 251–254 °C. IR (KBr, cm^{-1}): 3275, 1721, 1652 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.08–2.04 (m, 10H, 5CH₂), 2.21 (s, 9H, 3Me), 3.86 (m, 1H, CH–NH), 5.86 (br s, 1H, NH), 6.21 (s, 1H, CH–N), 6.81–7.78 (m, 10H, H–Ar), 7.80 (s, 2H, =CH and H–Ar). ^{13}C NMR (75 MHz, CDCl_3): δ : 20.3, 20.8, 25.0, 25.1, 25.4, 32.9, 33.2, 49.1, 59.0, 116.6,

118.0, 124.6, 126.6, 127.9, 128.3, 129.7, 131.1, 131.8, 132.1, 137.5, 138.4, 139.8, 141.8, 153.6, 157.8, 165.5, 169.0. HR-MS (ESI) calcd for $\text{C}_{33}\text{H}_{35}\text{N}_2\text{O}_4$ [M+H]⁺ 523.25911, found 523.25911; calcd for $\text{C}_{33}\text{H}_{34}\text{N}_2\text{NaO}_4$ [M+Na]⁺ 545.24100, found 545.24101; calcd for $\text{C}_{33}\text{H}_{34}\text{KN}_2\text{O}_4$ [M+K]⁺ 561.21492, found 561.21493.

N-((Cyclohexylcarbamoyl)(2-nitrophenyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5i): Yield 75%. Mp = 78–81 °C. IR (KBr, cm^{-1}): 3331, 1707, 1654, 1607 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.29–2.10 (m, 10H, 5CH₂), 3.94 (m, 1H, CH–NH), 7.00–7.90 (m, 15H, CH–N, NH and H–Ar), 8.0 (s, 1H, =CH). ^{13}C NMR (75 MHz, CDCl_3): δ : 24.9, 25.0, 25.4, 32.5, 33.1, 49.3, 61.7, 116.7, 117.9, 124.6, 125.1, 125.6, 128.3, 128.6, 128.7, 129.2, 129.3, 132.7, 132.8, 133.0, 138.1, 143.2, 149.6, 153.6, 158.7, 165.2, 166.9. HR-MS (ESI) calcd for $\text{C}_{30}\text{H}_{28}\text{N}_3\text{O}_6$ [M+H]⁺ 526.19762, found 526.19759; calcd for $\text{C}_{30}\text{H}_{27}\text{N}_3\text{NaO}_6$ [M+Na]⁺ 548.17946, found 548.17944; calcd for $\text{C}_{30}\text{H}_{27}\text{KN}_3\text{O}_6$ [M+K]⁺ 564.15325, found 564.15324.

N-((Cyclohexylcarbamoyl)(3-nitrophenyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5j): Yield 80%. Mp = 212–215 °C. IR (KBr, cm^{-1}): 3282, 1739, 1719, 1654 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.35–2.14 (m, 10H, 5CH₂), 3.97 (m, 1H, CH–NH), 6.51 (1H, s, CH–N), 7.07–7.64 (m, 12H, NH and H–Ar), 7.91 (s, 1H, =CH), 8.05 (dd, 1H, J = 8.1 Hz, H–Ar), 8.10 (d, 1H, J = 1.0 Hz, H–Ar). ^{13}C NMR (75 MHz, CDCl_3): δ : 24.9, 25.0, 25.4, 32.7, 33.1, 49.4, 64.7, 116.8, 117.8, 123.0, 125.1, 125.4, 125.7, 128.6, 128.7, 128.9, 129.1, 129.4, 133.1, 136.4, 136.8, 138.2, 143.2, 147.8, 153.7, 158.6, 165.6, 166.5. HR-MS (ESI) calcd for $\text{C}_{30}\text{H}_{28}\text{N}_3\text{O}_6$ [M+H]⁺ 526.19725, found 526.19725; calcd for $\text{C}_{30}\text{H}_{27}\text{N}_3\text{NaO}_6$ [M+Na]⁺ 548.17919, found 548.17919; calcd for $\text{C}_{30}\text{H}_{27}\text{KN}_3\text{O}_6$ [M+K]⁺ 564.15305, found 564.15306.

N-((Cyclohexylcarbamoyl)(4-nitrophenyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5k): Yield 84%. Mp = 225–227 °C. IR (KBr cm^{-1}): 3269, 1717, 1655, 1646 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.03–1.70 (m, 10H, 5CH₂), 3.59 (m, 1H, CH–NH), 6.31 (1H, s, CH–N), 7.02 (br s, 1H, NH), 7.04 (d, 2H, J = 8.1 Hz, H–Ar), 7.07 (d, 1H, J = 8.5 Hz, H–Ar), 7.18–7.33 (m, 4H, H–Ar), 7.48 (d, 2H, J = 8.1 Hz, H–Ar), 7.57 (t, 1H, J = 7.5 Hz, H–Ar), 7.70 (d, 1H, J = 8.5 Hz, H–Ar), 8.09 (s, 1H, =CH), 8.25 (d, 1H, J = 7.5 Hz, H–Ar). ^{13}C NMR (75 MHz, CDCl_3): δ : 24.4, 25.1, 30.7, 32.1, 48.1, 63.2, 116.2, 117.6, 123.0, 124.9, 125.0, 128.2, 129.1, 130.4, 131.2, 132.9, 138.5, 142.4, 142.8, 146.8, 152.9, 158.3, 164.5, 166.7 δ $\text{C}_{30}\text{H}_{27}\text{N}_3\text{O}_6$ (525.19): C, 68.56; H, 5.18; N, 8.00. Found: C, 68.36; H, 5.27; N, 7.87.

N-((Cyclohexylcarbamoyl)(2-(prop-2-ynyloxy)phenyl)methyl)-2-oxo-N-phenyl-2H-chromone-3-carboxamide (5l):

Yield 85%. Mp = 195–196 °C. IR (KBr cm^{-1}): 3300, 2124, 1726, 1652 cm^{-1} . ^1H NMR (300 MHz, acetone): δ = 1.20–1.75 (m, 10H, 5 CH_2 , cyclohexyl), 3.19 (t, 1H, J = 2.3 Hz, CH), 3.86 (m, 1H, CH–NH), 4.81 (d, 2H, J = 2.6 Hz, CH_2 –O), 6.55 (s, 1H, CH–N), 6.70 (t, 1H, J = 7.4 Hz, H–Ar), 6.90–6.97 (m, 4H, NH and H–Ar), 7.08–7.13 (m, 2H, H–Ar), 7.21–7.33 (m, 4H, H–Ar), 7.54–7.69 (m, 3H, H–Ar), 7.93 (s, 1H, =CH). ^{13}C NMR (75 MHz, acetone) δ : 25.6, 25.7, 26.3, 33.4, 33.6, 49.5, 56.4, 60.2, 77.4, 79.7, 112.2, 117.1, 119.1, 121.3, 124.5, 125.6, 128.3, 128.6, 129.7, 130.2, 131.1, 132.0, 133.3, 140.0, 142.59, 156.6, 169.0. HR-MS (ESI) calcd for $\text{C}_{33}\text{H}_{31}\text{N}_2\text{O}_5$ [$\text{M}+\text{H}$] $^+$ 535.22320, found 535.22316; calcd for $\text{C}_{33}\text{H}_{30}\text{N}_2\text{NaO}_5$ [$\text{M}+\text{Na}$] $^+$ 557.20517, found 557.20510; calcd for $\text{C}_{33}\text{H}_{30}\text{KN}_2\text{O}_5$ [$\text{M}+\text{K}$] $^+$ 573.17909, found 573.17901.

N-((tert-Butylcarbamoyl)(2-(prop-2-ynyloxy)phenyl)methyl)-2-oxo-N-phenyl-2H-chromene-3-carboxamide (5m): Yield 88%. Mp = 130–133 °C. IR (KBr cm^{-1}): 3397, 2113, 1713, 1668, 1650 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.46 (s, 9H, *t*-but), 2.51 (t, 1H, J = 2.2 Hz, CH), 4.60 (d, 2H, J = 2.2 Hz, CH_2 –O), 6.47 (s, 1H, CH–N), 6.57 (br s, 1H, NH), 6.78–6.83 (m, 2H, H–Ar), 6.96–6.97 (m, 3H, H–Ar), 7.12–7.26 (m, 6H, H–Ar), 7.39–7.48 (m, 2H, H–Ar), 7.77 (s, 1H, =CH). ^{13}C NMR (75 MHz, CDCl_3) δ : 28.7, 51.8, 55.8, 60.8, 111.4, 116.7, 118.1, 121.2, 123.1, 124.7, 126.2, 127.8, 128.1, 128.4, 129.6, 131.4, 132.3, 139.0, 141.1, 153.6, 155.4, 158.2, 165.3, 168.4. HR-MS (ESI) calcd for $\text{C}_{31}\text{H}_{29}\text{N}_2\text{O}_5$ [$\text{M}+\text{H}$] $^+$ 509.20763, found 509.20758; calcd for $\text{C}_{31}\text{H}_{28}\text{N}_2\text{NaO}_5$ [$\text{M}+\text{Na}$] $^+$ 531.18953, found 531.18949; calcd for $\text{C}_{31}\text{H}_{28}\text{KN}_2\text{O}_5$ [$\text{M}+\text{K}$] $^+$ 547.16356, found 547.16351.

N-((Cyclohexylcarbamoyl)(2-(prop-2-ynyloxy)phenyl)methyl)-N-(4-chlorophenyl)-2-oxo-2H-chromene-3-carboxamide (5n): Yield 85%. Mp = 195–196 °C. IR (KBr cm^{-1}): 3338, 2123, 1733, 1674, 1633 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.15–2.10 (m, 10H, 5 CH_2 , cyclohexyl), 2.53 (s, 1H, CH), 3.93 (m, 1H, CH–NH), 4.67 (s, 2H, CH_2 –O), 6.59 (s, 1H, CH–N), 6.77–6.90 (m, 5H, NH and H–Ar), 7.08–7.28 (m, 6H, H–Ar), 7.43–7.53 (m, 2H, H–Ar), 7.84 (s, 1H, =CH). ^{13}C NMR (75 MHz, CDCl_3) δ : 24.9, 25.0, 25.4, 32.7, 33.1, 49.0, 55.8, 60.1, 111.5, 116.8, 118.0, 121.3, 122.6, 124.9, 125.9, 128.3, 128.5, 131.1, 131.5, 132.7, 133.7, 137.2, 142.5, 153.7, 155.4, 158.3, 165.1, 168.1. HR-MS (ESI) calcd for $\text{C}_{33}\text{H}_{30}\text{ClN}_2\text{O}_5$ [$\text{M}+\text{H}$] $^+$ 569.18441, found 569.18435; calcd for $\text{C}_{33}\text{H}_{29}\text{ClN}_2\text{NaO}_5$ [$\text{M}+\text{Na}$] $^+$ 591.16612, found 591.16608; calcd for $\text{C}_{33}\text{H}_{29}\text{ClKN}_2\text{O}_5$ [$\text{M}+\text{K}$] $^+$ 607.14036, found 607.14030.

N-((tert-Butylcarbamoyl)(2-(prop-2-ynyloxy)phenyl)methyl)-N-(4-chlorophenyl)-2-oxo-2H-chromene-3-carboxamide (5o): Yield 78%. Mp = 97–100 °C. IR (KBr cm^{-1}):

3359, 3298, 2119, 1723, 1654, 1607 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.43 (s, 9H, *t*-but), 2.53 (t, 1H, J = 2.3 Hz, CH), 4.67 (d, 2H, J = 2.3 Hz, CH_2 –O), 6.35 (br s, 1H, NH), 6.44 (s, 1H, CH–N), 6.79–6.85 (m, 2H, H–Ar), 6.93 (d, 2H, J = 8.6 Hz, H–Ar), 7.11–7.26 (m, 6H, H–Ar), 7.42 (dd, 1H, J = 7.2, 1.4 Hz, H–Ar), 7.49 (dt, 1H, J = 7.8, 1.4 Hz, H–Ar), 7.78 (s, 1H, =CH). ^{13}C NMR (75 MHz, CDCl_3) δ : 28.7, 51.9, 55.7, 60.5, 111.6, 116.7, 118.0, 121.3, 122.8, 124.8, 125.9, 128.3, 128.4, 130.0, 131.1, 131.3, 132.5, 133.6, 137.4, 1342.2, 153.7, 155.3, 158.0, 165.1, 168.3. HR-MS (ESI) calcd for $\text{C}_{31}\text{H}_{28}\text{ClN}_2\text{O}_5$ [$\text{M}+\text{H}$] $^+$ 543.16869, found 543.16864; calcd for $\text{C}_{31}\text{H}_{27}\text{ClN}_2\text{NaO}_5$ [$\text{M}+\text{Na}$] $^+$ 565.15050, found 565.15046; calcd for $\text{C}_{31}\text{H}_{27}\text{ClKN}_2\text{O}_5$ [$\text{M}+\text{K}$] $^+$ 581.12444, found 581.12440.

N-((Cyclohexylcarbamoyl)(2-(prop-2-ynyloxy)phenyl)methyl)-N-(2,4-dichlorophenyl)-2-oxo-2H-chromene-3-carboxamide (5p): Yield 75%. Mp = 177–180 °C. IR (KBr cm^{-1}): 3338, 2123, 1733, 1674, 1633 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ = 1.28–2.16 (m, 10H, 5 CH_2 , cyclohexyl), 2.61 (t, 1H, J = 2.3 Hz, CH), 3.95 (m, 1H, CH–NH), 4.84 (d, 2H, J = 2.3 Hz, CH_2 –O), 6.87 (s, 1H, CH–N), 6.99 (d, 1H, J = 2.4 Hz, NH), 7.07 (m, 5H, H–Ar), 7.18–7.27 (m, 3H, H–Ar), 7.46–7.53 (m, 3H, H–Ar), 7.89 (s, 1H, =CH). ^{13}C NMR (75 MHz, CDCl_3) δ : 24.9, 52.0, 25.4, 32.7, 33.1, 49.2, 59.8, 60.9, 116.8, 117.9, 125.0, 125.8, 128.0, 128.5, 128.6, 128.7, 129.5, 129.6, 130.2, 130.5, 132.1, 132.8, 138.1, 142.6, 151.1, 153.6, 158.6, 165.2, 166.9. HR-MS (ESI) calcd for $\text{C}_{33}\text{H}_{29}\text{Cl}_2\text{N}_2\text{O}_5$ [$\text{M}+\text{H}$] $^+$ 603.14562, found 603.14555; calcd for $\text{C}_{33}\text{H}_{28}\text{Cl}_2\text{N}_2\text{NaO}_5$ [$\text{M}+\text{Na}$] $^+$ 625.12762, found 625.12754; calcd for $\text{C}_{33}\text{H}_{28}\text{Cl}_2\text{KN}_2\text{O}_5$ [$\text{M}+\text{K}$] $^+$ 641.10149, found 641.10142.

RESULTS AND DISCUSSION

Coumarin-3-carboxylic acid is an efficient starting material for the synthesis of functionalized coumarin skeleton. It appears in a vast range of natural products with high levels of biological activity and also was used for the synthesis of compounds with pharmaceutical activity.^{30–32} Coumarin-3-carboxylic acid **3** was obtained from the reaction of salicylaldehyde with Meldrum's acid refluxing in water.³³ Initially, the four-component reaction of benzaldehyde **1a**, aniline **2**, coumarin-3-carboxylic acid **3**, and cyclohexyl isocyanides **4a** were carried out smoothly in methanol at room temperature was selected as model reaction (*Scheme 1*). The desired product **5a** was isolated in 90% yield (*Table 1*, entry 1). Encouraged by this result, we turned our attention to other substituted benzaldehydes, amines and isocyanides. The desired amidated coumarins **5b–p** was obtained in 75–92%.

Table 1. Synthesis of amidated 3-substituted coumarins using Ugi 4-CR

Entry	Ar	R ¹	R ²	Product	Yield (%)*
1	Ph	Ph	Cyc**	5a	90
2	2-MeOC ₆ H ₄ -	Ph	Cyc	5b	87
3	4-MeOC ₆ H ₄ -	Ph	Cyc	5c	92
4	2,4-(MeO) ₂ C ₆ H ₃ -	Ph	Cyc	5d	93
5	4,5,6-(MeO) ₃ C ₆ H ₂ -	Ph	Cyc	5e	88
6	4-Me-C ₆ H ₄ -	Ph	Cyc	5f	83
7	4- <i>i</i> -pr-C ₆ H ₄ -	Ph	Cyc	5g	91
8	2,4,6-(Me) ₃ C ₆ H ₂ -	Ph	Cyc	5h	90
9	2-O ₂ N-C ₆ H ₄ -	Ph	Cyc	5i	75
10	3-O ₂ N-C ₆ H ₄ -	Ph	Cyc	5j	80
11	4-O ₂ N-C ₆ H ₄ -	Ph	Cyc	5k	77
12	2-(prop-2-ynyloxy)-C ₆ H ₄ -	Ph	Cyc	5l	85
14	2-(prop-2-ynyloxy)-C ₆ H ₄ -	Ph	<i>t</i> -butyl***	5m	85
15	2-(prop-2-ynyloxy)-C ₆ H ₄ -	4-Cl-C ₆ H ₄ -	Cyc	5n	88
16	2-(prop-2-ynyloxy)-C ₆ H ₄ -	4-Cl-C ₆ H ₄ -	<i>t</i> -butyl	5o	78
17	2-(prop-2-ynyloxy)-C ₆ H ₄ -	2,4-(Cl) ₂ C ₆ H ₃ -	Cyc	5p	75


*Yields of isolated products

**Cyclohexyl isocyanide

****tert*-butyl isocyanide

It was satisfying to observe that the reaction was complete in 6 hours and in many cases the reaction products precipitate during the reaction and the products were obtained after recrystallization. This approach was used for the synthesis of a chemical library of coumarins. It was found that these conditions were compatible with a range of electron-withdrawing and electron-donating substituents. The conditions were sufficiently mild to be tolerated by a number of functionalities including nitro and methoxy groups. Reaction with 2,4-dimethoxybenzaldehyde led to the highest yield (93% yields, Table 1, entry 4).

The structures of compounds **5a–p** were deduced from their spectroscopic data and also high-resolution mass spectrometry data (HR-ESI-MS). The ¹H NMR spectrum of the products showed a distinguished peak at region δ 6.21–6.60 ppm for the –CH protons and also a broad singlet signal at δ 5.86–7.02 ppm for the –NH moiety. Partial assignments of these resonances are given in the experimental section. The ¹³C NMR spectrum revealed two distinct peaks at δ(C) 157–165 ppm for the amide C=O groups, and 166–169 ppm for the carbonyl of lactone in coumarin skeleton. *O*-Propargyloxy benzaldehyde was used as starting material for the synthesis of products **5l–p**. We believe that this new functionality based multicomponent reactions have a huge potential in creating new molecules or simplifying the synthesis of existing compounds and the synthesized products have good potential for further reactions. The products **5l–p** could be used as starting material for some reac-

**Scheme 2.** Proposed mechanism for the synthesis of amidated 3-substituted coumarins **5a–p**.

tions such as: Click chemistry, Pauson-Khand reaction, and cyclization reaction.

According to the commonly accepted Ugi-4CR proposed mechanism, it seems that amine, the carbonyl compound and the acid are in equilibrium with the iminiumcarboxylate **6** in the reaction medium. The α -addition of the iminium carboxylate onto the carbenoid carbon of the isocyanide leads to the formation of the primary four-component adduct **7**, which undergoes an intramolecular acylation known as Mumm rearrangement to give the stable Ugi adduct **5** (Scheme 2).

Since some of coumarin derivatives could be used as fluorophores,^{34,35} we were interested to investigate the flu-




Figure 1. Emission spectra of compounds **5b**, **5d**, **5e**, **5g**, **5p** in chloroform.




Figure 2. Emission spectra of compounds **5b**, **5d**, **5e**, **5g**, **5p** in acetonitrile.

orescence property of the synthesized compounds. The fluorescence property of some novel amidated coumarins derivatives were investigated in chloroform, acetonitrile, and dimethyl formamide. The wavelengths of the absorption maxima and fluorescence emission maxima fluorescence property of compounds **5b**, **5d**, **5e**, **5g**, **5p** were investigated. Excitations were carried out with the four different wavelengths 200, 210, 220, and 230 nm, and the emission was investigated in the range 300–700 nm. In the case of **5d**, the maximum emission occurred at the excitation wavelength 200 nm, and the maximum emission occurred at the wavelength 542, 546, and 543 nm, respectively (Figs. 1–3). All of compounds displayed almost similar fluorescence emission in the range of 535–547 nm, and they could show a significant difference to coumarin-3-carboxylic acid emission spectra.




Figure 3. Emission spectra of compounds **5b**, **5d**, **5e**, **5g**, **5p** in dimethyl formamide.

CONCLUSION

In conclusion, we have synthesized a small library of 3-substituted coumarin carboxamides or pseudo-peptides using Ugi-4CR of coumarin-3-carboxylic acid, benzaldehydes, anilines and isocyanides. Some of these novel derivatives could be used for further reactions and have fluorescent spectra in the range 535–547 nm. The simplicity of the synthetic protocol and availability of diverse starting materials make this an attractive strategy for obtaining new coumarins via combinatorial techniques. Investigations of biological activity of these compounds are in progress.

Acknowledgments. Publication cost of this paper was supported by the Korean Chemical Society.

REFERENCES

- Murray, R. D. H.; Mendez, J.; Brown, S. A. *The Natural Coumarins: Occurrence, Chemistry and Biochemistry*; John Wiley: Chichester, U. K., 1982.
- Keating, G. J.; ÓKennedy, R. In *Coumarins: Biology, Applications and Mode of Action*; O'Kennedy, R., Thornes, R. D., Eds.; John Wiley and Sons: New York, 1997; pp 23–66.
- Murray, R. D. H. *Natu. Prod. Rep.* **1989**, 591.
- Zabradnik, M. *The Production and Application of Fluorescent Brightening Agents*; John Wiley and Sons: New York, 1992.
- Maeda, M. *Laser Dyes*; Academic Press: New York, 1994.
- Maxwell, A. *Mol. Microbiol.* **1993**, 9, 681.
- Dexeus, F. H.; Logothetis, C. J.; Sella, A.; Fitz, K.; Amato, R.; Reuben, J. M.; Dozier, N. *J. Clin. Oncol.* **1990**, 8, 325.
- Zembower, D. E.; Liao, S. Y.; Flavin, M. T.; Xu, Z. Q.; Stup, T. L.; Buckheit, R. W. *J. Med. Chem.* **1997**, 40, 1005.
- Yu, D.; Suzuki, M.; Xie, L.; Morris-Natschke, S. L.; Lee,

- K. H. *Med. Res. Rev.* **2003**, *23*, 322.
10. Kempen, I.; Papapostolou, D.; Thierry, N.; Pochet, L.; Counerotte, S.; Masereel, B.; Foidart, J.-M.; Reboud-Ravaux, M.; Noel, A.; Pirotte, B. *Br. J. Cancer* **2003**, *88*, 1111.
11. Finn, G.; Creaven, B.; Egan, D. *Eur. J. Pharmacol.* **2003**, *481*, 159.
12. Finn, G. J.; Creaven, B. S.; Egan, D. A. *Cancer Lett.* **2004**, *205*, 69.
13. Morita, H.; Dota, T.; Kobayashi, J. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 3665.
14. Kontogiorgis, C. A.; Hadjipavlou-Litina, D. *J. Enzyme Inhib. Med. Chem.* **2003**, *18*, 63.
15. Donnelly, A. C.; Mays, J. R.; Burlison, J. A.; Nelson, J. T.; Vielhauer, G.; Holzbeierlein, J.; Blagg, B. S. J. *J. Org. Chem.* **2008**, *73*, 8901.
16. Chimenti, F.; et al. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 4922.
17. Bylov, I. E.; Vasylyev, M. V.; Bilokin, Y. V. *Eur. J. Med. Chem.* **1999**, *34*, 997.
18. Kempen, I.; et al. *Eur. J. Med. Chem.* **2008**, *43*, 2735.
19. Reddy, N. S.; Gumireddy, K.; Mallireddigari, M. R.; Cosenza, S. C.; Venkatapuram, P.; Bell, S. C.; Reddy, E. P.; Reddy, M. V. R. *Bioorg. Med. Chem.* **2005**, *13*, 3141.
20. Melagraki, G.; Afantitis, A.; Igglessi-Markopoulou, O.; Detsi, A.; Koufaki, M.; Kontogiorgis, C.; Hadjipavlou-Litina, D. *J. Eur. J. Med. Chem.* **2009**, *44*, 3020.
21. Jung, H. S.; Ko, K. C.; Kim, G.-H.; Lee, A.-R.; Na, Y.-C.; Kang, C.; Lee, J. Y.; Kim, J. S. *Org. Lett.* **2011**, *13*, 1498.
22. Yu, T.; Zhang, P.; Zhao, Y.; Zhang, H.; Meng, J.; Fan, D. *Org. Elect.* **2009**, *10*, 653.
23. Katritzky, A. R.; Abdelmajeid, A.; Tala, S. R.; Amine, M. S.; Steel, P. J. *Synthesis* **2011**, 83.
24. Katritzky, A. R.; Ibrahim, T. S.; Tala, S. R.; Abo-Dya, N. E.; Abdel-Samii, Z. K.; El-Feky, S. A. *Synthesis* **2011**, 1494.
25. (a) Zhu, J.; Bienayme, H. Eds. *Multicomponent Reactions*; Wiley-VCH: Weinheim, 2005. (b) Tietze, L. F.; Brasche, G.; Gericke, K. M. *Domino Reactions in Organic Synthesis*; Wiley-VCH: Weinheim, 2006. (c) Orru, R. V. A.; Greef, M. *Synthesis*, **2003**, 1471. (d) Bienayme, H.; Oddon, C.; Schmidt, P. *Chem. Eur. J.* **2000**, *6*, 3321. (e) Schreiber, S. L. *Science* **2000**, *287*, 1964.
26. (a) Dömling, A.; Ugi, I. *Angew. Chem. Int. Ed.* **2000**, *39*, 3168. (b) Sunderhaus, J. D.; Martin, S. F. *Chem. Eur. J.* **2009**, *15*, 1300. (c) Isambert, N.; Lavilla, R. *Chem. Eur. J.* **2008**, *14*, 8444. (d) Ruijter, E.; Scheffelaar, R.; Orru, R. V. A. *Angew. Chem. Int. Ed.* **2011**, *50*, 6234.
27. (a) Dömling, A. *Chem. Rev.* **2006**, *106*, 17. (b) Tietze, L. F.; Brasche, G.; Gericke, K. M. *In Domino Reactions in Organic Synthesis*; Wiley-VCH, 2006. (c) Dömling, A.; Wang, W.; Wang, K. *Chem. Rev.* **2012**, *112*, 3083.
28. Che, C.; Li, S.; Jiang, X.; Quan, J.; Lin, S.; Yang, Z. *Org. Lett.* **2010**, *12*, 4682.
29. (a) Bararjanian, M.; Balalaie, S.; Rominger, F.; Movassagh, B.; Bijanzadeh, H. R. *J. Org. Chem.* **2010**, *75*, 2806. (b) Bararjanian, M.; Hosseinzadeh, S.; Balalaie, S.; Bijanzadeh, H. R. *Tetrahedron* **2011**, *67*, 2644. (c) Bararjanian, M.; Balalaie, S.; Rominger, F.; Barouti, S. A. *Helv. Chim. Acta.* **2010**, *93*, 777. (d) Khoshkholgh, M. J.; Lotfi, M.; Balalaie, S.; Rominger, F. *Tetrahedron* **2009**, *65*, 4228. (e) Bararjanian, M.; Hosseinzadeh, S.; Balalaie, S.; Bijanzadeh, H. R.; Wolf, E. *Tetrahedron Lett.* **2011**, *52*, 3329. (f) Khoshkholgh, M. J.; Balalaie, S.; Gleiter, R.; Rominger, F. *Tetrahedron* **2008**, *64*, 10924. (g) Ghabraie, E.; Balalaie, S.; Bararjanian, M.; Rominger, F.; Bijanzadeh, H. R. *Tetrahedron* **2011**, *67*, 5415.
30. Bhalla, M.; Hitkari, A.; Gujrati, V. R.; Bhalla, T. N.; Shanker, K. *Eur. J. Med. Chem.* **1994**, *29*, 713.
31. Bosignore, L.; Cottiglia, F.; Lavagna, S. M.; Loy, G.; Secci, D. *Eur. J. Med. Chem.* **1994**, *29*, 479.
32. Bosignore, L.; Cottiglia, F.; Lavagna, S. M.; Loy, G.; Secci, D. *Heterocycles* **1999**, *50*, 469.
33. Maggi, R.; Bigi, F.; Carloni, S.; Mazzacani, A.; Sartori, G. *Green. Chem.* **2001**, *3*, 173.
34. Key, J. A.; Koh, S.; Timerghazin, Q. K.; Brown, A.; Cairo, C. W. *Dyes & Pigments* **2009**, *82*, 196.
35. Zhao, Y. R.; Zheng, Q.; Dakin, K.; Xu, K.; Martinez, M. L.; Li, W.-H. *J. Am. Chem. Soc.* **2004**, *126*, 4653.