

A NOTE ON A REGULARIZED GAP FUNCTION OF QVI IN BANACH SPACES

SANGHO KUM*

ABSTRACT. Recently, Taji [7] and Harms et al. [4] studied the regularized gap function of QVI analogous to that of VI by Fukushima [2]. Discussions are made in a finite dimensional Euclidean space. In this note, an infinite dimensional generalization is considered in the framework of a reflexive Banach space. To do so, we introduce an extended quasi-variational inequality problem (in short, EQVI) and a generalized regularized gap function of EQVI. Then we investigate some basic properties of it. Our results may be regarded as an infinite dimensional extension of corresponding results due to Taji [7].

1. Introduction

Given a function $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and a multifunction $S : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ with closed convex values Sx for all $x \in \mathbb{R}^n$, the quasi-variational inequality problem (in short, QVI) is to find a vector $\bar{x} \in S\bar{x}$ such that

$$\langle F\bar{x}, y - \bar{x} \rangle \geq 0 \quad \text{for all } y \in S\bar{x}, \quad (1.1)$$

where $\langle \cdot, \cdot \rangle$ denote the inner product in \mathbb{R}^n . If C is a convex closed subset of \mathbb{R}^n and for all $x \in \mathbb{R}^n$, $Sx = C$, then QVI reduces to the standard variational inequality problem (VI). It is well-known that VI can be casted as the following equivalent problem:

$$\text{minimize } \phi(x) \quad \text{subject to } x \in C$$

where $\phi(x) = -\min_{y \in C} \langle Fx, y - x \rangle + \frac{1}{2} \|y - x\|^2$ is the regularized gap function by Fukushima [2]. One of the most important features, besides some excellent global properties, of the regularized gap function ϕ is

Received February 14, 2014; Accepted April 16, 2014.

2010 Mathematics Subject Classification: Primary 49J40, 47J20, 49J50, 52A41.

Key words and phrases: Quasi-variational inequalities, gap functions, subdifferential, convex function.

This work was supported by the research grant of the Chungbuk National University in 2012.

that it is continuously differentiable if F is so. At this point, a natural question arises: Is this approach still available for QVI? The answer is affirmative since Giannessi [3] proposed a gap function for QVI as follows:

$$g(x) := - \inf_{y \in Sx} \langle Fx, y - x \rangle.$$

But his original gap function is nondifferentiable, possibly extended-valued. So it is not suitable for dealing with optimization problems. To resolve this drawback, an extension of the regularized gap function by Fukushima [2] to QVIs is provided by Taji [7] as follows.

$$g_\alpha(x) := - \min_{y \in Sx} \langle Fx, y - x \rangle + \frac{\alpha}{2} \|y - x\|^2$$

where $\alpha > 0$ is a parameter. Taji [7] verified characteristic properties of the regularized gap function g_α . However, being different from the case of VI, g_α has no nice differentiability as ϕ does. So, recently, Harms et al. [4] exploited Taji's approach to take a closer look at the nondifferentiable points so that they could specialize the results to generalized Nash equilibrium problems. Of course, in [7, 4] only finite dimensional cases are treated.

Motivated by this observation, in this note, we consider the following question from a theoretical view point: What happens in an infinite dimensional case? We introduce an extended quasi-variational inequality problem (in short, EQVI) and a generalized regularized gap function of EQVI. Then we investigate some basic properties of it in a reflexive Banach space. Our results may be regarded as an infinite dimensional extension of corresponding results due to Taji [7].

2. Preliminaries

Consider the following extended quasi-variational inequality EQVI.

EQVI : Let E be a real Banach space and E^* be its dual space. Given a function $F : E \rightarrow E^*$, a multifunction $S : E \rightrightarrows E$ with closed convex values Sx for all $x \in E$ (for the simplicity of argument, Sx is assumed to be nonempty), and a convex and continuous function $f : E \rightarrow \mathbb{R}$, find $\bar{x} \in S\bar{x}$ such that

$$\langle F\bar{x}, x - \bar{x} \rangle \geq f(\bar{x}) - f(x) \quad \text{for all } x \in S\bar{x}, \quad (2.1)$$

where $\langle \cdot, \cdot \rangle$ denote the dual pairing on $E \times E^*$. If f is the zero function, EQVI (2.1) becomes QVI. The notion EQVI is adopted from Kum and Lee [6] (originally, Chen et al. [1]).

Throughout this paper, we make the following assumptions.

ASSUMPTION 2.1. Let $\Omega : E \times E \rightarrow \mathbb{R}$ be nonnegative and for each $x \in E$, $\Omega(x, \cdot)$ be continuously Gâteaux differentiable (not necessarily convex) on E . Assume that $\Omega(x, y) = 0$ if and only if $x = y$ and $\nabla_y \Omega(x, x) = 0$ for any $x \in E$.

Our concern is to show that the function $\phi : E \rightarrow \mathbb{R}$ is a continuous gap function for EQVI (2.1) under suitable conditions;

$$\phi(x) := - \inf_{y \in Sx} \{ \langle F(x), y - x \rangle + f(y) - f(x) + \Omega(x, y) \}. \tag{2.2}$$

Before going to the main result, it is necessary to recall continuity notions of multifunctions in Harms et al. [4, Definition 3.1].

DEFINITION 2.2. Let $X \subseteq E_1$, $Y \subseteq E_2$, and $T : X \rightrightarrows Y$ be a multifunction where E_1, E_2 are Banach spaces. Then T is called

- (i) lower semicontinuous at $\bar{x} \in X$ if for all sequences $\{x^k\} \subseteq X$ with $x^k \rightarrow \bar{x}$ and all $\bar{y} \in T\bar{x}$, there exists $k_0 \in \mathbb{N}$ and a sequence $\{y^k\} \subseteq Y$ with $y^k \rightarrow \bar{y}$ and $y^k \in Tx^k$ for all $k \geq k_0$;
- (ii) closed at $\bar{x} \in X$ if for all sequences $\{x^k\} \subseteq X$ with $x^k \rightarrow \bar{x}$ and all sequences $\{y^k\} \rightarrow \bar{y}$ with $y^k \in Tx^k$ for all $k \in \mathbb{N}$ sufficiently large, we have $\bar{y} \in T\bar{x}$;
- (iii) continuous at $\bar{x} \in X$ if it is lower semicontinuous and closed at $\bar{x} \in X$;
- (iv) lower semicontinuous, closed or continuous on X if it is lower semicontinuous, closed or continuous at every $x \in X$, respectively.

3. Main result

THEOREM 3.1. The function ϕ in (2.2) is a gap function of EQVI, that is, it satisfies

- (i) $\phi(x) \geq 0$ for all $x \in Sx$;
- (ii) \bar{x} is a solution of EQVI if and only if $\bar{x} \in S\bar{x}$ and $\phi(\bar{x}) = 0$.

Proof. (i) Taking $y = x$ in (2.2) yields the result.

- (ii) (\Rightarrow) If \bar{x} solves EQVI, then we have

$$\langle F\bar{x}, x - \bar{x} \rangle \geq f(\bar{x}) - f(x) \quad \text{for all } x \in S\bar{x}.$$

As Ω is nonnegative on $E \times E$,

$$\langle F\bar{x}, x - \bar{x} \rangle \geq f(\bar{x}) - f(x) - \Omega(\bar{x}, x) \quad \text{for all } x \in S\bar{x}.$$

Hence

$$\langle F\bar{x}, x - \bar{x} \rangle + f(x) - f(\bar{x}) + \Omega(\bar{x}, x) \geq 0 \quad \text{for all } x \in S\bar{x}.$$

This implies that $\phi(\bar{x}) \leq 0$. Obviously, $\bar{x} \in S\bar{x}$ and $\phi(\bar{x}) \geq 0$ by (i). Thus $\phi(\bar{x}) = 0$.

(\Leftarrow) Assume that $\bar{x} \in S\bar{x}$ and $\phi(\bar{x}) = 0$. Define a function

$$\tilde{\phi}(x) = - \inf_{y \in S\bar{x}} \{ \langle Fx, y - x \rangle + f(y) - f(x) + \Omega(x, y) \} \quad \text{for all } x \in E.$$

In fact, this function $\tilde{\phi}(x)$ is nothing but the gap function of EVI in Kum and Lee [6] for the fixed closed convex subset $S\bar{x}$ of E . Also, it is obvious that $\tilde{\phi}(x) \geq 0$ for all $x \in S\bar{x}$. Moreover, $\tilde{\phi}(\bar{x}) = \phi(\bar{x}) = 0$ and $\bar{x} \in S\bar{x}$. This means that \bar{x} is a solution of EVI by virtue of Kum and Lee [6, Theorem 3.1]. That is, $\langle F\bar{x}, x - \bar{x} \rangle \geq f(\bar{x}) - f(x)$ for all $x \in S\bar{x}$, which amounts to saying that \bar{x} is a solution of EQVI, as desired. \square

For more discussions of ϕ , from now on, it is further assumed that E is a reflexive Banach space and $\Omega(x, \cdot)$ is strongly convex in the second variable for each $x \in E$. Recall that a function $g : E \rightarrow \mathbf{R}$ is said to be *strongly convex* with modulus a ($a > 0$) if for all $x, y \in E$, and $\alpha \in [0, 1]$, we have

$$g(\alpha x + (1 - \alpha)y) \leq \alpha g(x) + (1 - \alpha)g(y) - \frac{1}{2}a\alpha(1 - \alpha)\|x - y\|^2.$$

Thus for each fixed x , the function

$$\psi(y) = \langle Fx, y - x \rangle + f(y) - f(x) + \Omega(x, y)$$

is strongly convex with respect to y , so it is coercive in the sense that $\lim_{\|y\| \rightarrow +\infty} \frac{\psi(y)}{\|y\|} \geq +\infty$. (For a proof, see Kum and Lee [6, Proposition 4.2].) As the closed unit ball B in E is weakly compact, $\psi(y)$ has a unique minimizer $z(x) \in Sx$ over the closed convex subset Sx so that

$$\begin{aligned} \phi(x) &= - \min_{y \in Sx} \psi(y) = -\psi(z(x)) \\ &= -\langle Fx, z(x) - x \rangle - f(z(x)) + f(x) - \Omega(x, z(x)). \end{aligned} \quad (3.1)$$

Then a solution \bar{x} of EQVI has the following fixed point characterization analogous to the case of VI [2]:

THEOREM 3.2. \bar{x} is a solution of EQVI if and only if $\bar{x} = z(\bar{x})$.

Proof. (\Rightarrow) Let \bar{x} be a solution of EQVI. Since $z(\bar{x})$ is a minimizer of the convex function $\psi(y)$ over the closed convex set $S\bar{x}$, we have

$$0 \in F\bar{x} + \partial f(z(\bar{x})) + \nabla_y \Omega(\bar{x}, z(\bar{x})) + N_{S\bar{x}}(z(\bar{x})) \tag{3.2}$$

where $\partial f(z(\bar{x}))$ is the subdifferential of f at $z(\bar{x})$ and $N_{S\bar{x}}(z(\bar{x})) = \{x^* \in E^* \mid \langle x^*, x - z(\bar{x}) \rangle \leq 0 \text{ for all } x \in S\bar{x}\}$ is the normal cone of $S\bar{x}$ at $z(\bar{x})$. Hence there exists $x^* \in \partial f(z(\bar{x}))$ such that

$$-F\bar{x} - x^* - \nabla_y \Omega(\bar{x}, z(\bar{x})) \in N_{S\bar{x}}(z(\bar{x})).$$

That is,

$$\langle F\bar{x} + x^* + \nabla_y \Omega(\bar{x}, z(\bar{x})), x - z(\bar{x}) \rangle \geq 0 \text{ for all } x \in S\bar{x}.$$

Taking $x = \bar{x} \in S\bar{x}$, we obtain

$$\begin{aligned} \langle x^* + \nabla_y \Omega(\bar{x}, z(\bar{x})), \bar{x} - z(\bar{x}) \rangle &\geq \langle F\bar{x}, z(\bar{x}) - \bar{x} \rangle \geq f(\bar{x}) - f(z(\bar{x})) \\ &\geq \langle x^*, \bar{x} - z(\bar{x}) \rangle \end{aligned}$$

because \bar{x} is a solution of EQVI and $x^* \in \partial f(z(\bar{x}))$. Thus we get

$$\langle \nabla_y \Omega(\bar{x}, z(\bar{x})), \bar{x} - z(\bar{x}) \rangle \geq 0. \tag{3.3}$$

From the strong convexity of $\Omega(\bar{x}, \cdot)$ with (3.3) it follows that

$$0 \leq \langle \nabla_y \Omega(\bar{x}, z(\bar{x})), \bar{x} - z(\bar{x}) \rangle + \mu \|\bar{x} - z(\bar{x})\|^2 \leq \Omega(\bar{x}, \bar{x}) - \Omega(\bar{x}, z(\bar{x})) \leq 0$$

for some $\mu > 0$. Here the second inequality comes from Kum and Lee [6, (4.3) of Proposition 4.2]. This implies that $\Omega(\bar{x}, \bar{x}) - \Omega(\bar{x}, z(\bar{x})) = -\Omega(\bar{x}, z(\bar{x})) = 0$, hence $\bar{x} = z(\bar{x})$ by Assumption 2.1.

(\Leftarrow) Assume that $\bar{x} = z(\bar{x})$. By (3.2), for some $x^* \in \partial f(z(\bar{x}))$, we again obtain that

$$\langle F\bar{x} + x^* + \nabla_y \Omega(\bar{x}, z(\bar{x})), x - z(\bar{x}) \rangle \geq 0 \text{ for all } x \in S\bar{x}.$$

Thus

$$\langle F\bar{x} + x^* + \nabla_y \Omega(\bar{x}, \bar{x}), x - \bar{x} \rangle \geq 0 \text{ for all } x \in S\bar{x}.$$

Hence

$$\langle F\bar{x} + x^*, x - \bar{x} \rangle \geq 0 \text{ for all } x \in S\bar{x}.$$

On the other hand,

$$\langle x^*, x - \bar{x} \rangle = \langle x^*, x - z(\bar{x}) \rangle \leq f(x) - f(z(\bar{x})) = f(x) - f(\bar{x}).$$

Therefore

$$\langle F\bar{x}, x - \bar{x} \rangle \geq \langle x^*, \bar{x} - x \rangle \geq f(\bar{x}) - f(x) \text{ for all } x \in S\bar{x},$$

which means that \bar{x} is a solution of EQVI. □

Now we deal with the continuity of the gap function ϕ of EQVI in terms of that of $z(x)$.

THEOREM 3.3. *Let the function $F : E \rightarrow E^*$ be continuous and let the multifunction $S : E \rightrightarrows E$ be continuous. Then the gap function $\phi : E \rightarrow \mathbb{R}$ has a closed graph in $E \times \mathbb{R}$.*

Proof. By Hogan [5, Theorem 8], $z : x \mapsto \{z(x)\}$ is closed on E , i.e., the function z has a closed graph in $E \times E$. It follows from (3.1) that $\phi(x) = -\langle Fx, z(x) - x \rangle - f(z(x)) + f(x) - \Omega(x, z(x))$ has the same property. \square

THEOREM 3.4. *Let F and S be as in Theorem 3.3. Assume that the image $S(E)$ is relatively compact. Then the gap function $\phi : E \rightarrow \mathbb{R}$ is continuous.*

Proof. By Hogan [5, Corollary 8.1], the function $z : E \rightarrow E$ is continuous, hence $\phi : E \rightarrow \mathbb{R}$ is continuous, too. \square

References

- [1] G. Y. Chen, C. J. Goh, and X. Q. Yang, *On gap functions and duality of variational inequality problems*, J. Math. Anal. Appl. **214** (1997), 658-673.
- [2] M. Fukushima, *Equivalent differentiable optimization problems and descent methods for asymmetric variational inequality problems*, Math. Programming **53** (1992), 99-110.
- [3] F. Giannessi, *Separation of sets and gap functions for quasi-variational inequalities*, in F. Giannessi and A. Maugeri (eds.): *Variational Inequality and Network Equilibrium Problems*, Plenum Press, New York, 1995, 101-121.
- [4] N. Harms, C. Kanzow, and O. Stein, *Smoothness properties of a regularized gap function for quasi-variational inequalities*, to appear in Optim. Meth. Software.
- [5] W. W. Hogan, *Point-to-set maps in mathematical programming*, SIAM Rev. **15** (1973), 591-603.
- [6] S. H. Kum and G. M. Lee, *On gap functions of variational inequality in a Banach space*, J. Korean Math. Soc. **38** (2001), 683-695.
- [7] K. Taji, *On gap functions for quasi-variational inequalities*, Abstr. Appl. Anal. (2008), Article ID 531361.

*

Department of Mathematics Education
 Chungbuk National University
 Cheongju 361-763, Republic of Korea
E-mail: shkum@cbnu.ac.kr