

**q -DEDEKIND-TYPE DAEHEE-CHANGHEE SUMS WITH
WEIGHT α ASSOCIATED WITH MODIFIED q -EULER
POLYNOMIALS WITH WEIGHT α**

JONG JIN SEO*, SERKAN ARACI**, AND MEHMET ACIKGOZ***

ABSTRACT. Recently, q -Dedekind-type sums related to q -Euler polynomials was studied by Kim in [T. Kim, Note on q -Dedekind-type sums related to q -Euler polynomials, Glasgow Math. J. 54 (2012), 121-125]. It is aim of this paper to consider a p -adic continuous function for an odd prime to inside a p -adic q -analogue of the higher order Dedekind-type sums with weight related to modified q -Euler polynomials with weight by using Kim's p -adic q -integral.

1. Introduction

Let p be a fixed odd prime number. Throughout this paper \mathbb{Z}_p , \mathbb{Q}_p , \mathbb{C} and \mathbb{C}_p will denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex numbers and the completion of algebraic closure of \mathbb{Q}_p , respectively.

Let v_p be normalized exponential valuation of \mathbb{C}_p with

$$|p|_p = p^{-v_p(p)} = \frac{1}{p}.$$

When one speaks of q -extension, q is variously considered as an indeterminate, a complex number $q \in \mathbb{C}$ or p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, we assume that $|q| < 1$. If $q \in \mathbb{C}_p$, we assume that $|1 - q|_p < 1$ (see [1-16]).

Received November 27, 2013; Accepted January 06, 2013.

2010 Mathematics Subject Classification: Primary 11B68, 11S80; Secondary 11M06.

Key words and phrases: Dedekind sums, q -Dedekind-type sums, p -adic q -integral, q -Euler numbers and polynomials, modified q -Euler numbers and polynomials.

Correspondence should be addressed to Serkan Araci, mts_rkn@hotmail.com.

A q -extension of p -adic Haar measure is defined by Kim as follows: for any positive integer N ,

$$\mu_q(a + p^N \mathbb{Z}_p) = (-q)^a \frac{(1+q)}{1+q^{p^N}}$$

for $0 \leq a < p^N$ and this can be extended to a measure on \mathbb{Z}_p (for details, see [1–4, 6–16]).

The modified q -Euler polynomials with weight α are defined by Rim and Jeong as follows:

$$(1.1) \quad \tilde{\mathcal{E}}_{n,q}^{(\alpha)}(x) = \int_{\mathbb{Z}_p} q^{-y} \left(\frac{1 - q^{\alpha(x+y)}}{1 - q^\alpha} \right) d\mu_q(y)$$

for $n \in \mathbb{Z}_+ := \{0, 1, 2, 3, \dots\}$. We note that

$$\lim_{q \rightarrow 1} \tilde{\mathcal{E}}_{n,q}^{(\alpha)}(x) = E_n(x)$$

where E_n are the famous Euler polynomials, which are defined by means of the following generating function:

$$\sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} = e^{tx} \frac{2}{e^t + 1}, \quad |t| < \pi$$

(for details, see [15]). Taking $x = 0$ into (1.1), then, we have $\tilde{\mathcal{E}}_{n,q}^{(\alpha)}(0) := \tilde{\mathcal{E}}_{n,q}^{(\alpha)}$ are called modified q -Euler numbers with weight α .

These numbers and polynomials have the following identities:

$$(1.2) \quad \tilde{\mathcal{E}}_{n,q}^{(\alpha)} = \frac{1+q}{(1-q^\alpha)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{1}{1+q^{\alpha l}},$$

$$(1.3) \quad \tilde{\mathcal{E}}_{n,q}^{(\alpha)}(x) = \frac{1+q}{(1-q^\alpha)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{q^{\alpha l x}}{1+q^{\alpha l}},$$

$$(1.4) \quad \tilde{\mathcal{E}}_{n,q}^{(\alpha)}(x) = \sum_{l=0}^n \binom{n}{l} q^{\alpha l x} \tilde{\mathcal{E}}_{l,q}^{(\alpha)} \left(\frac{1 - q^{\alpha x}}{1 - q^\alpha} \right)^{n-l}$$

and

$$(1.5) \quad \tilde{\mathcal{E}}_{n,q}^{(\alpha)}(x) = \left(\frac{1 - q^{\alpha d}}{1 - q^\alpha} \right) \sum_{a=0}^{d-1} (-1)^a \tilde{\mathcal{E}}_{n,q}^{(\alpha)} \left(\frac{x+a}{d} \right),$$

$d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$

(for more information, see [15]).

For any positive integer h, k and m , Dedekind-type DC sums are defined by Kim in [6], [7] and [8] as follows:

$$S_m(h, k) = \sum_{M=1}^{k-1} (-1)^{M-1} \frac{M}{k} \overline{E}_m\left(\frac{hM}{k}\right)$$

where $\overline{E}_m(x)$ are the m -th periodic Euler function. Kim gave some interesting properties Dedekind-type DC sums. He also constructed a p -adic continuous function for an odd prime number to contain a p -adic q -analogue of the higher order Dedekind-type DC sums $k^m S_{m+1}(h, k)$ in [7]. After Simsek also studied to q -analogue of Dedekind-type sums. He also derived their interesting properties. By the same motivation, we, by using p -adic q -integral on \mathbb{Z}_p , will construct weighted p -adic q -analogue of the higher order Dedekind-type DC sums $k^m S_{m+1}(h, k)$.

2. Weighted q -analogue of Dedekind-type Sums associated with modified q -Euler polynomials with weight α

Let w denotes the Teichmüller character (mod p). For $x \in \mathbb{Z}_p^* := \mathbb{Z}_p/p\mathbb{Z}_p$, set

$$\langle x : q \rangle = w^{-1}(x) \left(\frac{1 - q^x}{1 - q} \right).$$

Let a and N be positive integers with $(p, a) = 1$ and $p \mid N$. We now consider the following

$$\tilde{T}_q^{(\alpha)}(s, a, N : q^N) = w^{-1}(a) \langle x : q^\alpha \rangle^s \sum_{j=0}^{\infty} \binom{s}{j} q^{\alpha a j} \left(\frac{1 - q^{\alpha N}}{1 - q^{\alpha a}} \right)^j \tilde{\mathcal{E}}_{j, q^N}^{(\alpha)}.$$

In particular, if $m + 1 \equiv 0 \pmod{p - 1}$, then

$$\begin{aligned} & \tilde{T}_q^{(\alpha)}(m, a, N : q^N) \\ &= \left(\frac{1 - q^{\alpha a}}{1 - q^\alpha} \right)^m \sum_{j=0}^m \binom{m}{j} q^{\alpha a j} \tilde{\mathcal{E}}_{j, q^N}^{(\alpha)} \left(\frac{1 - q^{\alpha N}}{1 - q^{\alpha a}} \right)^j \\ &= \left(\frac{1 - q^{\alpha N}}{1 - q^\alpha} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha N(x + \frac{a}{N})}}{1 - q^{\alpha N}} \right)^m q^{-Nx} d\mu_{q^N}(x). \end{aligned}$$

That is, $\tilde{T}_q^{(\alpha)}(m, a, N : q^N)$ is a continuous p -adic extension of $\left(\frac{1 - q^{\alpha N}}{1 - q^\alpha} \right)^n \tilde{\mathcal{E}}_{n, q^N}^{(\alpha)}\left(\frac{a}{N}\right)$.

Let $[\cdot]$ be the Gauss' symbol and let $\{x\} = x - [x]$. Then, we consider q -analogue of the higher order Dedekind-type DC sums $\tilde{S}_{m,q}^{(\alpha)}(h, k : q^l)$ as

$$\begin{aligned} & \tilde{S}_{m,q}^{(\alpha)}(h, k : q^l) \\ &= \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^{\alpha k}} \right) \int_{\mathbb{Z}_p} q^{-lx} \left(\frac{1 - q^{\alpha(lx+l\{\frac{hM}{k}\})}}{1 - q^{\alpha l}} \right)^m d\mu_{q^l}(x). \end{aligned}$$

If $m + 1 \equiv 0 \pmod{p-1}$

$$\begin{aligned} & \left(\frac{1 - q^{\alpha k}}{1 - q^{\alpha}} \right)^{m+1} \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^{\alpha k}} \right) \\ & \quad \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha k(x+\frac{hM}{k})}}{1 - q^{\alpha k}} \right)^m q^{-kx} d\mu_{q^k}(x) \\ &= \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^{\alpha}} \right)^m \left(\frac{1 - q^{\alpha k}}{1 - q^{\alpha}} \right)^m \\ & \quad \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha k(x+\frac{hM}{k})}}{1 - q^{\alpha k}} \right)^m q^{-kx} d\mu_{q^k}(x) \end{aligned}$$

where $p \mid k$, $(hM, p) = 1$ for each M . From (1.1), we note that

$$\begin{aligned} (2.1a) \quad & \left(\frac{1 - q^{\alpha k}}{1 - q^{\alpha}} \right)^{m+1} \tilde{S}_{m,q}^{(\alpha)}(h, k : q^k) \\ &= \sum_{M=1}^{k-1} \left(\frac{1 - q^{\alpha M}}{1 - q^{\alpha}} \right) \left(\frac{1 - q^{\alpha k}}{1 - q^{\alpha}} \right)^m (-1)^{M-1} \\ & \quad \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha k(x+\frac{hM}{k})}}{1 - q^{\alpha k}} \right)^m q^{-kx} d\mu_{q^k}(x) \\ &= \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^{\alpha}} \right) \tilde{T}_q^{(\alpha)}(m, (hM)_k : q^k) \end{aligned}$$

where $(hM)_k$ denotes the integer x such that $0 \leq x < n$ and $x \equiv \alpha \pmod{k}$. It is not difficult to show that

(2.2)

$$\begin{aligned} & \int_{\mathbb{Z}_p} q^{-t} \left(\frac{1 - q^{\alpha(x+t)}}{1 - q^\alpha} \right)^k d\mu_q(t) \\ &= \left(\frac{1 - q^{\alpha m}}{1 - q^\alpha} \right)^k \frac{1 + q}{1 + q^m} \sum_{i=0}^{m-1} (-1)^i \int_{\mathbb{Z}_p} q^{-mt} \left(\frac{1 - q^{\alpha m(t + \frac{x+i}{m})}}{1 - q^{\alpha m}} \right)^k d\mu_{q^m}(t). \end{aligned}$$

By (2.1a) and (2.2), we easily see that

$$\begin{aligned} (2.3) \quad & \left(\frac{1 - q^{\alpha N}}{1 - q^\alpha} \right) \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha N(x + \frac{a}{N})}}{1 - q^{\alpha N}} \right)^m q^{-Nx} d\mu_{q^N}(x) \\ &= \frac{1 + q^N}{1 + q^{Np}} \sum_{i=0}^{p-1} (-1)^i \left(\frac{1 - q^{\alpha Np}}{1 - q^\alpha} \right)^m \\ & \quad \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha p N(x + \frac{a+iN}{pN})}}{1 - q^{\alpha p N}} \right) q^{-xpN} d\mu_{q^{pN}}(x) \end{aligned}$$

From (2.1a), (2.2) and (2.3), we note that the p -adic integration is given by

$$\begin{aligned} & \tilde{T}_q^{(\alpha)}(s, a, N : q^N) \\ &= \frac{1 + q^N}{1 + q^{Np}} \sum_{\substack{0 \leq i \leq p-1 \\ a+iN \not\equiv 0 \pmod{p}}} (-1)^i \tilde{T}_q^{(\alpha)}(s, (a+iN)_{pN}, p^N : q^{pN}) \end{aligned}$$

such that

$$\begin{aligned} & \tilde{T}_q^{(\alpha)}(m, a, N : q^N) \\ &= \left(\frac{1 - q^{\alpha N}}{1 - q^\alpha} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha N(x + \frac{a}{N})}}{1 - q^{\alpha N}} \right)^m q^{-Nx} d\mu_{q^N}(x) \\ & \quad - \left(\frac{1 - q^{\alpha Np}}{1 - q^\alpha} \right)^m \int_{\mathbb{Z}_p} \left(\frac{1 - q^{\alpha p N(x + \frac{a+iN}{pN})}}{1 - q^{\alpha p N}} \right)^m q^{-pNx} d\mu_{q^{pN}}(x) \end{aligned}$$

where $(p^{-1}a)_N$ denotes the integer x with $0 \leq x < N$, $px \equiv a \pmod{N}$ and m is integer with $m+1 \equiv 0 \pmod{p-1}$. Therefore, we procure the following

$$\begin{aligned}
& \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^{\alpha}} \right) \tilde{T}_q^{(\alpha)}(m, hM, k : q^k) \\
&= \left(\frac{1 - q^{\alpha k}}{1 - q^{\alpha}} \right)^{m+1} \tilde{S}_{m,q}^{(\alpha)}(h, k : q^k) \\
&\quad - \left(\frac{1 - q^{\alpha k}}{1 - q^{\alpha}} \right)^{m+1} \left(\frac{1 - q^{\alpha kp}}{1 - q^{\alpha k}} \right) \tilde{S}_{m,q}^{(\alpha)}((p^{-1}h), k : q^{pk})
\end{aligned}$$

where $p \nmid k$ and $p \nmid hm$ for each M . Thus, we state the following definition.

DEFINITION 2.1. Let h, k be positive integer with $(h, k) = 1$, $p \nmid k$. For $s \in \mathbb{Z}_p$, we define p -adic Dedekind-type DC sums as follows:

$$\tilde{S}_{p,q}^{(\alpha)}(s : h, k : q^k) = \sum_{M=1}^{k-1} (-1)^{M-1} \left(\frac{1 - q^{\alpha M}}{1 - q^{\alpha}} \right) \tilde{T}_q^{(\alpha)}(m, hM, k : q^k).$$

Then, we can give the following theorem.

THEOREM 2.2. For $m + 1 \equiv 0 \pmod{p-1}$ and $(p^{-1}a)_N$ denotes the integer x with $0 \leq x < N$, $px \equiv a \pmod{N}$, then, we have

$$\begin{aligned}
& \tilde{S}_{p,q}^{(\alpha)}(s : h, k : q^k) \\
&= \left(\frac{1 - q^{\alpha k}}{1 - q^{\alpha}} \right)^{m+1} \tilde{S}_{m,q}^{(\alpha)}(h, k : q^k) \\
&\quad - \left(\frac{1 - q^{\alpha k}}{1 - q^{\alpha}} \right)^{m+1} \left(\frac{1 - q^{\alpha kp}}{1 - q^{\alpha k}} \right) \tilde{S}_{m,q}^{(\alpha)}((p^{-1}h), k : q^{pk}).
\end{aligned}$$

For $\alpha = 1$, we have to Kim's results in [7]. This result seems to be interesting for further work in [6 – 8, 13].

References

- [1] S. Araci, Novel Identities for the q -Genocchi numbers and polynomials, *Journal of Function Spaces and Applications*, 2012 (2012), Article ID 214961, 12 pages.
- [2] S. Araci, M. Acikgoz, and K. H. Park, A note on the q -analogue of Kim's p -adic log gamma type functions associated with q -extension of Genocchi and Euler numbers with weight α , *Bull. Korean Math. Soc.* 50 (2013), no. 2, 583-588.
- [3] S. Araci, D. Erdal, and J. J. Seo, A study on the fermionic p -adic q -integral representation on \mathbb{Z}_p associated with weighted q -Bernstein and q -Genocchi polynomials, *Abstract and Applied Analysis*, **2011**, Article ID 649248, 10 pages.

- [4] S. Araci and M. Acikgoz, A note on the Frobenius-Euler numbers and polynomials associated with Bernstein polynomials, *Advanced Studies in Contemporary Mathematics* **22** (2012), no. 3, 399-406.
- [5] S. Araci, M. Acikgoz, A. Bagdasaryan, and E. Şen, The Legendre Polynomials Associated with Bernoulli, Euler, Hermite and Bernstein Polynomials, *Turkish Journal of Analysis and Number Theory*, no. 1 (1) (**2013**): 1-3. doi: 10.12691/tjant-1-1-1.
- [6] T. Kim, A note on p -adic q -Dedekind sums, *C. R. Acad. Bulgare Sci.* **54** (2001), 37-42.
- [7] T. Kim, Note on q -Dedekind-type sums related to q -Euler polynomials, *Glasgow Math. J.* **54** (2012), 121-125.
- [8] T. Kim, Note on Dedekind type DC sums, *Adv. Stud. Contemp. Math.* **18** (2009), 249-260.
- [9] T. Kim, The modified q -Euler numbers and polynomials, *Adv. Stud. Contemp. Math.* **16** (2008), 161-170.
- [10] T. Kim, q -Volkenborn integration, *Russ. J. Math. Phys.* **9** (2002), 288-299.
- [11] T. Kim, On p -adic interpolating function for q -Euler numbers and its derivatives, *J. Math. Anal. Appl.* **339** (2008), 598-608.
- [12] T. Kim, On a q -analogue of the p -adic log gamma functions and related integrals, *J. Number Theory* **76** (1999), 320-329.
- [13] Y. Simsek, q -Dedekind type sums related to q -zeta function and basic L -series, *J. Math. Anal. Appl.* **318** (2006), 333-351.
- [14] M. Acikgoz and Y. Simsek, On multiple interpolation function of the Nörlund-type q -Euler polynomials, *Abst. Appl. Anal.* **2009** (2009), Article ID 382574, 14 pages.
- [15] S. H. Rim and J. Jeong, A note on the Modified q -Euler Numbers and polynomials with weight α , *International Mathematical Forum*, **6** (2011), no. 65, 3245-3250.
- [16] C. S. Ryoo, A note on the weighted q -Euler numbers and polynomials, *Adv. Stud. Contemp. Math.* **21** (2011), 47-54.

*

Department of Applied Mathematics
Pukyong National University
Busan 608-737, Republic of Korea
E-mail: seo2011@pknu.ac.kr

**

Atatürk Street, TR-31290 Hatay, TURKEY
E-mail: mtsrkn@hotmail.com

Department of Mathematics
University of Gaziantep
27310 Gaziantep, TURKEY
E-mail: acikgoz@gantep.edu.tr