

# Chloroplast Genome Evolution in Early Diverged Leptosporangiate Ferns

Hyoung Tae Kim, Myong Gi Chung<sup>1</sup>, and Ki-Joong Kim\*

In this study, the chloroplast (cp) genome sequences from three early diverged leptosporangiate ferns were completed and analyzed in order to understand the evolution of the genome of the fern lineages. The complete cp genome sequence of *Osmunda cinnamomea* (Osmundales) was 142,812 base pairs (bp). The cp genome structure was similar to that of eusporangiate ferns. The gene/intron losses that frequently occurred in the cp genome of leptosporangiate ferns were not found in the cp genome of *O. cinnamomea*. In addition, putative RNA editing sites in the cp genome were rare in *O. cinnamomea*, even though the sites were frequently predicted to be present in leptosporangiate ferns. The complete cp genome sequence of *Diplazium glaucum* (Gleicheniales) was 151,007 bp and has a 9.7 kb inversion between the *trnL-CAA* and *trnV-GCA* genes when compared to *O. cinnamomea*. Several repeated sequences were detected around the inversion break points. The complete cp genome sequence of *Lygodium japonicum* (Schizaeales) was 157,142 bp and a deletion of the *rpoC1* intron was detected. This intron loss was shared by all of the studied species of the genus *Lygodium*. The GC contents and the effective numbers of codons (ENCs) in ferns varied significantly when compared to seed plants. The ENC values of the early diverged leptosporangiate ferns showed intermediate levels between eusporangiate and core leptosporangiate ferns. However, our phylogenetic tree based on all of the cp gene sequences clearly indicated that the cp genome similarity between *O. cinnamomea* (Osmundales) and eusporangiate ferns are symplesiomorphies, rather than synapomorphies. Therefore, our data is in agreement with the view that Osmundales is a distinct early diverged lineage in the leptosporangiate ferns.

Division of Life Sciences, School of Life Sciences, Korea University, Seoul 136-701, Korea. <sup>1</sup>Department of Biology and the Research Institute of Natural Science, Gyeongsang National University, Jinju 660-701, Korea

\*Correspondence: kimkj@korea.ac.kr

Received 15 October, 2013; revised 11 April, 2014; accepted 14 April, 2014; published online 14 May, 2014

**Keywords:** chloroplast genomes, *Diplazium glaucum*, early diverged leptosporangiate ferns, inversion mutations, *Lygodium japonicum*, *Osmunda cinnamomea*, *rpoC1* intron loss

## INTRODUCTION

Comparative chloroplast (cp) genomic studies provide an invaluable source of information for understanding plant evolution and plant phylogeny. Therefore, the cp genome is the most widely studied genome when compared to the two other genomes found in plant cells. Approximately 400 cp genome sequences for land plants are available from a public database, but the majority of them belonged to seed plants (<http://www.ncbi.nlm.nih.gov/genomes/GenomesGroup.cgi?opt=plastid&taxid=3193>). The cp genomes hold numerous important evolutionary features, including structural changes, gene content differences, and base substitution patterns.

Structural changes in the cp genome, such as gene rearrangements (Chumley et al., 2006; Tangphatsomruang et al., 2010; Wu et al., 2007), gene/intron losses or duplications (Guisinger et al., 2011; Hiratsuka et al., 1989; Jansen et al., 2007), and small inversions (Kim and Lee, 2004; Yi and Kim, 2012) are well known at the genus, family, or ordinal levels of seed plants. Therefore, the genome evolution and phylogenetic relationships of seed plants are relatively well understood. However, the cp genome studies in ferns are limited to just a few lineages.

One of the distinct features of cp genomes is its high levels of adenosine and thiamine (AT) content (Sablok et al., 2011; Smith, 2009). However, a relatively wide range of AT content variation was reported for a number of different plant lineages (Smith, 2009). The GC content differences in the cp genomes usually correlate well with codon usage bias. The effective number of codons (ENCs) represents a simple way to measure synonymous codon usage bias and is independent of coding region length and amino acid composition (Wright, 1990). Therefore, the comparative ENC values may show a broad spectrum of base usage patterns among major lineages of plant groups.

Ferns are an important plant group for the understanding plant evolution because of the long evolutionary history and the complicated phylogenetic relationships (Pryer et al., 2004). The extant ferns are composed of one monophyletic class and 11 monophyletic orders (Pryer et al., 2009). Since the physical map of the *Osmunda cinnamomea* cp genome was known (Palmer and Stein, 1982; 1986), many researchers tried to understand the ferns cp genome evolution. Recently, the complete cp genome sequences of four orders of eusporangiate ferns were analyzed, and the data aided in understanding the evolutionary history of eusporangiate ferns (Grewe et al., 2013; Karol et al., 2010). In leptosporangiate ferns, the six complete

cp genome sequences have been reported. These are *Alsophila spinulosa* (Gao et al., 2009), *Adiantum capillus-veneris* (Wolf et al., 2003), *Cheilanthes lindheimeri*, *Pteridium aquilinum* subsp. *aquilinum* (Wolf et al., 2011), *Lygodium japonicum*, and *Marsilea crenata* (Gao et al., 2013). In addition, the incomplete sequences were also used for the reconstruction of the fern trees (Wolf et al., 2010). Although the comparative cp genome studies of eusporangiate ferns and leptosporangiate ferns were published, it is still difficult to understand cp genome evolution from all fern lineages because there is a lack of cp genome data for the early diverged fern lineages. This appears as missing links in data.

In order to provide the data in the missing lineages, we report three complete cp genome sequences from the early diverged leptosporangiate ferns in this paper. Two are newly reported groups (Osmundales and Gleicheniales) and one (Schizaeales) is a previously reported group. Using these data, we address the following two questions about cp genome evolution of early diverged leptosporangiate ferns: (i) which of the cp genome structures are more similar to that of basal Osmundales, and (ii) whether or not Osmundales really have an intermediate-type cp genome that is between eusporangiate and leptosporangiate ferns.

Osmundales consists of a monophyletic family, three genera, and ca. 20 species (Smith et al., 2006), but it includes more than 150 fossil species (Tidwell and Ash, 1994). Many researchers consider the Osmundales to be closely related to eusporangiate ferns (Pryer et al., 2001; 2004; Schneider et al., 2004; Schuettpelz and Pryer, 2007; Wolf et al., 1995). Osmundales also have been considered as intermediate taxa between eusporangiate and leptosporangiate ferns based on their external appearance, and anatomical and meristem characteristics (Cross, 1931a; 1931b; Freeberg and Gifford Jr, 1984; Gifford Jr, 1983). Using fossil records, Osmundaceae could be traced back to the Late Permian period, but the genus *Osmunda* was known from the Late Triassic period. *O. cinnamomea* seemed to exist since the Late Cretaceous period (Taylor et al., 2009) and was identified as a sister species to the rest of the Osmundaceae (Metzgar et al., 2008; Yatabe et al., 1999). Considering the morphological characters and phylogenetic relationship of ferns, the cp genome of *O. cinnamomea* could be regarded as an ancestral type from leptosporangiate ferns.

Gleicheniales consists of three families, 10 genera, and ca. 140 species, and most of the species are members of Gleicheniaceae (Smith et al., 2006). Gleicheniaceae is considered as an old lineage originating from the Permian (Pryer et al., 2004; Taylor et al., 2009). We report the cp genome sequence of *Diplopterygium glaucum*, which was a synonym of *Gleichenia japonica* (Iwatsuki et al., 1995), and was widely distributed in the Asian tropics.

Schizaeales consists of three families, four genera, and ca. 155 species (Smith et al., 2006). The oldest Schizaeaceae fossil originated from the Jurassic period (Taylor et al., 2009), and Schizaeales diverged from the core leptosporangiate ferns in the Permian (Pryer et al., 2004). The genus *Lygodium* is considered as a basal group in the Schizaeales (Schuettpelz and Pryer, 2007), and the cp genome of *Lygodium japonicum* had two large inversions and gene deletions (Gao et al., 2013; Wolf et al., 2010). We report the cp genome sequences of a Korean population of *L. japonicum*.

In this study, the complete cp genome sequences of *O. cinnamomea* and *D. glaucum* filled the evolutionary gap between eusporangiate and leptosporangiate ferns and gave us information, such as gene/intron losses, inversions, codon usage bias,

and patterns of repeating units in early diverged leptosporangiate ferns, thus allowing us to understand cp genome evolution in these interesting phylogenetic groups. In addition, knowing the complete cp genome sequence of *L. japonicum* is helpful to understanding the cp genome evolution between Chinese and Korean populations.

## MATERIALS AND METHODS

### DNA extraction, sequencing, and assembling

*O. cinnamomea* (KUS 2006-0338), *D. glaucum* (KUS 2000-0022), and *L. japonicum* (KUS 2007-0451) were collected in Korea. All voucher specimens were kept in Korea University's Herbarium (KUS). The genomic DNAs of *O. cinnamomea* and *D. glaucum* were isolated from the fresh leaves by the CTAB method (Doyle and Doyle, 1987) and purified by ultracentrifugation in cesium chloride/ethidium bromide gradients. We designed common primer sets based on the cp genome sequence in ferns using known complete cp genome sequences. Using the primers, we amplified the 5-10 kb overlapping cp genome fragments using TaKaRa LA Taq by PCR. PCR products were sequenced by Big-Dye chemistry and ABI3730XL. For *L. japonicum*, the chloroplasts were isolated by sucrose step-gradient methods (Palmer, 1986), and the cp genome was isolated using 5x lysis buffer (Jansen et al., 2005). The cp genome of *L. japonicum* was sequenced using GS-FLX 454 at Macrogen Co. (Korea). A total of 108,241 sequence reads were generated. A total of 4,430 reads were fully incorporated into the assembly and 2,595 reads were partially incorporated. There were 400 contigs over 500 bp, and the largest contig was 13,610 bp. Gaps were filled by PCR. All sequenced contigs were de novo assembled using Geneious 6.1.2 (Kearse et al., 2012). Gene annotations were performed by DOGMA (Wyman et al., 2004) and tRNAscan (Lowe and Eddy, 1997). Then, the exact positions of all genes were determined by local BLAST searches using the gene database of ferns obtained from NCBI.


### Phylogenetic analyses and comparative sequence analyses of cp genomes

Thirty-five cp genome sequences were used for the phylogenetic analysis (Table 1). We sampled all of the published complete cp genome sequences from monilophytes (14), lycophytes (4), and bryophytes (5), and eight selected species from spermatophytes. Two charophytes were included as outgroups. In addition, two unpublished monilophytes sequences were also included in these taxon samplings (H.-T. Kim and K.-J. Kim, unpublished data). Eighty-nine genes, including 84 protein coding genes and five ribosomal RNA genes, were aligned using MUSCLE program (Edgar, 2004), and the phylogenetic trees were constructed using four different tree building methods. First, the maximum parsimony (MP) tree was generated by PAUP (Swofford, 2003) under the options of equal character weighting, random taxon addition, and TBR branch swapping options. Gaps were treated as missing. Second, the neighbor joining (NJ) tree was generated with Geneious 6.1.7 using the HKY genetic distance model. Third, for the maximum likelihood (ML) tree, we selected the optimal model with Modeltest 3.7 (Posada and Crandall, 1998). The ML tree was evaluated by the GTR + I + G model using RAxML (Stamatakis, 2006; Stamatakis et al., 2008) that is performed using the CIPRES Science Gateway (Miller et al., 2010). The strengths of all of the internal branches in MP, NJ, and ML analyses were evaluated by 1,000 bootstrap replications. Fourth, the Bayesian inference (BI) tree was reconstructed by MrBayes under the

**Table 1.** The list of complete chloroplast genome sequences and *rpoC1* sequences

Target	Taxa	Group	GenBank	
Phylogenetic analysis	<i>Arabidopsis thaliana</i>	Spermatophytes	NC000932	
	<i>Panax chinseng</i>	Spermatophytes	NC006290	
	<i>Nymphaea alba</i>	Spermatophytes	NC006050	
	<i>Amborella trichopoda</i>	Spermatophytes	NC005086	
	<i>Cycas revoluta</i>	Spermatophytes	NC020319	
	<i>Ginkgo biloba</i>	Spermatophytes	NC016986	
	<i>Welwitschia mirabilis</i>	Spermatophytes	NC010654	
	<i>Pinus koraiensis</i>	Spermatophytes	NC004677	
	<i>Adiantum capillus-veneris</i>	Polypodiales(core leptosporangiate ferns)	NC004766	
	<i>Cheilanthes lindheimeri</i>	Polypodiales(core leptosporangiate ferns)	NC014592	
	<i>Pteridium aquilinum</i> subsp. <i>aquilinum</i>	Polypodiales(core leptosporangiate ferns)	NC014348	
	<i>Alsophila spinulosa</i>	Cyatheaales(core leptosporangiate ferns)	NC012818	
	<i>Marsilea crenata</i>	Salviniales(core leptosporangiate ferns)	KC536646	
	<i>Lygodium japonicum</i> (K.-J. Kim et al. KUS 2006-0338)	Schizaeales (early diverged leptosporangiate ferns)	KF225593*	
	<i>Diplazium glaucum</i> (C.-H. Kim et al. KUS 2000-0022)	Gleicheniales (early diverged leptosporangiate ferns)	KF225594*	
	<i>Osmunda cinnamomea</i> (H.-W. Kim et al. KUS 2007-0451)	Osmundales (early diverged leptosporangiate ferns)	KF225592*	
	<i>Angiopteris evecta</i>	Marattiales (eusporangiate ferns)	NC008829	
	<i>Ophioglossum californicum</i>	Ophioglossales (eusporangiate ferns)	NC020147	
	<i>Mankyua chejuensis</i>	Ophioglossales (eusporangiate ferns)	NC017006	
	<i>Psilotum nudum</i> 1	Psilotales (eusporangiate ferns)	NC003386	
	<i>Psilotum nudum</i> 2	Psilotales (eusporangiate ferns)	KC117179	
	<i>Equisetum arvense</i> 1	Equisetales (eusporangiate ferns)	NC014699	
	<i>Equisetum arvense</i> 2	Equisetales (eusporangiate ferns)	JN968380	
	<i>Equisetum hyemale</i>	Equisetales (eusporangiate ferns)	NC020146	
	<i>Huperzia lucidula</i>	Lycophytes	NC006861	
	<i>Isoetes flaccida</i>	Lycophytes	NC014675	
	<i>Selaginella moellendorffii</i>	Lycophytes	NC013086	
	<i>Selaginella uncinata</i>	Lycophytes	AB197035	
	<i>Anthoceros formosae</i>	Bryophytes	NC004543	
	<i>Syntrichia ruralis</i>	Bryophytes	NC012052	
	<i>Physcomitrella patens</i> subsp. <i>patens</i>	Bryophytes	NC005087	
	<i>Marchantia polymorpha</i>	Bryophytes	NC001319	
	<i>Aneura mirabilis</i>	Bryophytes	NC010359	
	<i>Chaetosphaeridium globosum</i>	Charophytes	NC004115	
	<i>Chara vulgaris</i>	Charophytes	NC008097	
	<i>rpoC1</i> intron analysis	<i>Lygodium japonicum</i> (K.-J. Kim et al. TC 2010-0136)	Schizaeales (early diverged leptosporangiate ferns)	KF225595*
		<i>Lygodium polystachyum</i> (K.-J. Kim et al. TL 2008-1701)	Schizaeales (early diverged leptosporangiate ferns)	KF225596*
		<i>Vandenboschia striata</i> (K.-J. Kim et al. TC 2010-1362)	Hymenophyllales (early diverged leptosporangiate ferns)	KF225597*
		<i>Lygodium flexuosum</i> (K.-J. Kim et al. TL 2008-1886)	Schizaeales (early diverged leptosporangiate ferns)	Not sequenced
		<i>Lygodium microphyllum</i> (K.-J. Kim et al. TL 2008-1661)	Schizaeales (early diverged leptosporangiate ferns)	Not sequenced
<i>Lygodium salicifolium</i> (K.-J. Kim et al. TL 2008-1772)		Schizaeales (early diverged leptosporangiate ferns)	Not sequenced	

Asterisk on the GenBank accession numbers indicate newly reported sequences in this paper.



**Fig. 1.** Phylogenetic tree of ferns and related groups. The best ML tree was constructed using RAxML under the GTR + I + G base substitution model (A). The three consecutive numeric values on each internal node in (A) tree indicate the ML bootstrap support percentage, Bayesian probability, and MP bootstrap support percentage, respectively. Two alternative suboptimal tree topologies (B, C) that are observed frequently in MP, BI, and NJ analyses, were also generated using the topology constraint analyses option of RaxML using the same base substitution assumption. The lycophytes was a sister group of the euphyllophytes (spermatophytes + monilophytes) in the (A) tree, while the lycophytes was a sister group to the spermatophytes in the (B) tree. The Equisetales was a sister group to all other members of monilophytes in the (a) tree, while the Equisetales was a sister clade to the Psilotales and Ophioglossales in the (C) tree. The (D) tree topology represents the combination of the topology of the (B) and (C) trees.

following conditions: nst = 6, rates = invgamma, Ngen = 500,000 and sample = 100, using the CIPRES Science Gateway (Miller et al., 2010).

The cp genomes modifications, such as gene/intron gains or losses, inversion events, and the anticodon changes, were treated as binary characters. A total of 30 variable evolutionary events were recorded from the fern lineages. Next, the character states were plotted on the ML tree topology in order to deduce the evolutionary direction of these characteristics. The evolutionary directions were accounted on the ACCTRAN criteria on the parsimony analysis using PAUP (Swofford, 2003).

The complete cp genome sequences of 194 species of land plants were used to analyze the GC contents of coding sequences in the cp genome (Supplementary Table 1). All cp genome sequences were obtained from NCBI Organellar Genome Resources. The GC contents of the entire coding gene (GCall), first position (GC1), second position (GC2), and third position (GC3), and the effective numbers of codons (ENCs) (Wright, 1990) were calculated using Acua 1.0 (Vetrivel et al., 2007). We also analyzed dispersed repeats using REPuter (Kurtz et al., 2001). Then, each repeat sequence was sorted by similarity. These repeat sequences were reanalyzed using a DNA pattern search ([http://www.geneinfinity.org/sms/sms\\_DNApatterns.html#](http://www.geneinfinity.org/sms/sms_DNApatterns.html#)) for calculating the exact numbers of repeat sequences in the complete cp genome (Kim et al., 2009; Yi et al., 2012).

### Analysis of *rpoC1* intron loss

Five species of the genus *Lygodium* and one accession of *Vandenboschia striata* were collected from Laos, China and Korea for *rpoC1* intron loss analysis. All specimens were kept in the KUS (Table 1). Their genomic DNAs were obtained using the same method as described for *O. cinnamomea* and *D. glaucum*. The primer set was designed based on the consensus *rpoC1* sequence among 15 fern cp genomes. The forward primer 'FrpoC1 exon1' (5'-GAAAGCCYAGTATTGCGA-3') was located at the end of exon1, and the reverse primer 'FrpoC1 exon2' (5'-ATGCARACGAATRGCRGCTCC-3') was in the middle of exon2 in *rpoC1*. We amplified and sequenced the region. The exon/intron of *rpoC1* was annotated by DOGMA, and the *rpoC1* partial sequences were aligned with 14 full fern *rpoC1* sequences by MUSCLE alignment program (Edgar, 2004). We also amplified and sequenced the region for *L. japonicum* as a reference, even though the species was subjected to the completed sequencing of the cp genome.

## RESULTS

### Phylogenetic analysis of ferns and related groups


The aligned sequences of 89 cp genes from 35 taxa consisted of 94,790 bp. Among them, 31,312 sites (33.0%) were constant, 10,740 sites (11.3%) were parsimony-uninformative, and 52,738



**Table 3.** Potential and detected RNA editing sites in chloroplast genome of ferns

Group	Taxon	No. of alternative start codons	No. of genes with internal stop codons	Maximum no. of internal stop codons in gene	No. of RNA editing sites <sup>a</sup>
Core leptosporangiate ferns	<i>Adiantum capillus-veneris</i>	25	18	4	349
	<i>Cheilanthes lindheimeri</i>	26	22	4	-
	<i>Pteridium aquilinum</i> subsp. <i>Aquilinum</i>	29	25	8	-
	<i>Alsophila spinulosa</i>	22	30	10	-
	<i>Marsilea crenata</i>	28	21	3	-
Early diverged leptosporangiate ferns	<i>Lygodium japonicum</i> *	21	17	2	-
	<i>Diplazium glaucum</i> *	19	33	15	-
	<i>Osmunda cinnamomea</i> *	5	0	0	-
Eusporangiate ferns	<i>Angiopteris evecta</i>	1	0	0	-
	<i>Psilotum nudum</i>	0	0	0	-
	<i>Ophioglossum californicum</i>	7	1	1	-
	<i>Mankyua chejuensis</i>	7	3	1	-
	<i>Equisetum arvense</i>	2	0	0	-
	<i>Equisetum hyemale</i>	1	0	0	-

<sup>a</sup>The numbers of RNA editing sites were reported by Wolf et al. (2004).




**Fig. 3.** GC contents and the effective numbers of codons (ENCs). (A) The scatter diagram of the GC1, GC2, and GC3 against all GC contents. The regression lines and the equations are shown on the diagram. First (blue), second (green), and third (red) codon positions, and the taxonomic groups are distinguished by colors and symbols (five different symbols). (B) The boxplot of the GC contents (the upper diagram) and the ENCs (the lower diagram) for taxonomic groups. Seed plant lineages usually show small ranges of variation, while the fern lineages, especially the fern allies, show a wide range of variation both in GC contents and ENCs. The core leptosporangiate ferns show higher ENCs than early diverged leptosporangiate ferns.

### Patterns of GC contents and ENCs in early diverged leptosporangiate ferns

We compared the GC contents of the cp genome coding sequence of bryophytes (8 spp.), Lycopodiopsida (4 spp.), Polypodiopsida (14 spp.), gymnosperms (26 spp.), and angiosperms (142 spp.; Supplementary Table 1). The GC content ranged from 29.5 to 39.2% in bryophytes, from 36.8 to 54.4% in Lycopodiopsida, from 33.6 to 42.4% in Polypodiopsida, from 35.1 to 40.0% in gymnosperms, and from 34.4 to 41.3% in angiosperms. We analyzed the GC content for each codon position and the ENCs using a box plot for each taxonomic group. In the GC position-plot, almost all data were distributed near the regression line, and the slope of GC3 was twice as high as the slope of GC1 and GC2 (Fig. 3A). The GC3 showed wider variation than the GC1 and the GC2 (Fig. 3B). The median value of GC3 showed a little variation among seed plants. However, the range of GC3 in Polypodiopsida showed substantial variation. The GC3 value seemed to increase from eusporangiate ferns to leptosporangiate ferns. The ENCs showed a similar distribution pattern when compare to the GC3 values. The ENCs of seed plants were concentrated between 45 and 50, but the ENCs of Polypodiopsida ranged from 41 to 54 (Fig. 3B).

### Repeat sequences in the cp genomes of early diverged leptosporangiate ferns

The *D. glaucum* cp genome contained more than 100 dispersed repeat sequences. Most of them were located around tRNA genes, especially between *trnL-CAA* and *trnI6*. The cp genome of *D. glaucum* had a 9.7 kb inversion mutation between *trnL-CAA* and *trnV-GAC*. As a result, the position of *trnL-CAA* was moved to the IR region near the *trnI6* gene. The repeat sequences between *trnL-CAA* and *trnI6* usually had a repeating backbone sequence of GGAC-NNNN-AATCC. The sequence was repeated 43 times in the intergenic spacer (IGS) region between *trnL-CAA* and *trnI6*, and 31 of them were AGGAC-NNN-AAATCCT. The 15 bp sequence was similar to the tRNA anticodon loop sequences of *trnF-GAA* and *trnC-GCA* (Fig. 4). The first 5 bp and the last 7 bp in the sequence



**Fig. 4.** The IGS region between *tmL* and *rm16* in *D. glaucum*. A 15 bp sequence is repeated 31 times, and the repeating unit is similar to the anticodon loop sequences of tRNA. The number in the circles indicates the number of duplications in each repeating unit.

were conserved, and the middle 3 bp sequence were variable. The IGS region between *tmL-CAA* and *ndhB* also contained many repeat sequences. TCNATGTAGAAA was repeated 10 times, and GAAATAGTAGGGTTGACATT was repeated four times in this IGS. The *O. cinnamomea* cp genome contained fewer repeat sequences than *D. glaucum*. GGAC-NNNN-AAATCC was detected 11 times, with six of them located between *tmL-CAA* and *ycf2*, but AGGAC-NNN-AAATCCT was not repeated in *D. glaucum*. The *L. japonicum* cp genome contained the smallest number of repeat sequence. The GGAC-NNNN-AAATCC sequence was repeated six times, with four of the repeats being located between the *tmR-ACG* and *ndhB* genes. AGGACNNN-AAATCCT was repeated four times and three of them located between the *tmR-ACG* and *ndhB* genes. Tandem repeats over 20 bp in length were also detected in all three investigated species. The *D. glaucum* and *L. japonicum* cp genomes contained only AT dinucleotide repeats, while the *O. cinnamomea* cp genome contained di-, hexa-, 7-, 12-, and 17-nucleotide repeats.

#### Intraspecific cp genome difference of *L. japonicum* and *rpoC1* intron loss in the genus *Lygodium*

The *L. japonicum* cp genome from a Korean population was 118 bp shorter than the Chinese population (KC 536645). The difference between two populations were due to tandem re-


peats between *psbM* and *tmD-GUC* in LSC and between *ndhB* and *tmR-ACG* in IR. A total of 19 single nucleotide polymorphisms (SNPs) were detected between the two populations. In addition, one small inversion was detected between *rpl32* and *tmP-GGG*.

*RpoC1* intron loss was described in the *L. japonicum* cp genome (Gao et al., 2013). We extended the intron survey of *rpoC1* to other species of *Lygodium* (Schizaeales) and *Vandenboschia* (Hymenophyllales) using PCR and sequencing methods. The amplified *rpoC1* sequences from five species of the genus *Lygodium* were about 960 bp, but it was over 1.5 kb in *Vandenboschia*. The amplified region included a whole *rpoC1* intron and the 5' end of *rpoC1* exon 2. We generated the sequences from all amplified products. However, background noise and double peaks at both ends were present in three sequences of the genus *Lygodium*, so these sequences were excluded from further analyses. Therefore, the sequences from *L. japonicum* (China), *L. poystachyum*, and *V. striata* were aligned with the 14 complete cp genome sequences from ferns (Fig. 5). All surveyed species of the genus *Lygodium* lost the *rpoC1* intron, while all other fern lineages have the *rpoC1* intron.


## DISCUSSION

### Phylogenetic relationship of monilophytes

Monilophytes consist of four orders of eusporangiate ferns and a clade of leptosporangiate ferns (Smith et al., 2006). The phylogenetic relationships among the four eusporangiate ferns were uncertain even though most of the recent data indicate they are paraphyletic assemblages. Specifically, the phylogenetic position of Equisetales is largely different among the data sets, and this relationship remains to be resolved (Karol et al., 2010; Pryer et al., 2001). Our phylogenetic tree developed based on the most comprehensive cp genome data so far placed the Equisetales at the most basal position among the members of monilophytes, even though they did not show a large ML value difference (Fig. 1). The branch length leading to the *Equisetum* is relatively long when compared to other monilophyte lineages, and it is comparable with the branch lengths of *Selaginella* and *Welwitschia*. In contrast, several recent molecular phylogenetic analyses based on cp gene sequences placed the Equisetales either as a sister group to marattioid ferns (Pryer et al., 2001; 2004) or as a sister group to Psilotales (Karol et al., 2010). We believe that the tree differences may due to the taxon sampling for the long branched taxa. The *Equisetum* lineages were recognized as a distinct phylum for a long time because of the distinct morphological characteristics of both sporophytes and gametophytes (Bold et al., 1987). Their ancestry shows a long evolutionary history that dates back to the Devonian period, and they flourished during the Carboniferous periods in fossil history (Good and Taylor, 1975; Schweitzer, 1972). The *Equisetum* is the only extant lineages of the group. Addition of more complete cp genome sequences from eusporangiate ferns and basal leptosporangiate ferns will be helpful to resolve the different phylogenetic hypotheses. Our data support the monophyly of leptosporangiate ferns. We added two new complete cp genome sequences from early diverged leptosporangiate ferns: *Ophioglossum* (Ophioglossales) and *Diplopterygium* (Gleicheniales). Osmundales was the first diverged leptosporangiate fern in the tree, and this result concordant to the previous studies (Pryer et al., 2004; Schuettpelz and Pryer, 2007). In addition, Geicheniales was placed between Osmundales and Schizaeales. The addition of two early diverged leptosporangiate ferns helped us to understand not only cp genome evolution, but also the evolution of



**Fig. 5.** Alignments of *rpoC1* gene regions in ferns. We designed a primer set at the end of exon 1 and in the middle of exon 2 (indicated by black arrows) in order to test the presence or the absence of the *rpoC1* intron. The intron is lost only in the genus *Lygodium*. All eusporangiate ferns and all other leptosporangiate ferns contained the intron.



**Fig. 6.** The major evolutionary changes of cp genomes among fern lineages. The gene/intron gains or losses, inversion events, and the anticodon changes were plotted on the abbreviated phylogenetic tree of ferns as shown on Fig. 1A. The evolutionary events were accounted for using the ACCTRAN criteria during the parsimony analysis using MacClade program. The solid and empty circles indicate the presence and the absence of each character states, respectively.



leptosporangiate fern groups. Our phylogenetic analyses also suggest that two alternative positions of Lycophytes, either as a sister group to euphyllophytes or as a sister group to spermatophytes, are not significantly different in terms of ML values and other support values.

### Cp genome structure evolution in early diverged leptosporangiate ferns

Several cp genome structural modifications, including gene/intron loss and inversion, have been reported for various ferns (Gao et al., 2009; 2013; Hasebe and Iwatsuki, 1990; Wolf et al., 2003; 2010). However, most of these studies were focused on the core leptosporangiate ferns. The complete cp DNA sequences from three early diverged leptosporangiate ferns provide us with new information on the evolution of the cp genome and the phylogenetic relationships of ferns. Figure 6 shows the genome evolutionary history on the phylogenetic tree. The coding gene losses occurred mainly for eusporangiate ferns. In contrast, the tRNA gene losses and anticodon substitutions usually occur on non-Osmundalean leptosporangiate ferns. Large inversions among IR-LSC are characteristic of the early diverged leptosporangiate ferns.

The inversion between *tmL*-CAA and *tmV*-GCA in the cp genome of *D. glaucum* is interesting. Due to this inversion, the position of *tmL*-CAA is moved from the LSC to an IR. In addition, many repeat sequences are located in the IGS between *rm16* and *tmL*-CAA. The repeating AGGAC-NNN-AAATCCT backbone sequence makes up the repeating sequences within this region. The repeat sequence has also been detected in *L. japonicum*. This 15 bp backbone sequence was also reported in the IGS between *tmT*-UGU and *tmR*-ACG in *Alsophila spinulosa* (Gao et al., 2009). However, the sequence was not detected in other core leptosporangiate ferns. Therefore, we hypothesize that the 15 bp repeat occurred widely in early diverged leptosporangiate ferns and then the repeat sequences were degraded due to slipped mispairing (Moore, 1983).

*RpoC1* intron loss was detected only from the genus *Lygodium* (Fig. 5) in fern lineages. *RpoC1* intron losses occur in various plant lineages. They are recognized as a synapomorphic characteristic of the subfamily Cactoideae of Cactaceae (Wallace and Cota, 1996) and the tribes Drosanthemeae and Ruschieae of Aizoaceae (Thiede et al., 2007). However, the intron losses occur independently even within a single genus depending on the species (Downie et al., 1996). Similar to *rpoC1* intron losses in angiosperm, the analyses of the *rpoC1* intron loss in ferns will help to indicate some taxonomic grouping. The primer set we developed in this study work well for fern species. So far, we tested the limited samples of ferns and found intron loss only from *Lygodium*. All species of *Lygodium* share this intron loss. This survey needs to expand to more fern species.

The patterns of GC contents by codon position and the ENCs of ferns are different from those of seed plants (Fig. 3). Early diverged ferns show low GC contents and ENCs when compared to the recently originated group. They also show a wide range of variation in GC and ENC values, while seed plants were similar to each other. The difference among groups may be due to the sampling error because many cp genome sequences are reported in seed plants, but only fifteen cp genome sequences are reported in ferns. Nevertheless, the value of GC3 and ENCs are notably different between ferns and seed plants. Furthermore, the GC3 and ENCs values are markedly different between the early diverged leptosporangiate and the core leptosporangiate ferns. We need more information about

the cp genome sequences from ferns in order to address this question properly.

### The molecular characteristics of the *O. cinnamomea* cp genome

Osmundales have several common characteristics with eusporangiate ferns. However, it is normally recognized as a member of leptosporangiate ferns based on other morphological characteristics. However, the cp genome of *Osmunda* is notably different from other leptosporangiate ferns in the following characters: 1) The cp genome structure is similar to that of *E. arvense*, which is eusporangiate fern; 2) the gene/intron losses that occurred in non-Osmundalean leptosporangiate ferns do not occur in the cp genome of *O. cinnamomea*; and 3) RNA editing sites in the cp genome are frequently predicted or detected in leptosporangiate ferns, but almost no RNA editing sites occur in the *O. cinnamomea* cp genome.

Molecular characteristics are frequently used to indicate specific taxonomic groups. The large inversion between *psbM* and *ycf2* is recognized as a synapomorphic characteristic of seed plants and ferns (Raubeson and Jansen, 1992). In addition, a 9-bp insertion in *rps4* is also a monophyletic characteristic of ferns (Pryer et al., 2001). Based on molecular characters, leptosporangiate ferns can be divided into two groups: Osmundalean ferns and non-Osmundalean ferns. Osmundales is also differentiated from other leptosporangiate ferns by echina-bearing tubercles on the surfaces of spores (Tryon and Lugaron, 1991). Therefore, it is reasonable to distinguish non-Osmundalean leptosporangiate ferns from Osmundalean ferns, or vice versa.

### CONCLUSION

The complete cp DNA sequences from three major lineages of basal leptosporangiate ferns provide us a substantial information not only on evolution of the cp genomes and also on the phylogenetics of fern lineages. Our phylogenetic analysis, which was based on the largest numbers of complete cp genomes of Monilophytes so far, showed the paraphyly of the eusporangiate ferns. The Equisetales was the sister group to all other members of monilophytes. The results were consistent for the majority of the other analyses. In contrast to the paraphyly of eusporangiate ferns, the leptosporangiate ferns from a monophyletic group. Within the eusporangiate ferns, the cp genome structures, gene/intron contents, and RNA editing sites of *Osmunda cinnamomea* (Osmundales) were similar to that of the eusporangiate ferns. Therefore, these cp genome characteristics in *Osmunda* are symplesiomorphic conditions. Several lines of morphological and anatomical data, both from sporophyte and gametophyte of *Osmunda*, also support the intermediate or the distinctive natures of Osmundalean ferns from other leptosporangiate ferns. Therefore, the Osmundalean ferns can be recognized as living fossil lineages from leptosporangiate ferns. The GC contents and ENCs in ferns vary significantly when compared to that of seed plants. The both values in the early diverged leptosporangiate ferns showed intermediate levels between eusporangiate and core leptosporangiate ferns. The cp genome of *Diplazium glaucum* (Gleicheniales) has a large unique inversion between the *tmL*-CAA and *tmV*-GCA genes. Several repeated sequences were detected around the inversion break points. The cp genome of *Lygodium japonicum* (Schizaeales) showed *rpoC1* intron loss, which is shared among all *Lygodium* species.

Note: Supplementary information is available on the *Molecules and Cells* website ([www.molcells.org](http://www.molcells.org)).

## ACKNOWLEDGMENTS

This research was supported by a research grant (#KRF-2010-0011796) from the Korea Research Foundation and by the Eco-Innovation project of the Ministry of Environment (416-111-007) from KEITI to Ki-Joong Kim. A Ph.D. fellowship to HTK was granted by the graduate student education program from NIBR (2012-2013). DNA materials used in this study were deposited in and are available from the Plant DNA Bank of Korea (PDBK).

## REFERENCES

- Bold, H.C., Alexopoulos, C.J., and Delevoryas, T. (1987). *Morphology of Plants and Fungi*, 4th ed. (New York: Harper and Row).
- Chumley, T.W., Palmer, J.D., Mower, J.P., Fourcade, H.M., Calie, P.J., Boore, J.L., and Jansen, R.K. (2006). The complete chloroplast genome sequence of *Pelargonium x hortorum*: organization and evolution of the largest and most highly rearranged chloroplast genome of land plants. *Mol. Biol. Evol.* **23**, 2175-2190.
- Cross, G.L. (1931a). Embryology of *Osmunda cinnamomea*. *Bot. Gaz.* **92**, 210-217.
- Cross, G.L. (1931b). Meristem in *Osmunda cinnamomea*. *Bot. Gaz.* **91**, 65-76.
- Downie, S.R., Llanas, E., and Katz-Downie, D.S. (1996). Multiple independent losses of the *rpoC1* intron in angiosperm chloroplast DNA's. *Syst. Bot.* **21**, 135-151.
- Doyle, J.J., and Dolye, J.L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* **19**, 11-15.
- Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **32**, 1792-1797.
- Freeberg, J.A., and Gifford Jr, E.M. (1984). The root apical meristem of *Osmunda regalis*. *Am. J. Bot.* **71**, 558-563.
- Gao, L., Yi, X., Yang, Y.X., Su, Y.J., and Wang, T. (2009). Complete chloroplast genome sequence of a tree fern *Alsophila spinulosa*: insights into evolutionary changes in fern chloroplast genomes. *BMC Evol. Biol.* **9**, 130. doi:10.1186/1471-2148-9-130
- Gao, L., Wang, B., Wang, Z.W., Zhou, Y., Su, Y.J., and Wang, T. (2013). Plastome sequences of *Lygodium japonicum* and *Marsilea crenata* reveal the genome organization transformation from basal ferns to core leptosporangiates. *Genome Biol. Evol.* **5**, 1403-1407.
- Gifford Jr, E. (1983). Concept of apical cells in bryophytes and pteridophytes. *Annu. Rev. Plant Physiol.* **34**, 419-440.
- Good, C.W., and Taylor, T.N. (1975). The morphology and systematic position of calamitean elater-bearing spores. *Geosci. Man* **11**, 133-139.
- Grewe, F., Guo, W., Gubbels, E.A., Hansen, A.K., and Mower, J.P. (2013). Complete plastid genomes from *Ophioglossum californicum*, *Psilotum nudum*, and *Equisetum hyemale* reveal an ancestral land plant genome structure and resolve the position of Equisetales among monilophytes. *BMC Evol. Biol.* **13**, 8.
- Guisinger, M.M., Kuehl, J.V., Boore, J.L., and Jansen, R.K. (2011). Extreme reconfiguration of plastid genomes in the angiosperm family Geraniaceae: rearrangements, repeats, and codon usage. *Mol. Biol. Evol.* **28**, 583-600.
- Hasebe, M., and Iwatsuki, K. (1990). Chloroplast DNA from *Adiantum capillus-veneris* L., a fern species (Adiantaceae); clone bank, physical map and unusual gene localization in comparison with angiosperm chloroplast DNA. *Curr. Genet.* **17**, 359-364.
- Hiratsuka, J., Shimada, H., Whittier, R., Ishibashi, T., Sakamoto, M., Mori, M., Kondo, C., Honji, Y., Sun, C.R., Meng, B.Y., et al. (1989). The complete sequence of the rice (*Oryza sativa*) chloroplast genome: intermolecular recombination between distinct tRNA genes accounts for a major plastid DNA inversion during the evolution of the cereals. *Mol. Gen. Genet.* **217**, 185-194.
- Iwatsuki, K., Yamazaki, T., Boufford, D., and Ohba, H. (1995). *Flora of Japan*. Vol. I, Pteridophyta and Gymnospermae (Tokyo: Kodansha).
- Jansen, R.K., Raubeson, L.A., Boore, J.L., dePamphilis, C.W., Chumley, T.W., Haberle, R.C., Wyman, S.K., Alverson, A.J., Peery, R., Herman, S.J., et al. (2005). Methods for obtaining and analyzing whole chloroplast genome sequences. *Methods Enzymol.* **395**, 348-384.
- Jansen, R.K., Cai, Z., Raubeson, L.A., Daniell, H., Depamphilis, C.W., Leebens-Mack, J., Muller, K.F., Guisinger-Bellian, M., Haberle, R.C., Hansen, A.K., et al. (2007). Analysis of 81 genes from 64 plastid genomes resolves relationships in angiosperms and identifies genome-scale evolutionary patterns. *Proc. Natl. Acad. Sci. USA* **104**, 19369-19374.
- Karol, K.G., Arumuganathan, K., Boore, J.L., Duffy, A.M., Everett, K.D., Hall, J.D., Hansen, S.K., Kuehl, J.V., Mandoli, D.F., Mishler, B.D., et al. (2010). Complete plastome sequences of *Equisetum arvense* and *Isoetes flaccida*: implications for phylogeny and plastid genome evolution of early land plant lineages. *BMC Evol. Biol.* **10**, 321.
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., et al. (2012). Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647-1649.
- Kim, K.J., and Lee, H.L. (2004). Complete chloroplast genome sequences from Korean ginseng (*Panax schinseng* Nees) and comparative analysis of sequence evolution among 17 vascular plants. *DNA Res.* **11**, 247-261.
- Kim, Y.K., Park, C.W., and Kim, K.J. (2009). Complete chloroplast DNA sequence from a Korean endemic genus, *Megaleranthis saniculifolia*, and its evolutionary implications. *Mol. Cells* **27**, 365-381.
- Kurtz, S., Choudhuri, J.V., Ohlebusch, E., Schleiermacher, C., Stoye, J., and Giegerich, R. (2001). REPuter: the manifold applications of repeat analysis on a genomic scale. *Nucleic Acids Res.* **29**, 4633-4642.
- Lowe, T.M., and Eddy, S.R. (1997). tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* **25**, 955-964.
- Metzgar, J.S., Skog, J.E., Zimmer, E.A., and Pryer, K.M. (2008). The paraphyly of *Osmunda* is confirmed by phylogenetic analyses of seven plastid loci. *Syst. Bot.* **33**, 31-36.
- Miller, M.A., Pfeiffer, W., and Schwartz, T. (2010). Creating the CIPRES Science Gateway for inference of large phylogenetic trees; in Gateway Computing Environments Workshop (GCE), 2010 (IEEE).
- Moore, G.P. (1983). Slipped-mispairing and the evolution of introns. *Trends Biochem. Sci.* **8**, 411-414.
- Palmer, J.D. (1986). Isolation and structural analysis of chloroplast DNA. *Methods Enzymol.* **118**, 167-186.
- Palmer, J.D., and Stein, D.B. (1982). Chloroplast DNA from the fern *Osmunda cinnamomea*: physical organization, gene localization and comparison to angiosperm. *Curr. Genet.* **5**, 165-170.
- Palmer, J.D., and Stein, D.B. (1986). Conservation of chloroplast genome structure among vascular plants. *Curr. Genet.* **10**, 823-833.
- Posada, D., and Crandall, K.A. (1998). MODELTEST: testing the model of DNA substitution. *Bioinformatics* **14**, 817-818.
- Pryer, K.M., Schneider, H., Smith, A.R., Cranfill, R., Wolf, P.G., Hunt, J.S., and Sipes, S.D. (2001). Horsetails and ferns are a monophyletic group and the closest living relatives to seed plants. *Nature* **409**, 618-622.
- Pryer, K.M., Schuettpelz, E., Wolf, P.G., Schneider, H., Smith, A.R., and Cranfill, R. (2004). Phylogeny and evolution of ferns (monilophytes) with a focus on the early leptosporangiate divergences. *Am. J. Bot.* **91**, 1582-1598.
- Pryer, K.M., Smith, A.R., and Rothfels, C. (2009). Polypodiopsida Cronquist, Takht. & Zimmerm. 1966. *Ferns*. Version 14, January 2009 (under construction). <http://tolweb.org/Polypodiopsida/20615/2009.01.14> in The Tree of Life Web Project, <http://tolweb.org/>.
- Raubeson, L.A., and Jansen, R.K. (1992). Chloroplast DNA evidence on the ancient evolutionary split in vascular land plants. *Science* **255**, 1697-1699.
- Sablok, G., Nayak, K.C., Vazquez, F., and Tatarinova, T.V. (2011). Synonymous codon usage, GC(3), and evolutionary patterns across plastomes of three pooid model species: emerging grass genome models for monocots. *Mol. Biotechnol.* **49**, 116-128.
- Schneider, H., Schuettpelz, E., Pryer, K.M., Cranfill, R., Magallon,

- S., and Lupia, R. (2004). Ferns diversified in the shadow of angiosperms. *Nature* 428, 553-557.
- Schuettpelz, E., and Pryer, K.M. (2007). Fern phylogeny inferred from 400 leptosporangiate species and three plastid genes. *Taxon* 56, 1037-1050.
- Schweitzer, H.-J. (1972). Die Mitteldevon-Flora von Lindlar (Rheinland). 3. Filicinae-Hyenia elegans Kräusel & Weyland. *Palaeontographica Abteilung B*, 154-175.
- Smith, D.R. (2009). Unparalleled GC content in the plastid DNA of *Selaginella*. *Plant Mol. Biol.* 71, 627-639.
- Smith, A.R., Pryer, K.M., Schuettpelz, E., Korall, P., Schneider, H., and Wolf, P.G. (2006). A classification for extant ferns. *Taxon* 55, 705-731.
- Stamatakis, A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22, 2688-2690.
- Stamatakis, A., Hoover, P., and Rougemont, J. (2008). A rapid bootstrap algorithm for the RAxML web servers. *Syst. Biol.* 57, 758-771.
- Swofford, D.L. (2003). PAUP; Phylogenetic Analysis Using Parsimony, version 4.0 b10 (Sunderland, Massachusetts: Sinauer Associates).
- Tangphatsornruang, S., Sangsrakru, D., Chanprasert, J., Uthapaisanwong, P., Yoocha, T., Jomchai, N., and Tragoonrung, S. (2010). The chloroplast genome sequence of mungbean (*Vigna radiata*) determined by high-throughput pyrosequencing: structural organization and phylogenetic relationships. *DNA Res.* 17, 11-22.
- Taylor, E.L., Taylor, T.N., and Krings, M. (2009). *Paleobotany: The Biology and Evolution of Fossil Plants* (Burlington, Massachusetts: Elsevier Ltd.).
- Thiede, J., Schmidt, S.A., and Rudolph, B. (2007). Phylogenetic implication of the chloroplast *rpoC1* intron loss in the Aizoaceae (Caryophyllales). *Biochem. Syst. Ecol.* 35, 372-380.
- Tidwell, W.D., and Ash, S.R. (1994). A review of selected Triassic to Early Cretaceous ferns. *J. Plant Res.* 107, 417-442.
- Tryon, A., and Lugardon, B. (1991). *Spores of the Pteridophyta*. (New York: Springer-Verlag).
- Vetrivel, U., Arunkumar, V., and Dorairaj, S. (2007). ACUA: a software tool for automated codon usage analysis. *Bioinformation* 2, 62-63.
- Wallace, R.S., and Cota, J.H. (1996). An intron loss in the chloroplast gene *rpoC1* supports a monophyletic origin for the subfamily Cactoideae of the Cactaceae. *Curr. Genet.* 29, 275-281.
- Wolf, P.G., Pryer, K.M., Ueda, K., Ito, M., Sano, R., Gastony, G., Yokoyama, J., Manhart, J., Murakami, N., and Crane, E. (1995). Fern phylogeny based on *rbcL* nucleotide sequences. *Am. Fern J.* 85, 134-181.
- Wolf, P.G., Rowe, C.A., Sinclair, R.B., and Hasebe, M. (2003). Complete nucleotide sequence of the chloroplast genome from a leptosporangiate fern, *Adiantum capillus-veneris* L. *DNA Res.* 10, 59-65.
- Wolf, P.G., Rowe, C.A., and Hasebe, M. (2004). High levels of RNA editing in a vascular plant chloroplast genome: analysis of transcripts from the fern *Adiantum capillus-veneris*. *Gene* 339, 89-97.
- Wolf, P.G., Roper, J.M., and Duffy, A.M. (2010). The evolution of chloroplast genome structure in ferns. *Genome* 53, 731-738.
- Wolf, P.G., Der, J.P., Duffy, A.M., Davidson, J.B., Grusz, A.L., and Pryer, K.M. (2011). The evolution of chloroplast genes and genomes in ferns. *Plant Mol. Biol.* 76, 251-261.
- Wright, F. (1990). The 'effective number of codons' used in a gene. *Gene* 87, 23-29.
- Wu, C.S., Wang, Y.N., Liu, S.M., and Chaw, S.M. (2007). Chloroplast genome (cpDNA) of *Cycas taitungensis* and 56 cp protein-coding genes of *Gnetum parvifolium*: insights into cpDNA evolution and phylogeny of extant seed plants. *Mol. Biol. Evol.* 24, 1366-1379.
- Wyman, S.K., Jansen, R.K., and Boore, J.L. (2004). Automatic annotation of organellar genomes with DOGMA. *Bioinformatics* 20, 3252-3255.
- Yatabe, Y., Nishida, H., and Murakami, N. (1999). Phylogeny of Osmundaceae inferred from *rbcL* nucleotide sequences and comparison to the fossil evidences. *J. Plant Res.* 112, 397-404.
- Yi, D.K., and Kim, K.J. (2012). Complete chloroplast genome sequences of important oilseed crop *Sesamum indicum* L. *PLoS One* 7, e35872.
- Yi, D.K., Lee, H.L., Sun, B.Y., Chung, M.Y., and Kim, K.J. (2012). The complete chloroplast DNA sequence of *Eleutherococcus senticosus* (Araliaceae); comparative evolutionary analyses with other three asterids. *Mol. Cells* 33, 497-508.