



Effects of a Chelated Copper as Growth Promoter on Performance and Carcass Traits in Pigs

J. Zhao*, G. Allee¹, G. Gerlemann¹, L. Ma¹, M. I. Gracia², D. Parker,
M. Vazquez-Anon, and R. J. Harrell³

Novus International Inc, Saint Charles, MO, USA 63304

ABSTRACT: Three studies were conducted to investigate whether a chelated Cu can replace CuSO₄ as a growth promoter in pigs. In Exp. 1, a total of 240 piglets (Large White×Landrace, 7.36±0.10 kg) were randomly allocated to 1 of 3 treatments with 8 replicates and 10 piglets per pen. Treatments included a NRC control (CuSO₄, 6 mg/kg), two Cu supplementations from either CuSO₄ or Cu(HMTBa)₂ at 170 mg/kg. Pigs fed Cu(HMTBa)₂ were 6.0% heavier than pigs fed either the NRC control or 170 mg/kg CuSO₄ (p = 0.03) at the end of the experiment. During the 42 days of experimental period, pigs fed Cu(HMTBa)₂ gained 9.0% more (p = 0.01), tended to eat more feed (p = 0.09), and had better feed efficiency (p = 0.06) than those fed CuSO₄. Compared with the 6 mg/kg CuSO₄ NRC control, liver Cu was increased 2.7 times with 170 mg/kg CuSO₄ supplementation, and was further increased with Cu(HMTBa)₂ (4.5 times, p<0.05). In Exp. 2, a total of 616 crossbred piglets (PIC, 5.01±0.25 kg) were randomly allocated to 1 of 4 treatments with 7 replicates and 22 piglets per pen. Treatments included a NRC control (from CuSO₄), and three pharmaceutical levels of Cu (150 mg/kg) supplemented either from CuSO₄, tri-basic copper chloride (Cu₂[OH]₃Cl), or Cu(HMTBa)₂. Pigs fed CuSO₄ or Cu(HMTBa)₂ had better feed efficiency (p = 0.01) and tended to gain more (p = 0.08) compared with those fed the NRC control. Pigs fed Cu₂(OH)₃Cl were intermediate. Pigs fed Cu(HMTBa)₂ had the highest liver Cu, which was significantly higher than those fed (Cu₂[OH]₃Cl) or the negative control (p = 0.01). In Exp. 3, a total of 1,048 pigs (PIC, 32.36±0.29 kg) were allotted to 6 treatments with 8 replicates per treatment and 20 to 22 pigs per pen. The treatments included a NRC control with 4 mg/kg Cu from CuSO₄, a positive control with 160 mg/kg Cu from CuSO₄, and incremental levels of Cu(HMTBa)₂ at 20, 40, 80, and 160 mg/kg. During the overall experimental period of 100 days, no benefit from 160 mg/kg CuSO₄ was observed. Pigs fed Cu(HMTBa)₂ had increased ADG (linear and quadratic, p≤0.05) and feed efficiency (linear and quadratic, p≤0.05) up to 80 mg/kg and no further improvement was observed at 160 mg/kg for the whole experimental period. Pigs fed 80 mg/kg Cu(HMTBa)₂ weighed 1.8 kg more (p = 0.07) and were 2.3 kg heavier in carcass (p<0.01) compared with pigs fed 160 mg/kg CuSO₄. In addition, loin depth was increased with increased Cu(HMTBa)₂ supplementation with pigs fed 80 mg/kg Cu(HMTBa)₂ had the greatest loin depth (p<0.05). In summary, Cu(HMTBa)₂ can be used to replace high CuSO₄ as a growth promoter in nursery and grower-finisher pigs. (**Key Words:** Chelated Trace Mineral, Cu(HMTBa)₂, Copper, Growth Promoter, Swine)

INTRODUCTION

Requirement of Cu as a nutrient is low and NRC (1998) recommends 3 to 6 mg/kg Cu for nursery and grower-

finisher pigs. In practice, pharmaceutical Cu from CuSO₄ (125 to 250 mg/kg) is routinely added in nursery pigs as a growth promoter and its benefits on feed intake and weight gain have been well documented (Bunch et al., 1961; Hawbaker et al., 1961). However, high dietary Cu presents an environmental concern when excess Cu is excreted in feces (Kornegay and Harper, 1997). Roof and Mahan (1982) reported a 14-fold increase of fecal Cu excretion with 250 mg/kg CuSO₄ supplementation compared with the non-supplemented control. Accumulation of Cu and Zn in the soil has been implicated to reduce crop yield (Tucker,

* Corresponding Author: Junmei Zhao. 20 Research Park Dr., Saint Charles, MO USA 63304. Tel: +1-314-8257211, E-mail: zhaojunmei@hotmail.com

¹ University of Missouri, Columbia, MO, USA 65211.

² Imasde Agroalimentaria, Madrid, Spain 28224.

³ Carthage Veterinary Service, Carthage, IL USA 62321.

Submitted Jul. 13, 2013; Revised Sept. 25, 2013; Accepted Nov. 27, 2013

1997) and pose an environmental threat. In addition, it has been demonstrated that high dietary Cu from inorganic sources antagonizes other nutrients utilization, such as Zn (Zhao et al., 2008) and P (Banks et al., 2004). Due to the negative impact of high dietary CuSO₄, the Commission of the European Communities regulate maximum allowed total Cu in feed with 170 mg/kg in piglets up to 12 weeks of age and 25 mg/kg in all other pigs. Other countries are following with similar regulations.

The mode of action of high Cu might due to its antibacterial function in the lumen (Shurson et al., 1990) directly or indirectly through absorption from lumen to liver, secretion to the bile, and then back to the lumen loop. This theory is partially supported by Zhou et al. (1994), whose research indicated that intravenous injection of copper histidinate, which bypass the lumen, had similar growth promoting effect as dietary supplementation. Compared with inorganic trace minerals, chelated trace minerals are more bioavailable, likely due to reduced incidence of antagonism with other dietary constituents in the gastrointestinal tract (Wang et al., 2007; Zhao et al., 2010). Our previous study suggested that Cu(HMTBa)₂ is better at avoiding nutrient antagonism and less is needed to achieve similar or better performance compared with inorganic minerals in broilers (Zhao et al., 2010) and nursery pigs (Zhao et al., 2009). The objective of these studies was to evaluate a chelated Cu (Cu[HMTBa]₂) as a growth promoter on performance, liver Cu and carcass traits in nursery and grower-finisher pigs.

MATERIALS AND METHODS

Exp. 1 was conducted in Segovia, Spain and met appropriate current quality standards and guidance of the European Food Safety Authority (EFSA). Exp. 2 and 3 were conducted at the University of Missouri (Columbia, MO). All animal procedures were approved by the University of Missouri Animal Care and Use Committee. Cu(HMTBa)₂ (Mintrex Cu, Novus International Inc., St Charles, MO) is a chelate of one Cu atom coordinated with two molecules of 2-hydroxy-4-methylthio butanoic acid (HMTBa). Methionine activity from Cu(HMTBa)₂ (80% Met activity,

Yi et al., 2007) was accounted for and all diets were adjusted to be equal in methionine. Tribasic Cu chloride (Cu₂[OH]₃Cl) is a more concentrated form of Cu (58%), less soluble in water (<1%), and is viewed as a less destructive form of Cu when combined with vitamins.

Exp. 1: Benefits of Cu(HMTBa)₂ as a growth promoter in nursery pigs

A total of 240 crossbred piglets (Large White×Landrace, 26 d of age, 7.36±0.10 kg) were randomly allocated to 1 of 3 experimental treatments based on sex, litter of origin, and BW with 8 replicates for each treatment and 10 piglets per pen. All pigs were fed a common creep feed from 21 to 25 days of age, then initiated treatments at 26 days of age. The treatments included a NRC control (Cu 6 mg/kg, Mn 4 mg/kg, and Zn 100 mg/kg for phase I; Cu 5 mg/kg, Mn 3 mg/kg, and Zn 80 mg/kg for phase II from sulfates), as well as the two treatments at European maximum level (total Cu 170 mg/kg, Mn and Zn at 150 mg/kg from sulfates) with added Cu came either from CuSO₄ or Cu(HMTBa)₂. Table 1 listed detailed treatment descriptions. The nursery rooms were lit by a combination of daylight (through skylights) and artificial light (non-programmable). The pens were slatted and ventilation was achieved by single, variable-speed fans linked to temperature sensors. Diets were provided *ad libitum* throughout the trial from hoppers and water from two nipple drinkers per pen.

Typical European mixed-grain diets (barley, wheat, corn, and soybean meal) were formulated to meet or exceed NRC (1998) nutrient recommendation for piglets (Table 2). Pigs were fed two phase diets (d 0 to 14 and d 15 to 42). No veterinary antibiotics or veterinary zinc oxide was included in any diets. Diets were analyzed for Cu content using atomic absorption spectrometry (AOAC 968.08), ash by incineration (AOAC 942.05), protein by Kjeldahl (AOAC 984.13), and ether extract by Soxhlet fat analysis (AOAC 920.39) as described by the AOAC (2000). Weight gains, feed intakes and feed conversion were determined at the end of each feeding period. At the end of the experiment, one medium-size piglet per pen was slaughtered and tissue sampled (muscle, liver, kidney, skin and fat, and tibia bone)

Table 1. Treatment description (Exp. 1)¹

Trt	Phase I (d 0 to 14)	Phase II (d 15 to 42)
1	NRC control (CuSO ₄ at 6 mg/kg)	Cu 6 mg/kg, Mn 4 mg/kg, Zn 100 mg/kg from sulfates
2	CuSO ₄ at 170 mg/kg	Cu 5 mg/kg, Mn 3 mg/kg, Zn 80 mg/kg from sulfates
3	Cu(HMTBa) ₂ at 170 mg/kg ²	Cu 170 mg/kg, Mn 150 mg/kg, Zn 150 mg/kg from sulfates
		Same as phase I
		Same as phase I

¹ Analyzed Cu using atomic absorption spectrometry: 15, 149, and 169 mg/kg for treatment 1 to 3 in phase I, and 11, 145, and 160 mg/kg in phase II, respectively.

² Chelate of one Cu atom coordinated with 2 molecules of 2-hydroxy-4-methylthio butanoic acid (HMTBa).

Table 2. Basal diet composition and nutrient profile (Exp. 1)

Ingredients (%)	Phase I (d 0 to 14)	Phase II (d 15 to 42)
Barley	9.70	37.00
Cooked wheat	30.25	-
Wheat	-	4.22
Cooked corn	16.50	-
Corn	-	27.35
Extruded soybean	15.78	8.00
Soybean meal	-	11.90
Dried whey	11.70	-
Soy protein	12.00	5.00
Soybean oil	0.637	2.289
MHA ¹	0.313	0.313
L-lysine-50	0.751	0.759
L-threonine	0.186	0.203
L-tryptophan	0.034	0.030
Calcium carbonate	0.429	0.783
Dicalcium phosphate	0.890	1.298
Sodium chloride	0.400	0.425
Vitamin and mineral premix ²	0.40	0.40
Mineral sources ³	0.03	0.03
Analyzed chemical composition (as-fed basis)		
CP (%)	18.5	17.3
Ether extract (%)	5.8	6.6
Ash (%)	5.9	5.4
Calculated nutrients ⁴		
NE (kcal/kg)	2,470	2,450
Ca (%)	0.70	0.75
Available P (%)	0.40	0.36
Standardized ileal digestible Lys (%)	1.25	1.15
TSAA/Lys	64	66
Thr/Lys	65	65

¹ MHA is a calcium salt of 2-hydroxy-4-methylthio butanoic acid.

² Provided per kg of diet: vitamin A 10,000 IU; vitamin D 2,000 IU; vitamin E (α-tocopherol) 100 mg; vitamin K₃ 5 mg; vitamin B₁ 1.5 mg; vitamin B₂ 5 mg; vitamin B₆ 4 mg; vitamin B₁₂ 30 µg; nicotinic acid 25 mg; pantothenic acid 20 mg; folic acid 1 mg; biotin 100 µg; choline chloride 200 mg; Co 0.50 mg; Fe 100 mg; I 2.00 mg; Se 0.20 mg.

³ Supplemented Zn, Mn, and Cu from either sulfates or Cu(HMTBa)₂ following experimental design.

⁴ Based on FEDNA (2003) values for feed ingredients.

for Cu deposition analysis (AOAC 968.08).

Exp. 2: Comparison of different Cu sources as growth promoter in nursery pigs

A total of 616 crossbred piglets (PIC, 20 days of age, 5.02±0.25 kg) were randomly allocated to 4 experimental treatments based on sex and BW with 7 replicates for each treatment and 22 piglets per pen. The treatments included a NRC negative control (6 mg/kg Cu from CuSO₄), and 3 treatments supplemented with different Cu sources (CuSO₄,

Cu₂[OH]₃Cl, and Cu[HMTBa]₂) at 150 mg/kg. The pens were slatted and ventilation was achieved by single, variable-speed fans linked to temperature sensors. Diets were provided *ad libitum* throughout the trial from hoppers and water from 2 nipple drinkers per pen.

Typical North American commercial nursery diets (corn-soybean meal) were formulated to meet or exceed NRC (1998) requirement (Table 3). High Zn (3,000 mg/kg) from zinc oxide was included during the first seven days post-weaning as typical commercial practice. Diets were analyzed for Cu content using atomic absorption spectrometry (AOAC 968.08). Weight gain, feed intake and feed efficiency were determined for the whole experimental period (21 days post-weaning). At the end of the trial, one medium-size piglet per pen was slaughtered for liver mineral analysis.

Table 3. Basal diet formula and nutrient profile (Exp. 2)

Ingredients (%)	Phase I (d 0 to 7)	Phase II (d 7 to 21)
Corn	38.38	46.26
Soybean meal	15.0	25.0
Dairy-Lac 80	24.4	12.2
Poultry meal	7.38	6.16
Fish meal	5.50	3.00
Spray-dried plasma protein	3.0	-
Choice white grease	4.0	4.0
Mono calcium phosphate	-	0.91
Limestone	0.11	0.29
L-lysine-HCL, 78%	0.30	0.30
Threonine	0.16	0.14
MHA ¹	0.26	0.23
Salt	0.40	0.40
ZnO	0.42	0.42
Vitamin premix ²	0.25	0.25
Mineral premix ³	0.15	0.15
Antibiotics	0.29	0.29
Analyzed chemical composition (as-fed basis)		
CP (%)	22.06	21.64
Ether extract (%)	7.39	7.32
Ash (%)	7.53	6.95
Calculated nutrients		
ME (kcal/kg)	3,256	3,217
Ca (%)	0.80	0.80
Available P (%)	0.47	0.45
Standardized ileal digestible Lys (%)	1.40	1.35
TSAA/Lys	60	60
Thr/Lys	65	65

¹ MHA is a calcium salt of 2-hydroxy-4-methylthio butanoic acid.

^{2,3} Provide per kg of final diet: vitamin A 10,000 IU; vitamin D 8,800 IU; vitamin E 220 IU; vitamin K 22 mg; vitamin B₁₂ 19.25 mg; niacin 250 mg; pantothenic acid 137 mg; riboflavin 41.25 mg; I 0.26 mg; Fe 140 mg; Mn 30 mg; Se 0.25 mg; and Zn 140 mg.

Exp. 3: Benefits of Cu(HMTBa)₂ on growth performance and carcass traits in grower-finisher pigs

The third experiment was designed to investigate whether Cu(HMTBa)₂ can replace high dietary CuSO₄ as growth promoter in grower-finisher pigs. A total of 1,048 pigs (PIC C22, 32.36±0.29 kg) were allotted to 1 of 6 dietary treatments in a randomized complete block design with 8 pens per treatment and 20 to 22 pigs per pen based on BW and gender. Gilts and barrows were housed separately in an environmentally controlled room with access to feed and water *ad libitum*. All pigs were fed a Cu deficient diet for the first 21 days then switched to experimental diets. The treatments included a basal control with 4 mg/kg Cu from CuSO₄, a positive control with 160 mg/kg Cu from CuSO₄, and incremental levels of Cu(HMTBa)₂ at 20, 40, 80, or 160 mg/kg. Different Cu(HMTBa)₂ doses were mixed by blending the highest level (160 mg/kg) with the basal to make targeted doses in the final diets. Pigs were fed five phases diets of d 0 to 21 (30 to 50 kg), d 22 to 42 (50 to 70 kg), d 43 to 62 (70 to 90 kg), d 63 to 80 (90 to 100 kg), and d 81 to 100 (100 to 120 kg). Corn-soybean meal-dried distillers grains with solubles (DDGs) diets were made to satisfy NRC (1998) requirements (Table 4). Ractopamine (4.5 mg/kg) was used in the last 28 days before marketing. Pen weight and feed intake were collected for each feeding phase and feed

efficiency was calculated and adjusted for mortality. At the end of the study, all pigs were harvested at the same time at a Missouri process plant and carcass traits were recorded on a pen basis.

Statistical analyses

Data were analyzed as a randomized completely block design ANOVA using GLM of SAS 9.0 (SAS Institute, 2002). In experiment three, linear and quadratic effects of Cu(HMTBa)₂ were analyzed. Pen was the experimental unit. Statistical significance was declared at $p \leq 0.05$, with $0.05 < p \leq 0.10$ considered as a near-significant trend.

RESULTS

Analyzed dietary Cu level

Analyzed dietary Cu levels in Exp. 1 were 15, 149, and 169 mg/kg in phase I and 11, 145, and 160 mg/kg in phase II, which were similar to the targeted levels of 6, 170, and 170 mg/kg added Cu, respectively. In Exp. 2, the analyzed Cu levels were 19, 117, 162, and 208 mg/kg in phase I and 14, 154, 140, and 188 mg/kg in phase II, respectively. Dietary Cu was not analyzed in Exp. 3.

Exp. 1

At the end of the study, pigs fed Cu(HMTBa)₂ were

Table 4. Basal diets composition and nutrient profile (Exp. 3)

Ingredients (%)	Phase I	Phase II	Phase III	Phase IV	Phase V
Corn	57.91	63.88	67.65	77.43	69.77
Dried distillers grain with solubles	20.00	20.00	20.00	10.00	16.68
Soybean meal	17.83	11.86	8.12	8.00	10.00
Choice white grease	1.93	1.93	1.93	2.19	1.00
Limestone	1.06	1.09	1.10	0.90	0.87
Sodium chloride	0.50	0.50	0.50	0.50	0.50
L-lysine-HCL, 78%	0.40	0.40	0.38	0.32	0.40
Threonine	0.11	0.09	0.07	0.06	0.12
Alimet ¹	0.07	0.03	0.00	0.00	0.07
Mono calcium phosphate	0.04	0.07	0.10	0.45	0.40
Mineral and vitamin premix ²	0.13	0.13	0.13	0.13	0.13
Phytase	0.02	0.02	0.02	0.02	0.02
Ractopamine hydrochloride					0.04
Calculated nutrient profile					
ME (kcal/kg)	3,400	3,400	3,400	3,428	3,374
CP (%)	19.30	16.85	15.34	13.34	17.00
Ca (%)	0.5	0.5	0.5	0.5	0.5
Available P (%)	0.3	0.2	0.2	0.2	0.2
Standardized ileal digestible Lys (%)	1.00	0.85	0.74	0.67	0.95
Met/Lys	58	64	68	72	66
Thr/Lys	65	65	65	65	65
Trp/Lys	16	15	15	15	15

¹ Alimet: Brand name of 2-hydroxy-4-methylthio butanoic acid.

² Provided per kg of final diet: vitamin A 10,000 IU; vitamin D 8,800 IU; vitamin E 220 IU; vitamin K 22 mg; vitamin B₁₂ 19.25 mg; niacin 250 mg; pantothenic acid 137 mg; riboflavin 41.25 mg; I 0.26 mg; Fe 140 mg; Mn 30 mg; Se 0.25 mg; and Zn 140 mg.

Table 5. Effects of Cu(HMTBa)₂ and CuSO₄ on performance in Exp. 1¹

	NRC control CuSO ₄ , (6 mg/kg)	CuSO ₄ 170 mg/kg	Cu(HMTBa) ₂ 170 mg/kg	SEM	p-value
Pig body weight (kg)					
d 0	7.36	7.36	7.35	0.10	0.99
d 14	10.44	10.54	11.27	0.24	0.06
d 42	22.07 ^b	22.07 ^b	23.40 ^a	0.37	0.03
d 0 to 14					
ADG (g)	220 ^b	227 ^b	279 ^a	12	<0.01
ADFI (g)	310 ^b	324 ^b	352 ^a	9	0.01
Gain:feed	0.709 ^a	0.701 ^a	0.793 ^b	0.026	0.05
d 15 to 42					
ADG (g)	408	405	393	12	0.17
ADFI (g)	644	601	608	16	0.17
Gain:feed	0.633	0.674	0.646	0.017	0.31
d 0 to 42					
ADG (g)	346 ^b	346 ^b	378 ^a	7	0.01
ADFI (g)	537	510	544	10	0.09
Gain:Feed	0.644	0.678	0.695	0.013	0.06

^{a,b} Means in a row with no common superscripts significantly differ (p<0.05).

¹ Values represent the means of 8 pens with 10 pigs per pen.

6.0% heavier than pigs fed the NRC control or CuSO₄ at the same level (p = 0.03, Table 5). During phase I (d 0 to 14), piglets fed Cu(HMTBa)₂ gained 22.9% more weight (279 vs 227 g/d, p<0.01), ate 8.6% more feed (352 vs 324 g/d, p = 0.01), and had better feed conversion (0.79 vs 0.70, p = 0.05) than animals fed 170 mg/kg CuSO₄. No difference was observed between pigs fed the NRC control (Cu, Mn, and Zn) and CuSO₄ at EU maximum level. In phase II (d 14 to 42), no differences were observed among treatments (p>0.17). Overall, piglets fed Cu(HMTBa)₂ gained 9.0% more (p = 0.01) and tended to eat more feed (p = 0.09) than animals fed 170 mg/kg CuSO₄. Feed conversion rate tended to be improved with Cu(HMTBa)₂ supplementation compared with the other two treatments (p = 0.06).

Liver Cu was increased 2.7 times with 170 mg/kg CuSO₄ supplementation compared with the NRC control, and was further increased with Cu(HMTBa)₂ (4.5 times, p<0.05, Table 6). No significant differences were observed on Cu deposition in tibia bone, kidney, muscle, and skin and fat (p>0.23).

Exp. 2

Pigs fed CuSO₄ and Cu(HMTBa)₂ tended to have better ADG (p = 0.08) and improved feed efficiency (p = 0.01) compared with the NRC control (Table 7). The Cu₂(OH)₃Cl was intermediate and did not differ from the NRC control. No difference was observed on feed intake among treatments (p = 0.81).

Liver Cu was significantly increased with Cu(HMTBa)₂ compared with the control and Cu₂(OH)₃Cl while CuSO₄ was intermediate (p = 0.01, Table 8). No difference was observed on liver Fe, Zn, and Mn concentration (p>0.27).

Exp. 3

Pigs fed 80 mg/kg Cu(HMTBa)₂ tended to be heavier than those fed either the NRC control or 160 mg/kg CuSO₄ at the end of experiment (p = 0.07, Table 9). Compared with the negative control, 160 mg/kg CuSO₄ improved weight gain and feed intake during the phase (d 21 to 42) immediately after 21 days of depletion (p<0.05). No benefit of 160 mg/kg CuSO₄ was observed during all other phases

Table 6. Effects of Cu(HMTBa)₂ and CuSO₄ on tissue Cu concentration (DM basis, Exp. 1)¹

Tissues Cu ² mg/kg	NRC control CuSO ₄ , 6 mg/kg	CuSO ₄ 170 mg/kg	Cu(HMTBa) ₂ 170 mg/kg	SEM	p-value
Tibia bone	0.64	0.64	0.76	0.07	0.40
Kidney	3.59	3.69	4.58	0.42	0.23
Liver	6.83 ^b	18.37 ^{ab}	30.46 ^a	0.41	<0.01
Muscle	0.56	0.51	0.53	0.03	0.53
Skin and fat	1.06	1.41	1.40	0.19	0.35

^{a,b} Means in a row with no common superscripts significantly differ (p<0.05).

¹ Values represent the means of 8 pigs per treatment.

² Cu content was analyzed using atomic absorption spectrometry following AOAC 968.08.

Table 7. Effects of different Cu supplementation on performance (Exp. 2)¹

	NRC control	CuSO ₄	Cu ₂ (OH) ₃ Cl	Cu(HMTBa) ₂	SEM	p-value
Body weight (kg)						
d 0	5.02	5.02	5.01	5.01	0.25	1.00
d 21	10.04	10.71	10.64	10.82	0.43	0.59
d 0 to 21						
ADG (g)	238	271	266	280	11	0.08
ADFI (g)	316	335	334	335	16	0.81
Gain:feed	0.753 ^b	0.814 ^a	0.798 ^{ab}	0.825 ^a	0.015	0.01

^{a,b} Means in a row with no common superscripts significantly differ (p<0.05).

¹ Values represent the means of 7 pens with 22 pigs per pen.

Table 8. Effects of different Cu sources on liver mineral concentrations (DM basis, Exp. 2)¹

Parameters ²	NRC control	CuSO ₄	Cu ₂ (OH) ₃ CL	Cu(HMTBa) ₂	SEM	p-value
Cu (mg/kg)	39.82 ^b	52.12 ^{ab}	48.80 ^b	73.32 ^a	8.52	0.01
Fe (mg/kg)	436.3	501.8	475.4	550.5	48.7	0.27
Mn (mg/kg)	10.12	8.38	9.00	9.43	0.78	0.46
Zn (mg/kg)	776.7	909.7	918.9	916.4	108.9	0.73

^{a,b} Means in a row with no common superscripts significantly differ (p<0.05).

¹ Values represent the means of 7 pigs per treatment.

² Mineral content was analyzed using atomic absorption spectrometry following AOAC 968.08.

and overall. Due to variation in different phases, only overall effects of Cu(HMTBa)₂ are discussed in this paper. Linear and quadratic dose response was observed on weight gain (p<0.01) and feed efficiency (p = 0.05) with Cu(HMTBa)₂ supplementation with the best performance was observed at 80 mg/kg. Pigs fed 80 mg/kg Cu(HMTBa)₂ tended to have improved weight gain (p = 0.07) and feed efficiency (p = 0.10) compared with those fed either the NRC control or 160 mg/kg Cu from CuSO₄.

Pigs fed 80 mg/kg Cu(HMTBa)₂ had the heaviest carcass weight, which was significant differ from those fed either the NRC control or CuSO₄ at 160 mg/kg (p<0.01, Table 10). Linear dose response of Cu(HMTBa)₂ was observed on loin depth with 55.73, 55.96, 56.64, 58.12, and 58.12 mm for the NRC control, and Cu(HMTBa)₂ at 20, 40, 80, or 160 mg/kg, respectively (p = 0.04). Loin depth from pigs fed 80 mg/kg and 160 mg/kg Cu(HMTBa)₂ were significantly greater than pigs fed 160 mg/kg CuSO₄ and the NRC control. No treatment difference was observed on back fat, yield, and lean percentage (p>0.37). In summary, no benefit of 160 mg/kg CuSO₄ was observed on carcass traits and BW. Growth performance, final carcass weight, and loin depth increased with increased Cu(HMTBa)₂ supplementation with the best performance was observed with 80 mg/kg Cu(HMTBa)₂.

DISCUSSION

High level of CuSO₄ is typically used in the swine industry as growth promoter and its benefits on weight gain and feed efficiency have been well documented (Bunch et

al., 1961;1963). Braude (1967) recorded 83 studies involving a total of 1,215 pigs in each treatment (0 vs 250 mg/kg) and reported that CuSO₄ improved daily weight gain on average 8.1%, and feed efficiency on average 5.4%. The optimum dose of CuSO₄ as growth promoter is around 242 mg/kg and 125 mg/kg of Cu was approximately 75% as effective (Cromwell et al., 1989). In agreement with previous studies, CuSO₄ improved weight gain and feed efficiency in the current nursery studies. In addition, Cu(HMTBa)₂ is more effective compared with CuSO₄ as growth promoter in stimulating gain and feed efficiency in both nursery and finisher pigs. These results agree with Coffey et al. (1994) and Veum et al. (2004) that chelated Cu is more effective than CuSO₄ and less is required to achieve similar performance. With limited Cu supplementation allowance in swine diets, using chelated Cu could be an alternative to pharmaceutical Cu to maintain growth promoting effects and reduce Cu excretion to the environment. The growth advantage of chelated Cu is perhaps due to the fact it is more available for absorption and less antagonism in the lumen as indicated by increased liver Cu storage and better antibacterial function.

It has been proposed that CuSO₄ acts as an anti-bacterial agent by shifting the microbial population within the gastrointestinal tract (Apgar et al., 1995; Arias and Koutsos, 2006), by quantitatively modifying some gram-positive bacterial populations in the gut. Evidence that Cu produces a growth promoting effect possibly through the microbial gut flora is supported by Shurson et al. (1990), who observed a positive effect of high concentration (283 mg/kg) of Cu in the diet on daily growth rate and feed

Table 9. Effects of CuSO₄ and Cu(HMTBa)₂ on performance in grower-finisher pigs (Exp. 3)^{1,2}

Treatments mg/kg	CuSO ₄		Cu(HMTBa) ₂				SEM	p-value		
	4	160	20	40	80	160		Treatment	Linear	Quadratic
Body weight (kg)										
d 0	32.29	32.42	32.52	32.33	32.22	32.40	0.29	0.98	-	-
d 21	52.57	52.58	52.52	52.60	52.54	52.56	0.30	1.00	0.99	0.99
d 42	72.16	73.43	72.00	71.75	72.41	72.88	0.42	0.08	0.89	0.99
d 62	89.45	90.54	89.35	89.52	89.38	90.27	0.43	0.24	0.88	0.75
d 80	101.69	101.88	100.69	100.92	101.98	102.48	0.57	0.24	0.96	0.75
d 100	117.54	117.86	117.28	118.22	119.67	118.91	0.60	0.07	0.14	0.25
d 0 to 21										
ADG (kg)	0.967	0.960	0.952	0.960	0.968	0.956	0.007	0.62	0.36	0.37
ADFI (kg)	1.899	1.865	1.870	1.865	1.881	1.874	0.015	0.61	0.83	0.85
Gain:feed	0.511	0.516	0.509	0.516	0.513	0.510	0.004	0.67	0.55	0.51
d 21 to 42										
ADG (kg)	0.932 ^{bcd}	0.988 ^a	0.927 ^{cd}	0.907 ^d	0.947 ^{bc}	0.961 ^{ab}	0.011	<0.01	0.65	0.95
ADFI (kg)	2.405 ^{bc}	2.486 ^a	2.411 ^{bc}	2.363 ^c	2.418 ^{abc}	2.445 ^{ab}	0.026	0.04	0.95	0.82
Gain:feed	0.389 ^{abc}	0.398 ^a	0.384 ^c	0.385 ^{bc}	0.394 ^{ab}	0.394 ^{ab}	0.003	0.04	0.19	0.32
d 42 to 62										
ADG (kg)	0.865	0.849	0.868	0.889	0.847	0.873	0.011	0.11	0.26	0.27
ADFI (kg)	2.561	2.555	2.536	2.553	2.525	2.555	0.026	0.92	0.83	0.80
Gain:feed	0.339	0.335	0.343	0.350	0.335	0.343	0.004	0.16	0.36	0.40
d 62 to 80										
ADG (kg)	0.680 ^{ab}	0.630 ^{bc}	0.619 ^c	0.634 ^{bc}	0.699 ^a	0.665 ^{abc}	0.017	0.02	<0.01	<0.01
ADFI (kg)	2.434	2.391	2.386	2.277	2.459	2.458	0.008	0.12	0.46	0.62
Gain:feed	0.280	0.263	0.258	0.289	0.285	0.271	0.01	0.31	0.19	0.19
d 80 to 100										
ADG (kg)	0.835	0.791	0.816	0.837	0.876	0.821	0.02	0.10	0.06	0.06
ADFI (kg)	2.340	2.266	2.332	2.399	2.343	2.292	0.05	0.58	0.74	0.60
Gain:feed	0.358	0.350	0.351	0.350	0.374	0.359	0.007	0.16	0.09	0.11
d 0 to 100										
ADG (kg)	0.858	0.852	0.840	0.856	0.872	0.857	0.007	0.07	<0.01	<0.01
ADFI (kg)	2.315	2.308	2.303	2.289	2.319	2.317	0.015	0.75	0.81	0.89
Gain:feed	0.370	0.371	0.365	0.373	0.378	0.369	0.003	0.10	0.05	0.05

^{a,b,c,d} Means in a row with no common superscripts significantly differ (p<0.05).

¹ Values represent the means of 8 pens with 20 to 22 pigs per pen.

² All pigs were fed a Cu deficient diet for the first 21 days (d 0 to 21).

conversion rate in conventional pigs and a negative effect in germ-free pigs. On the other hand, similar growth-promoting effect was obtained when copper histidinate or histidine solution was administered by intravenous injection and thus bypassed the gastrointestinal tract (Zhou et al.,

1994).

The authors suspect that injected extra Cu is secreted to the bile and back to the gastrointestinal tract, and has similar impact on bacteria population in the lumen as through dietary supplementation. If this is the case, dietary Cu needs

Table 10. Benefits of Cu(HMTBa)₂ on carcass traits in grower-finisher pigs (Exp 3.)¹

Treatments mg/kg	CuSO ₄		Cu(HMTBa) ₂				SEM	p-value		
	4	160	20	40	80	160		Treatment	Linear	Quadratic
Carcass wt. (kg)	88.13 ^b	88.24 ^b	88.13 ^b	88.13 ^b	90.57 ^a	89.43 ^{ab}	0.51	<0.01	0.07	0.17
Back fat (mm)	17.91	18.29	17.81	18.11	17.88	18.36	0.30	0.72	0.94	0.80
Yield (%)	74.62	74.92	75.16	73.78	74.77	74.73	0.44	0.37	0.70	0.66
Lean (%)	54.63	54.33	54.72	54.64	54.93	54.56	0.19	0.38	0.58	0.55
Loin depth (mm)	55.73 ^c	55.19 ^c	55.96 ^{bc}	56.64 ^{abc}	58.12 ^a	58.12 ^{ab}	0.58	<0.01	0.04	0.11

^{a,b,c} Means in a row with no common superscripts significantly differ (p<0.05).

¹ There were 8 replicates per treatment with 20-22 pigs per pen. All pigs were marketed at the same day.

to be absorbed first in order to be effective, which is the case in this study that Cu(HMTBa)₂ was more effective than CuSO₄ as growth promoter in both nursery and finisher pigs. Arnold (2011) reported that antimicrobial effects of bile determined by measuring the diameter of the zone of inhibition tended to be higher in Cu(HMTBa)₂ treatments compared with CuSO₄ at 75 mg/kg. However bile components, intestinal bacterial populations, and small intestine gene expression profiles associated with Cu absorption and homeostasis (Ctr-1, Atox-1, Cox-17, ATP7a, and ATP7b) were not affected. More research is needed to understand the effects of bile Cu recycle pathway on growth performance and antibacterial function.

Even though CuSO₄ is widely used in the swine industry, there are negative effects of adding high CuSO₄. CuSO₄ is highly water soluble and hygroscopic, which drawing moisture to the surface of each crystal. In vitamin premix and diet, CuSO₄ acts as a hot spot for reaction that break down labile organic compounds such as vitamins, enzymes, fat and oil (Dove and Ewan, 1991; Marchetti et al., 2000). *In vivo*, CuSO₄ dissociates and generates free Cu in the stomach, which is very active pro-oxidative substrate and leads to lipid oxidation and cell damage (Ajuwon et al., 2011). Absorption of Cu by the stomach is much less than what takes place in the intestine and is further hindered by binding with phytic acid and forming insoluble compounds. Banks et al. (2004) reported that 250 mg/kg CuSO₄ reduced apparent P retention 0.029 percentage-units in broilers, by forming an insoluble Cu-phytate chelate complex. The chelate is insoluble at the pH of the small intestine, and therefore none of the components of the complex are available for absorption. In addition, Cu competes with other trace minerals for absorption and utilization, such as Zn, Fe, and Mn. Hall et al. (1979) reported that a 20% decrease in ⁶⁵Zn absorption occurred when dietary Cu was raised from 3 to 24 mg/kg in rats. One advantage of chelated mineral is that chelation between ligand(s) and mineral preventing mineral dissociation and complex with other nutrients and consequently avoiding antagonism with other nutrients in the lumen. Previous studies suggested that chelated minerals are more bioavailable and better at avoiding antagonism *in vivo* based on tissue mineral, gene expression, and functional assays compared with inorganic trace minerals (Richards et al., 2010; Zhao et al., 2010). Because of its higher availability, less Cu(HMTBa)₂ can be used to replace high CuSO₄ and have similar or better growth promoter effects in pigs. Our follow-up study indicated that pigs fed 50 mg/kg Cu(HMTBa)₂ performed equally as pigs fed 250 mg/kg CuSO₄ (Zhao et al., 2009). Similar results were reported in broilers that reduced level of chelated trace minerals can be used to replace inorganic trace mineral and have similar or even better performance (Zhao et al., 2010). Additional benefit of using chelated

trace mineral is reduced mineral excretion to the environment (Managi et al., 2010).

Using high CuSO₄ in grower-finisher pigs is less common and research results are controversial. Cromwell et al. (1978) reported 3% improvement of gain and feed efficiency with 250 mg/kg CuSO₄ supplementation. No significant benefit of 250 mg/kg CuSO₄ was observed by Gipp et al. (1973) and Dove and Ewan (1991). This is in agreement with the current study that the only benefit of 160 mg/kg CuSO₄ was observed during the first 21 days immediately after a three-week depletion period. Interestingly, quadratic dose response was observed on growth performance, carcass weight, and loin depth with pigs fed 80 mg/kg Cu(HMTBa)₂ had the best gain and the greatest loin depth. Mechanism of Cu(HMTBa)₂ improved loin depth is not well understood. Zhou et al. (1994) suggested that Cu stimulated the growth of longissimus muscle more than the growth of the whole body, potentially through stimulating growth hormone secretion. The same authors also reported increased Cu concentration in the brain, which indicated that Cu may cross the blood-brain barrier and regulate hormone secretion. Similar results were reported by Myres and Bowland (1973), who observed that pigs fed high Cu diets had larger loin cross-sections than those receiving no supplemental Cu. Improved loin eye area suggests higher lean percentage and better carcass value. Further research is needed to better understand the effect of high Cu on improving loin muscle growth.

IMPLICATION

Cu(HMTBa)₂ is more available based on performance and liver Cu concentration, so less Cu(HMTBa)₂ is needed to have similar growth promoter effects as CuSO₄ when fed to nursery pigs. For grower-finisher pigs, 80 mg/kg Cu(HMTBa)₂ can be used as growth promoter and pigs fed Cu(HMTBa)₂ had greater loin depth compared with 160 mg/kg CuSO₄. By using less chelated Cu in pig diets, less Cu may be excreted to the environment although Cu concentration in feces was not analyzed in these trials.

REFERENCES

- Ajuwon, O. R., O. M. O. Idowu, S. A. Afolabi, B. O. Kehinde, O. O. Oguntola, and K. O. Olatunbosun. 2011. The effects of dietary copper supplementation on oxidative and antioxidant systems in broiler chickens. *Arch. Zootec.* 60:275-282.
- AOAC. 2000. Official methods of analysis (17th Ed). Association of Official Analytical Chemists. Arlington, VA.
- Apgar, G. A., E. T. Kornegay, M. D. Lindemann, and D. R. Notter. 1995. Evaluation of copper sulfate and a copper lysine complex as growth promoters for weanling swine. *J. Anim. Sci.* 73:2640-2646.
- Arias, V. J. and E. A. Koutsos. 2006. Effects of copper source and

- level on intestinal physiology and growth of broiler chickens. *Poult. Sci.* 85:999-1007.
- Arnold, M. A. 2011. Interaction of copper source, bile composition on microbial growth, and Cu protein homeostasis. Master thesis. Colorado State Univ. Fort Collins, Colorado.
- Banks, K. M., K. L. Thompson, J. K. Rush, and T. J. Applegate. 2004. Effects of copper source on phosphorus retention in broiler chicks and laying hens. *Poult. Sci.* 83:990-996.
- Braude, R. 1967. Copper as a stimulant in pig feeding. *World Rev. Anim. Prod.* 3:69-82.
- Bunch, R. J., V. C. Speer, V. W. Hays, and J. T. McCall. 1963. Effects of high levels of copper and chlortetracycline on performance of pigs. *J. Anim. Sci.* 22:56-60.
- Bunch, R. J., V. C. Speer, V. W. Hays, J. H. Hawbaker, and D. V. Catron. 1961. Effects of copper sulfate, copper oxide and chlortetracycline on baby pig performance. *J. Anim. Sci.* 20:723-726.
- Coffey, R. D., G. L. Cromwell, and H. J. Monegue. 1994. Efficacy of a copper-lysine complex as a growth promotant for weanling pigs. *J. Anim. Sci.* 72:2880-2886.
- Cromwell, G. L., T. S. Stahly, and H. J. Monegue. 1989. Effects of source and level of copper on performance and liver copper stores in weanling pigs. *J. Anim. Sci.* 67:2996-3002.
- Cromwell, G. L., V. W. Hays, and T. L. Clark. 1978. Effect of copper sulfate, copper sulfide and sodium sulfide on performance and copper stores of pigs. *J. Anim. Sci.* 46:692-698.
- Dove, C. R. and R. C. Ewan. 1991. Effect of vitamin E and copper on the vitamin E status and performance of growing pigs. *J. Anim. Sci.* 69:2516-2523.
- FEDNA, 2003. Guidelines of the Spanish Foundation for Development of Animal Nutrition for the formulation of compound feeds (Ed. C. de Blas, P. Garcia, and G. G. Mateos). Madrid Polytechnical University, Spain.
- Gipp, W. F., W. G. Pond, and E. F. Walker, Jr. 1973. Influence of diet composition and mode of copper administration on the response of growing-finishing swine to supplemental copper. *J. Anim. Sci.* 36:91-99.
- Hall, A. C., B. W. Young, and I. Bremner. 1979. Intestinal metallothionein and the mutual antagonism between copper and zinc in the rat. *J. Inorg. Biochem.* 11:57-66.
- Hawbaker, J. A., V. C. Speer, V. W. Hays, and D. V. Catron. 1961. Effect of copper sulfate and other chemotherapeutics in growing swine rations. *J. Anim. Sci.* 20:163-167.
- Kornegay, E. T. and A. F. Harper. 1997. Environmental nutrition: Nutrient management strategies to reduce nutrient excretion of swine. *Prof. Anim. Sci.* 13:99-111.
- Marchetti, M., H. D. Ashmead, N. Tossani, S. Marchetti, and S. D. Ashmead. 2000. Comparison of the rates of vitamin degradation when mixed with metal sulphates or metal amino acid chelates. *J. Food Composit. Anal.* 13:875-884.
- Manangi, M. K., T. Hampton, P. Fisher, J. D. Richards, M. Vazquez-Anon, and K. D. Christensen. 2010. Impact of feeding lower levels of chelated minerals vs. industry levels of inorganic trace minerals on broiler performance, yield, foot pad health, and litter minerals concentration. International Poultry Scientific Forum. Southern Poultry Science Society (SPSS) Annual Meeting and the Southern Conference on Avian Diseases Annual Meeting. Atlanta, GA, January 24-25, 2010.
- Myres, A. W. and J. P. Bowland. 1973. Effects of environmental temperature and dietary copper on growth and lipid metabolism in pigs. Concentration and composition of plasma free fatty acids. *Can. J. Anim. Sci.* 53:127-132.
- NRC. 1998. Nutrient requirements of swine 10th ed. National Academy Press. Washington, DC.
- Richards, J. D., J. Zhao, R. J. Harrell, C. A. Atwell, and J. J. Dibner. 2010. Trace mineral nutrition in poultry and swine. *Asian Australas. J. Anim. Sci.* 23:1527-1534.
- Roof, M. D. and D. C. Mahan. 1982. Effect of carbadox and various dietary copper levels for weanling swine. *J. Anim. Sci.* 55:1109-1117.
- Shurson, G. C., P. K. Ku, G. L. Waxler, M. T. Yokoyama, and E. R. Miller. 1990. Physiological relationships between microbiological status and dietary copper levels in the pig. *J. Anim. Sci.* 68:1061-1071.
- Tucker, M. R. 1997. Experiences with metal toxicities in North Carolina. Pages 97-100 in Proc. 40th Annual Meet. Soil Sci. Soc. North Carolina, Raleigh.
- Veum, T. L., M. S. Carlson, C. W. Wu, D. W. Bollinger, and M. R. Ellersieck. 2004. Copper proteinate in weanling pig diets for enhancing growth performance and reducing fecal copper excretion compared with copper sulfate. *J. Anim. Sci.* 82:1062-1070.
- Wang, Z., S. Cerrate, C. Coto, F. Yan, and P. W. Waldroup. 2007. Evaluation of Mintrex copper as a source of copper in broiler diets. *Int. J. Poultry Sci.* 6:308-313.
- Yi, G. F., C. A. Atwell, J. A. Hume, J. J. Dibner, C. D. Knight, and J. D. Richards. 2007. Determining the methionine activity of Mintrex organic trace minerals in broiler chicks by using radiolabel tracing or growth assay. *Poult. Sci.* 86:877-887.
- Zhao, J., R. B. Shirley, M. Vazquez-Anon, J. J. Dibner, J. D. Richards, P. Fisher, T. Hampton, K. D. Christensen, J. P. Allard, and A. F. Giesen. 2010. Effects of chelated trace minerals on growth performance, breast meat yield and foot pad health in commercial meat broilers. *J. Appl. Poultry Res.* 19:365-372.
- Zhao, J., R. J. Harrell, G. Allee, B. Hinson, P. Winkelbauer, C. Atwell, J. D. Richards, and M. Vazquez-Anon. 2009. Effect of an organic copper source on growth performance and tissue copper concentration in nursery pigs. Mid-west Animal Science Meeting. Des Moines, IA.
- Zhao, J., R. B. Shirley, T. R. Hampton, J. D. Richards, R. J. Harrell, J. J. Dibner, C. D. Knight, M. Vazquez-Anon, and A. F. Giesen. 2008. Benefits of an organic trace mineral on performance with dietary Cu antagonism in broilers. *Poult. Sci.* 87:53(Abstr.).
- Zhou, W., E. T. Kornegay, M. D. Lindemann, J. W. Swinkels, M. K. Welten, and E. A. Wong. 1994. Stimulation of growth by intravenous injection of copper in weanling pigs. *J. Anim. Sci.* 72: 2395-2403.