

LETTER to the EDITOR

Helicobacter Pylori vacA d1 Genotype is associated with Gastric Cancer but not Peptic Ulcers in Kurdistan Region, Northern Iraq*Asian Pac J Cancer Prev*, **15 (14)**, 5965-5966**Dear Editor**

We read with interest the paper by Basiri and colleagues studying the prevalence of *vacA d1* and *d2* genotypes in the *H. pylori* isolates from patients with gastric adenocarcinoma, peptic ulcer disease and gastritis in East Azerbaijan region, where the incidence of gastric cancer (GC) is high. They found that 43.4% of the isolate carried the *d1* allele and 56.6% carried subtype *d2*. A significant correlation between *vacA d1* and gastric adenocarcinoma or PUD was shown. The study suggested that the *H. pylori vacA d1* genotype helps predict risk for gastric adenocarcinoma and PUD in East Azerbaijan, Iran (Basiri et al., 2014).

Despite the geographical proximity of Iraq and Iran the incidence of gastric cancer differs hugely between these countries; in Iran it ranges from 38-69/10⁵ compared to 5/10⁵ in Iraq (GLOBOCAN, 2002; Sadjadi et al., 2005; Yavari et al., 2006). Previous studies of *H. pylori* virulence factors were not successful in the explanation of such a difference in cancer rate. We hypothesised that *H. pylori vacA d1* may help understanding this variation in cancer rate in these two neighbouring countries. DNA was extracted from 157 *H. pylori* strains isolated from patients with different clinical outcome (non-ulcer disease (NUD): 81; gastric ulcer (GU): 31; duodenal ulcer (DU):38 and gastric cancer:5). Using the same methodology mentioned in the Basiri et al paper, we typed these strains for *vacA d* subtypes. Overall, (45/157) 28.6% of our samples typed as *vacA d1* and (112/157) 71.4% as *vacA d2*. It was found that (21/81) 26% of the strains isolated from NUD typed as *vacA d1*. In addition, *vacAd1* genotype was found in (9/38) 23.6% and (10/31) 30.3% of DU and GU samples, respectively. Interestingly, all the isolates obtained from GC patients were *vacAd1*. No association was found between *vacA d* genotypes and DU and GU (Chi square test; $p > 0.05$). However, a statistical significant relationship was found between *vacA d1* and gastric cancer (Exact Fischer test; $p = 0.002$).

H. pylori strains possessing *cagA* are associated with a significantly increased risk for the development of atrophic gastritis, peptic ulcer disease (PUD) and gastric cancer (Atherton, 2006). Previous study comparing *H. pylori* strains from these two countries showed that no difference in prevalence of *cagA+* strains between unselected dyspeptic populations from these countries (Hussein et

al., 2008). The vacuolating cytotoxin (VacA) is a well-established *H. pylori* virulence factor which has multiple effects including vacuolation of cultured epithelial cells, inducing apoptosis, increasing permeability of epithelial monolayers, forming pores in cells and suppressing immune cell function (Atherton, 2006). The *vacA* gene is polymorphic within its signal, intermediate and mid regions. Also, no significant differences between Iranian and Iraqi populations in other *vacA* genotypes was found, and in particular in the *i* region type which has recently been linked with gastric cancer risk in Iran (Rhead et al., 2007). Recently, a fourth disease-related region between the *i* and *m* regions has been indentified and named the deletion (*d*) region. The *d* region is divided into *d1* and *d2* with *d1* identified a risk factor for GC and PUD in Western strains. Nearly all *H. pylori* isolates from East Asia have been found to have a genotype of *vacA d1* (Ogiwara et al., 2009). Our results showed that the overall prevalence of *vacA d1* genotype is lower than that found in Iran this could contribute to the differences in gastric cancer rates seen between these communities.

In agreement with findings from Basiri et al's paper and results from Western countries, this study showed a significant relationship between *vacA d1* and gastric cancer. This study therefore supports the proposal of Basiri et al's conclusion that the *H. pylori vacA d1* genotype might be a new risk marker for gastric cancer and peptic ulcer.

References

- Atherton J (2006). The pathogenesis of *H. pylori*-induced gastro-duodenal diseases. *Ann Rev Pathology: Mechanisms of Disease*, **1**, 63-96.
- Basiri Z, Safaralizadeh R, Bonyadi MJ, et al (2014). *Helicobacter pylori vacA d1* genotype predicts risk of gastric adenocarcinoma and peptic ulcers in Northwestern Iran. *Asian Pac J Cancer Prev*, **15**, 1575-79.
- GLOBOCAN IARC (2002). Cancer Map: Male Stomach Cancer, Age-Standardized Incidence Rate per 100,000.
- Hussein NR, Mohammadi M, Talebkhan Y, et al (2008). Differences in virulence markers between *Helicobacter pylori* strains from Iraq and those from Iran: potential importance of regional differences in *H. pylori*-associated disease. *J Clin Microbiology*, **46**, 1774-79.
- Ogiwara H, Sugimoto M, Ohno T, et al (2009). Role of deletion located between the intermediate and middle regions of the *Helicobacter pylori vacA* gene in cases of gastroduodenal diseases. *J Clin Microbiology*, **47**, 3493-500.

Nawfal Rasheed Hussein

Rhead JL, Letley DP, Mohammadi M, et al (2007). A new *Helicobacter pylori* vacuolating cytotoxin determinant, the intermediate region, is associated with gastric cancer. *Gastroenterology*, **133**, 926-36.

Sadjadi A, Nouraie M, Mohagheghi MA, et al (2005). Cancer occurrence in Iran in 2002, an international perspective. *Asian Pac J Cancer Prev*, **6**, 359-63.

Yavari P, Hislop TG, Bajdik C, et al (2006). Comparison of cancer incidence in Iran and Iranian immigrants to British Columbia, Canada. *Asian Pac J Cancer Prev*, **7**, 86-90.

Nawfal Rasheed Hussein

Department of Internal Medicine, School of Medicine, Faculty of Medical Sciences, University of Duhok, Kurdistan Region, Iraq
**For correspondence: nawfal.hussein@yahoo.com*