

## TOEPLITZ TYPE OPERATOR IN $\mathbb{C}^n$

KI SEONG CHOI\*

ABSTRACT. For a complex measure  $\mu$  on  $B$  and  $f \in L^2_a(B)$ , the Toeplitz operator  $T_\mu$  on  $L^2_a(B, d\nu)$  with symbol  $\mu$  is formally defined by  $T_\mu(f)(w) = \int_B f(w) \overline{K(z, w)} d\mu(w)$ . We will investigate properties of the Toeplitz operator  $T_\mu$  with symbol  $\mu$ . We define the Toeplitz type operator  $T_\psi^r$  with symbol  $\psi$ ,

$$T_\psi^r f(z) = c_r \int_B \frac{(1 - \|w\|^2)^r}{(1 - \langle z, w \rangle)^{n+r+1}} \psi(w) f(w) d\nu(w).$$

We will also investigate properties of the Toeplitz type operator with symbol  $\psi$ .

### 1. Introduction

Integration with respect to area measure is denoted with  $dA$ . The set of all analytic functions on  $D$  will be denoted by  $H(D)$  or simply  $H$ . The Bergman space  $L^2_a$  of the unit disk is the space of all functions analytic in  $D$  which belongs to  $L^2 = L^2(dA)$ , that is,  $L^2_a = L^2 \cap H$ . The inner product in  $L^2$  is denoted by  $\langle f, g \rangle = \frac{1}{\pi} \int_D f(z) \overline{g(z)} dA(z)$ .

Let  $P$  be the orthogonal projection from  $L^2$  to  $L^2_a$ . It satisfies  $Pf(w) = \langle f, K_w \rangle$  for all  $f \in L^2$  where the function  $K_z(w) = K(z, w) = (1 - \bar{w}z)^{-2}$  is the Bergman kernel. In particular, if  $f \in L^2_a$ , then  $f(w) = \langle f, K_w \rangle$ .

McDonald and Sundberg defined Toeplitz operators on the Bergman space  $L^2_a$  by  $T_\psi f = P(\psi f)$ , where  $\psi$  is a function on the interior of the disk and  $P$  is the Bergman projection from  $L^2$  to  $L^2_a$  (See [10]).

In the Bergman space, one can have  $\psi f \in L^2$  for all  $f \in L^2_a$  even if  $\psi$  is unbounded. Moreover, the formula for the Bergman projection as an integral can be applied even when the product  $\psi f$  is only in  $L^1$ . The formula for the Toeplitz operator,

---

Received September 22, 2014; Accepted October 20, 2014.

2010 Mathematics Subject Classification: Primary 32H25; Secondary 30H05.

Key words and phrases: Bergman space, Toeplitz operator, Toeplitz type operator.

$$P(\psi f)(z) = \int_D \frac{\psi(w)f(w)}{(1-wz)^2} dA(z),$$

allows one to extend the notion of Toeplitz operators to symbols that are measures: simply replace  $\psi dA$  with  $d\mu$  in the formula.

Let  $\mu$  be any complex regular Borel measure on the unit disk  $D$  in the complex plane  $\mathbb{C}$ . The Toeplitz operator on  $L^2_a$  with symbol  $\mu$  is denoted by  $T_\mu$  and is formally defined by

$$T_\mu(f)(w) = \frac{1}{\pi} \int_D \frac{f(z)}{(1-\bar{z}w)^2} d\mu(z).$$

If  $\mu$  has the form  $\psi dA$  for some bounded measurable function  $\psi$ , then  $T_\mu$  is defined by  $T_\psi$  and satisfies  $T_\psi f = P(\psi f), f \in L^2_a$ . For arbitrary measures on  $D$ ,  $T_\mu$  may be only densely defined because the above integral can only be guaranteed to converge for bounded  $f$ .

In this paper, we will consider the Toeplitz operators in the complex space  $\mathbb{C}^n$ . Throughout this paper,  $\mathbb{C}^n$  will be the Cartesian product of  $n$  copies of  $\mathbb{C}$ . For  $z = (z_1, z_2, \dots, z_n)$  and  $w = (w_1, w_2, \dots, w_n)$  in  $\mathbb{C}^n$ , the inner product is defined by  $\langle z, w \rangle = \sum_{j=1}^n z_j \bar{w}_j$  and the norm by  $\|z\|^2 = \langle z, z \rangle$ .

Let  $B$  be the open unit ball in the complex space  $\mathbb{C}^n$ . Let  $\nu$  be the Lebesgue measure in  $\mathbb{C}^n$  normalized by  $\nu(B) = 1$ . We let  $L^2(B, d\nu)$  be the usual space of Lebesgue square-integrable complex valued functions on  $B$ . The Bergman space  $L^2_a(B, d\nu)$  is defined to be the subspace of  $L^2(B, d\nu)$  consisting of analytic functions.

Fix a point  $z \in B$ . Since the functional  $e_z$  given by  $e_z(f) = f(z), f \in L^2_a(B, d\nu)$ , is continuous, there exists a function  $k_z \in L^2_a(B, d\nu)$  such that

$$f(z) = \langle f, k_z \rangle = \int_B f(w) \overline{k_z(w)} d\nu(w)$$

by the Riesz representation theorem. The function  $k_z(w) = K(z, w)$  is called the Bergman reproducing kernel in  $L^2_a(B, d\nu)$ .

Let  $P$  be the Bergman projection defined by

$$Pf(z) = \int_B f(w) K(w, z) d\nu(w).$$

The Bergman projection is used in many areas related with Hankel operators and Toeplitz operators(See [1, 3, 7, 8, 12, 13]).

For a complex measure  $\mu$  on  $B$  and  $f \in L^2_a(B, d\nu)$ , the Toeplitz operator on  $L^2_a(B, d\nu)$  with symbol  $\mu$  is denoted by  $T_\mu$  and is formally

defined by

$$T_\mu(f)(z) = \int_B f(w) \overline{K(z, w)} d\mu(w).$$

We will view  $T_\mu$  as an operator defined on the dense subset of polynomials with range in the set of all analytic functions.

In Section 2, we will investigate properties of the Toeplitz operator  $T_\mu$  with symbol  $\mu$ . In particular, we will show that if  $\mu$  is a positive Borel measure on  $B$  such that  $\|\mu\|_r = \sup_{z \in B} \frac{\mu(E(z, r))}{\nu(E(z, r))} < \infty$ , then  $T_\mu$  is bounded.

Let  $L_a^p(B, d\nu)$  be the Bergman space of analytic functions in  $L^p(B, d\nu)$ .  $L_a^p(B, d\nu)$  is a Banach space for all  $1 \leq p < \infty$ . Given a function  $f$  on  $B$ , the Toeplitz operator  $T_f$  is defined by  $T_f g = P(fg)$  where  $P$  is the Bergman projection. We let  $H^\infty(B)$  denote the space of bounded analytic functions in  $B$ .

Zhu used the Toeplitz operator to complete characterization for the multipliers of the Bloch and the little Bloch space of the open unit ball in  $\mathbb{C}^n$  (See [13]).

The measure  $\mu_r$  is the weighted Lebesgue measure:

$$d\mu_r(z) = c_r(1 - \|z\|^2)^r d\nu(z)$$

where  $r > -1$  is fixed, and  $c_r$  is a normalization constant such that  $\mu_r(B) = 1$ .

If we equip  $L_{a,r}^2 = L_a^2(B, d\mu_r)$  with the norm  $\|f\|_{2,r} = \sqrt{\int_B |f|^2 d\mu_r}$ , then  $L_{a,r}^2$  is a Banach space for each  $r > -1$ . It was shown in [6] that if  $f \in L_{a,r}^1, r > -1$ , then

$$\begin{aligned} f(z) &= \int_B f(w) \overline{k_{r,z}(w)} d\mu_r(w) \\ &= c_r \int_B \frac{(1 - \|w\|^2)^r}{(1 - \langle z, w \rangle)^{n+r+1}} f(w) d\nu(w) \end{aligned}$$

where  $\overline{k_{r,z}(w)} = \frac{1}{(1 - \langle z, w \rangle)^{n+r+1}}$ . Suppose  $1 \leq p < +\infty$  and  $r > 0$ . Let  $L_{a,r}^p$  be the subspace of  $L^p(B, d\mu_r)$  consisting of analytic functions.

For some bounded measurable function  $\psi$ , we will define the Toeplitz type operator  $T_\psi^r$  with symbol  $\psi$ ,

$$\begin{aligned} T_\psi^r f(z) &= \int_B \psi(w) f(w) \overline{k_{r,z}(w)} d\mu_r(w) \\ &= c_r \int_B \frac{(1 - \|w\|^2)^r}{(1 - \langle z, w \rangle)^{n+r+1}} \psi(w) f(w) d\nu(w). \end{aligned}$$

In section 3, we will investigate properties of the Toeplitz type operator with symbol  $\psi$ . In particular, we will show that the Toeplitz type operator  $T_\psi^r$  with symbol  $\psi$  is bounded on  $L^p(B, d\nu)$  for  $1 \leq p < +\infty$  and  $r > 0$ .

## 2. Toeplitz operator with symbol $\mu$

The function  $k_z(w) = K(z, w)$  is the Bergman representing kernel in  $L_a^2(B, d\nu)$ . The function  $K(\cdot, \cdot)$  is actually defined and continuous on  $B \times \bar{B}$  (where  $\bar{B}$  is the closure of  $B$  in  $\mathbb{C}^n$ ). Since  $k_z \in L_a^2$ , we have

$$\begin{aligned} k_z(w) &= \overline{K(z, w)} \\ &= \int_B \overline{K(z, \varsigma)} K(w, \varsigma) dV(\varsigma) \\ &= \overline{\int_B \overline{K(w, \varsigma)} K(z, \varsigma) dV(\varsigma)} = K(w, z). \end{aligned}$$

**THEOREM 2.1.** *If  $T_\mu$  is the Toeplitz operator on  $L_a^2(B, d\nu)$  with symbol  $\mu$ , then*

$$T_\mu = 0 \text{ if and only if } \mu = 0.$$

*Proof.* Suppose  $f$  and  $g$  are in  $C(B)$  which is the set of continuous functions on  $B$ . Then

$$\begin{aligned} &\langle T_\mu f, g \rangle \\ &= \int_B T_\mu f(z) \overline{g(z)} d\nu(z) \\ &= \int_B \int_B f(w) \overline{K(w, z)} d\mu(w) \overline{g(z)} d\nu(z) \\ &= \int_B f(w) \overline{\int_B g(z) \overline{K(z, w)} d\nu(z)} d\mu(w) \\ &= \int_B f(w) \overline{g(w)} d\mu(w). \end{aligned}$$

Since the closure of  $\text{span}\{w^k, \bar{w}^n\}_{k, n \geq 0} = C(B)$ ,  $T_\mu = 0$  if and only if  $\mu = 0$ . □

We define the Berezin transform of  $\mu$  by

$$\tilde{\mu}(z) = \int_B |k_z(w)|^2 d\mu(w)$$

and consider the usual supremum  $\|\tilde{\mu}\|_\infty \equiv \sup_{z \in B} |\tilde{\mu}(z)|$ .

The Bergman metric  $\beta(\cdot, \cdot)$  gives the usual topology on  $B$  (See [7, p52]). Moreover, the closed metric balls

$$E(z, r) = \{w : \beta(z, w) \leq r\}$$

are compact (See [7, p56]). For any fixed  $r > 0$ , we define

$$\|\mu\|_r = \sup_{z \in B} \frac{\mu(E(z, r))}{\nu(E(z, r))}.$$

Let

$$\|\mu\|_{s,p} = \sup\{\|h\|_\mu^p / \|h\|^p : h \in L_a^p(B, d\nu), h \neq 0\}$$

where  $\|h\|^p = \int_B |h(w)|^p d\nu(w)$  and  $\|h\|_\mu^p = \int_B |h(w)|^p d\mu(w)$ .

**THEOREM 2.2.** *Suppose  $\mu$  is a finite positive Borel measure on  $\Omega$ ,  $p > 1$  and  $r > 0$ . Then quantities  $\|\mu\|_r$ ,  $\|\tilde{\mu}\|_\infty$  and  $\|\mu\|_{s,p}$  are all equivalent (There exist some constants  $a, b$  and  $c$  such that  $\|\mu\|_r \leq a \|\tilde{\mu}\|_\infty \leq b \|\mu\|_{s,p} \leq c \|\mu\|_r$ ).*

*Proof.* See [4, Theorem 6]. □

**THEOREM 2.3.**  *$T_\mu$  is the Toeplitz operator on  $L_a^2(B, d\nu)$  with symbol  $\mu$ . If  $\mu$  is a positive Borel measure on  $B$  such that  $\|\mu\|_r = \sup_{z \in B} \frac{\mu(E(z, r))}{\nu(E(z, r))} < \infty$ , then  $T_\mu$  is bounded.*

*Proof.* For any two polynomials  $f$  and  $g$ ,

$$\begin{aligned} & |\langle T_\mu f, g \rangle| \\ &= \left| \int_B f(w) \overline{g(w)} d\mu(w) \right| \\ &\leq \int_B |f(w) \overline{g(w)}| d\mu(w) \\ &\leq c \int_B |f(w) \overline{g(w)}| d\nu(w) \\ &\leq c \|f\|^2 \|g\|^2 \end{aligned}$$

where third inequality follows from Theorem 2.2. Thus  $T_\mu$  is bounded. □

### 3. Toeplitz type operator with symbol $\psi$

THEOREM 3.1. *If  $f \in L^1_{a,r}$ ,  $r > -1$ , then*

$$f(z) = c_r \int_B \frac{(1 - \|w\|^2)^r}{(1 - \langle z, w \rangle)^{n+r+1}} f(w) d\nu(w).$$

*Proof.* See [6, Theorem 2]. □

The notation  $a(z) \sim b(z)$  means that the ratio  $a(z)/b(z)$  has a positive finite limit as  $\|z\| \rightarrow 1$ .

THEOREM 3.2. *For  $z \in B$ , a real  $c$  and  $t > -1$ , define*

$$I_{c,t}(z) = \int_B \frac{(1 - \|w\|^2)^t}{|1 - \langle z, w \rangle|^{n+1+c+t}} d\nu(w).$$

Then,

- (i)  $I_{c,t}(z)$  is bounded in  $B$  if  $c < 0$ ;
- (ii)  $I_{0,t}(z) \sim -\log(1 - \|z\|^2)$  as  $\|z\| \rightarrow 1^-$ ;
- (iii)  $I_{c,t}(z) \sim (1 - \|z\|^2)^{-c}$  as  $\|z\| \rightarrow 1^-$  if  $c > 0$ .

*Proof.* See [11, Proposition 1.4.10]. □

THEOREM 3.3. *Let  $X$  and  $Y$  be Banach spaces. Let  $\mathfrak{L}(X, Y)$  be the set of bounded linear transformations from  $X$  to  $Y$ . If  $A^*$  is the adjoint of  $A \in \mathfrak{L}(X, Y)$ , then:*

- (1)  $\|A^*\| = \|A\|$ ;
- (2) If  $A$  is invertible, then  $A^*$  is invertible and  $(A^*)^{-1} = (A^{-1})^*$ .

*Proof.* See [5, Proposition 1.4]. □

Recall that, for some bounded measurable function  $\psi$ , the Toeplitz type operator  $T_\psi^r$  with symbol  $\psi$  is defined by

$$T_\psi^r f(z) = c_r \int_B \frac{(1 - \|w\|^2)^r}{(1 - \langle z, w \rangle)^{n+r+1}} \psi(w) f(w) d\nu(w).$$

THEOREM 3.4. *For  $r > 0$ , the Toeplitz type operator  $T_\psi^r$  with symbol  $\psi$  is bounded on  $L^1(B, d\nu)$ .*

*Proof.*

$$\begin{aligned} & \langle T_\psi^r f, g \rangle \\ &= \int_B T_\psi^r f(z) \overline{g(z)} d\nu(z) \\ &= \int_B \int_B \frac{c_r(1 - \|w\|^2)^r \psi(w) f(w)}{(1 - \langle z, w \rangle)^{n+r+1}} d\nu(w) \overline{g(z)} d\nu(z) \\ &= \int_B f(w) c_r(1 - \|w\|^2)^r \psi(w) \int_B \frac{\overline{g(z)}}{(1 - \langle z, w \rangle)^{n+r+1}} d\nu(z) d\nu(w) \end{aligned}$$

where  $g \in L^\infty(B)$ . Let  $T_\psi^{r*}$  be the adjoint of  $T_\psi^r$  under the usual integral pairing. Then above result shows that

$$T_\psi^{r*} g(w) = c_r(1 - \|w\|^2)^r \psi(w) \int_B \frac{g(z)}{(1 - \langle w, z \rangle)^{n+r+1}} d\nu(z).$$

By Theorem 3.2, if  $r > 0$

$$\sup_{w \in B} (1 - \|w\|^2)^r \int_B \frac{d\nu(z)}{(1 - \langle w, z \rangle)^{n+r+1}} < \infty.$$

This shows that if  $r > 0$ , then  $T_\psi^{r*}$  is bounded on  $L^\infty(B, d\nu)$ . By Theorem 3.3,  $T_\psi^r$  is bounded on  $L^1(B, d\nu)$  if  $r > 0$ .  $\square$

**COROLLARY 3.5.** *If  $r < 0$ , then  $T_\psi^r$  is not bounded.*

*Proof.*

$$\begin{aligned} & \int_B T_\psi^{r*} 1(w) d\nu(w) \\ &= \int_B c_r(1 - \|w\|^2)^r \int_B \frac{\psi(z) d\nu(z)}{(1 - \langle w, z \rangle)^{n+r+1}} d\nu(w) \end{aligned}$$

By Theorem 3.2,  $\int_B \frac{d\nu(z)}{(1 - \langle w, z \rangle)^{n+r+1}} d\nu(w)$  is bounded. But  $\int_B (1 - \|w\|^2)^r d\nu(w)$  is not finite for  $r \leq -1$ . This shows that  $T_\psi^r$  is not bounded.  $\square$

**THEOREM 3.6.** *Suppose  $(X, \mu)$  is a measure space and  $\Psi$  is a measurable function on  $X \times X$ . Let  $T$  be the integral operator induced by  $\Psi$ , that is,*

$$Tf(x) = \int_X \Psi(x, y) f(y) d\mu(y).$$

Suppose  $1 < p < +\infty$  with  $\frac{1}{p} + \frac{1}{q} = 1$ . If there is a constant  $c > 0$  and a positive measurable function  $h$  on  $X$  such that

$$\int_X |\Psi(x, y)| h(y)^q d\mu(y) \leq ch(x)^q$$

for  $\mu$ -almost every  $x$  in  $X$  and

$$\int_X |\Psi(x, y)| h(x)^p d\mu(x) \leq ch(y)^p$$

for  $\mu$ -almost every  $y$  in  $X$ , then  $T$  is bounded on  $L^p(X, d\mu)$  with norm less than or equal to  $c$ .

*Proof.* See [14, Theorem 3.2.2]. □

**THEOREM 3.7.** Suppose  $1 < p < +\infty$  and  $r > 0$ . Then the Toeplitz type operator  $T_\psi^r$  with symbol  $\psi$  is bounded on  $L^p(B, d\nu)$ .

*Proof.* For  $f \in L^p(B, d\nu)$ ,

$$T_\psi^r f(z) = \int_B \frac{c_r(1 - \|w\|^2)^r f(w) \psi(w)}{|1 - \langle z, w \rangle|^{n+r+1}} d\nu(w).$$

For  $h(z) = (1 - \|z\|^2)^{-\frac{1}{pq}}$  and  $\Psi(z, w) = \frac{c_r(1 - \|w\|^2)^r f(w) \psi(w)}{|1 - \langle z, w \rangle|^{n+r+1}}$ ,

$$\begin{aligned} & \int_B |\Psi(z, w) \psi(w)| h(w)^q d\nu(w) \\ &= \int_B \left| \frac{c_r(1 - \|w\|^2)^r}{|1 - \langle z, w \rangle|^{n+r+1}} \psi(w) \right| (1 - \|w\|^2)^{-\frac{1}{p}} d\nu(w) \\ &\leq c(1 - \|z\|^2)^{-\frac{1}{p}} \\ &= ch(z)^q \end{aligned}$$

where the second inequality follows from Theorem 3.2.

$$\begin{aligned} & \int_B |\Psi(z, w) \psi(w)| h(z)^q d\nu(z) \\ &= c_r(1 - \|w\|^2)^r |\psi(w)| \int_B \frac{(1 - \|z\|^2)^{-\frac{1}{q}}}{|1 - \langle z, w \rangle|^{n+r+1}} d\nu(z) \\ &= c_r(1 - \|w\|^2)^r |\psi(w)| \int_B \frac{(1 - \|z\|^2)^{-\frac{1}{q}}}{|1 - \langle z, w \rangle|^{n+(r+1/q)+1-1/q}} d\nu(z) \\ &\leq c_r |\psi(w)| (1 - \|w\|^2)^{-\frac{1}{q}} \\ &\leq ch(w)^p \end{aligned}$$



where the fourth inequality follows from Theorem 3.2. By Theorem 3.6,  $T_\psi^r$  is bounded on  $L^p(B, d\nu)$  if  $r > 0$ .  $\square$

### References

- [1] J. Arazy, S. D. Fisher, and J. Peetre, *Hankel operators on weighted Bergman spaces*, Amer. J. Math. **110** (1988), 989-1054.
- [2] K. S. Choi, *Lipschitz type inequality in Weighted Bloch spaces  $\mathfrak{B}_q$* , J. Korean Math. Soc. **39** (2002), no. 2, 277-287.
- [3] K. S. Choi, *Little Hankel operators on Weighted Bloch spaces*, Commun. Korean Math. Soc. **18** (2003), no. 3, 469-479.
- [4] K. S. Choi, *Notes on Carleson Measures on bounded symmetric domain*, Commun. Korean Math. Soc. **22** (2007), no. 1, 65-74.
- [5] J. B. Conway, *A course in Functional Analysis*, Springer Verlag, New York, 1985.
- [6] K. T. Hahn and K. S. Choi, *Weighted Bloch spaces in  $\mathbb{C}^n$* , J. Korean Math. Soc. **35** (1998), 177-189.
- [7] L. Helgason, *Differential Geometry, Lie Groups and Symmetric Spaces*, Academic Press, New York/London, 1978.
- [8] D. H. Luecking, *A Technique for characterizing Carleson measures on Bergman spaces*, Proc. Amer. Math. Soc. **87** (1983), 656-660.
- [9] D. H. Luecking, *Finite rank Toeplitz operators on the Bergman Space*, Proc. Amer. Math. Soc. **136** (2008), 1717-1723.
- [10] G. McDonald and C. Sundberg, *Toeplitz operators on the disc*, Indiana Univ. Math. J. **28** (1979), 595-611.
- [11] W. Rudin, *Function theory in the unit ball of  $\mathbb{C}^n$* , Springer Verlag, New York, 1980.
- [12] K. H. Zhu, *Duality and Hankel operators on the Bergman spaces of bounded symmetric domains*, J. Funct. Anal. **81** (1988), 260-278.
- [13] K. H. Zhu, *Multipliers of BMO in the Bergman metric with applications to Toeplitz operators*, J. Funct. Anal. **87** (1989), 31-50.
- [14] K. H. Zhu, *Operator theory in function spaces*, Marcel Dekker, New York, 1990.

\*

Department of Information Security  
Konyang University  
Nonsan 320-711, Republic of Korea  
*E-mail*: ksc@konyang.ac.kr