Spiro-β-lactams Synthesized from Imines Derived from 8-Azabicyclo[3.2.1]octan-3-ones

Ju-Hyun Song, Yeon-geun Kwon, Hyun-Ok Lee,[†] Dai-II Jung,^{*} and Jung-Tai Hahn[‡]

Department of Chemistry, Dong-A University, Busan 604-714, Korea. *E-mail: dijung@dau.ac.kr *Korea Kumho Pertrochemical Co., Ltd. Korea *Department of Beautycare, Young-Dong University, Chungcheongbuk-do 370-701, Korea Received July 8, 2014, Accepted August 14, 2014

Key Words : Spiro- β -lactam, 8-Azabicyclo[3.2.1]octan-3-one, Anticonvulsant, Tropane alkaloid

Tropane alkaloids have attracted considerable attention because of their remarkable pharmaceutical significance.¹⁻⁴ In particular, a series of tropanes exhibited anticonvulsant activity against pentylenetetrazol-induced convulsions in mice and antiarrhythmic activity in rabbits previously treated with ouabain.5-8 Therefore, a variety of synthetic approaches to prepare tropane alkaloids have been investigated. In addition, the importance of β -lactams for the treatment of bacterial infections has been well established.9-10 Considerable efforts have been made for the synthesis and structural modification of the β -lactam nucleus to increase its antimicrobial activity. Apart from their therapeutic utility, β -lactams are versatile synthons that are used for the preparation of α - and β -amino acids.¹⁰ Although many functionalized monocyclic β -lactams have been reported thus far, there have been relatively few efforts toward the synthesis of spiro- β -lactams.¹¹ Among the different strategies employed for the construction of β lactams,¹² the reaction of acid chlorides with imines (Staudinger synthesis)¹³ is one of the most popular procedures. As a part of our study on the improvement of anticonvulsants and the development of new drugs, here, we report the synthesis of a new class of spiro- β -lactams having an 8-azabicyclo[3.2.1]octan-3-one ring. We have already published a few reports on the synthetic study of pharmacologically interesting 8-azabicyclo[3.2.1]octan-3-one compounds derived from the reaction of corresponding amines 1 with 2,5-dimethoxytetrahydrofuran and 1,3-acetonedicarboxylic acid in the presence of HCl and H₂O.¹⁴⁻¹⁸ Imines 3 derived from 8azabicyclo[3.2.1]octan-3-one (Scheme 1) may serve as an interesting starting material for the synthesis of a new class of spiro- β -lactams such as 4. Furthermore, in these structures, the inherent reactivity of the 8-azabicyclo[3.2.1]octan-3-one framework can be combined with the synthetic versatility of the β -lactam moiety in a single compound. To the best of our knowledge, the Staudinger synthesis of compounds 3 has not been reported previously; therefore, our paper might be the first to report results in this field. Phenyl-(8-phenyl-8azabicyclo[3.2.1]oct-3-ylidene)-amine 3a was synthesized by the reaction of 8-phenyl-8-azabicyclo[3.2.1]octan-3-one 2a (1.0 g, 5 mmol) in benzene with aniline (0.5 g, 5 mmol) and montmorillonite (5 g) by using the Dean-Stark apparatus. The reaction mixture was refluxed for a further 24 h and then cooled to room temperature. After filtering the montmorillonite, the organic layer was concentrated. The residue was chromatographed on a silica gel (n-hexane:dichloromethane = 1:10, v/v) to yield **3a** (0.84 g, 61%) as a brown sticky oil. The structures and yields of the synthesized imines (3a-3e) are summarized in Scheme 1 and Table 1.



Scheme 1. Synthesis of spiro- β -lactams 4 from amines 1. Reagent and reaction conditions: (I) 2,5-dimethoxytetrahydrofuran, 1,3-acetonedicarboxylic acid, HCl, H₂O, rt; (II) 4-substituted-aniline, montmorillonite, benzene, reflux; (III) phenoxyacetyl chloride, triethylamine, toluene.

Table 1. Physical data from 8-substituted-8-azabicyclo[3.2.1]octan-3-ones **2** to spiro- β -lactams **4**

Entry	Reactant	Product	Yield $(\%)^a$	mp (°C)
1	1a	2a	51	99–100
2	1b	2b	85	51-52
3	1c	2c	85	88-89
4	1d	2d	85	155-156
5	1e	2e	78	Liq.
6	2a	3 a	61	Liq.
7	2b	3b	31	Liq
8	2c	3c	82	157-158
9	2c'	3c'	34	148–149
10	2d	3d	55	Liq.
11	2e	3e	15	Liq.
12	3a	4 a	91	127-128
13	3b	4b	69	96–97
14	3c	4c	30	164–165
15	3c'	4c'	82	120-121
16	3e	4e	73	136–137

^aYields are isolated yields.

The Staudinger synthesis of compounds 3 with the phenoxyacetyl chloride afforded spiro- β -lactams 4 (Scheme 1). 3-Phenoxy-1,8'-diphenyl-8'-azaspiro[azetidine-2,3'-bicyclo-[3.2.1]octan]-4-one 4a was synthesized by the reaction of phenyl-(8-phenyl-8-azabicyclo[3.2.1]oct-3-ylidene)-amine **3a** (25 mg, 9×10^{-2} mmol) in toluene (50 mL) with triethylamine and phenoxyacetyl chloride (16 mg, 9.5×10^{-2} mmol). The reaction mixture was stirred at room temperature for a further 5 h. After stirring for 5 h, the organic layer was filtered and concentrated. The residue was chromatographed on a silica gel (*n*-hexane:EtOAc = 4:1, v/v) to yield 4a (34 mg, 91%) as a light-yellow solid. The structures and yields of the synthesized spiro- β -lactams (4a-4c') are summarized in Scheme 1 and Table 1. These were obtained as single diastereomers, and no traces of the corresponding isomeric $exo-\beta$ -lactam 5 were observed.



It is noteworthy that this stereochemical outcome of β lactam formation with phenoxyacetyl chloride under Staudinger synthesis conditions was opposite to that expected from a simple [2+2]-cycloaddition reaction,^{19,20} which should have occurred from the exo face of compound **2** to afford compounds **4**.²¹ A plausible mechanism for the formation of spiro- β -lactams **4** is shown in Scheme 2.

The initial acylation of imines **3** followed by the addition of a chloride ion²¹ from the exo-face of the iminium ion should afford intermediate **A**. Enolization followed by intramolecular endo-displacement of the chloride should afford spiro- β -lactams **4**. This is the reaction pathway observed for the highly reactive oxygen-substituted enolate as a phenoxyacetyl chloride.²² In conclusion, the Staudinger synthesis of imines derived from 8-azabicyclo[3.2.1]octan-3-one with phenoxyacetyl chloride affords the corresponding spiro- β lactams, but with an unexpected exo stereochemistry. Studies aimed at gaining a greater understanding of this Staudinger synthesis of imines with new structures as well as those aimed at the synthetic application of these findings are underway and these will be reported in due course.

Experimental Section

Melting points were determined using an electrothermal capillary melting point apparatus and uncorrected. Thin layer chromatography (TLC) was performed on glass plates coated with silicon oxide (silica gel 60F₂₅₄), and compounds were visualized using a UV lamp. ¹H and ¹³C NMR spectra were



Scheme 2. Plausible mechanism for the formation of β -lactams 4 from imines 3.

Notes

obtained with Bruker AC 2000 (200 MHz) and Varian Gemini (200 or 300 MHz) spectrometers. Mass spectra were measured with HP 5890 GC/Mass (70 eV, EI). The organic solvents and chemicals were obtained from commercial products and purified by the appropriate methods before use. Except where explicitly stated, all the starting materials were purchased from Aldrich, Fluka, Fisher, Lancaster, or TCI chemical companies and used as received.

3-Phenoxy-1,8'-diphenyl-8'-azaspiro[azetidine-2,3'-bi-cyclo[3.2.1]octan]-4-one (4a). Yield: 91%; mp 127–128 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.26 (m, 7H), 7.04 (s, 1H), 6.95 (m, 2H), 6.81 (t, 1H), 6.77 (s, 2H), 6.74 (d, *J* = 7.9 Hz, 2H), 4.60 (s, 1H), 4.37 (m, 2H), 4.29 (s, 2H), 2.63 (dd, *J* = 3.2, 17.1 Hz, 1H), 2.34 (m, 1H), 2.16 (s, 1H), 2.14 (m, 1H), 1.78 (s, 1H), 1.61 (s, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 167.10, 157.99, 146.50, 133.72, 129.52, 129.38, 128.84, 126.44, 125.16, 121.42, 121.41, 116.42, 114.70, 66.74, 53.47, 34.15, 33.02, 29.80; GC/MS *m/z* 410 (M+); Anal. Calcd. for C₂₇H₂₆N₂O₂: C, 79.00; H, 6.38; N, 6.82; O, 7.79. Found: C, 78.70; H, 6.46; N, 6.77; O, 7.83.

3-Phenoxy-1-phenyl-8'-*p***-ethylphenyl-8'-azaspiro[azetidine-2,3'-bicyclo[3.2.1]octan]-4-one (4b).** Yield: 69%; mp 96–97 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.25 (m, 7H), 7.01 (s, 1H), 6.89 (m, 2H), 6.75 (s, 2H), 6.71 (d, , *J* = 7.9 Hz, 2H), 4.59 (s, 1H), 4.34 (m, 2H), 4.28 (s, 2H), 2.61 (dd, , *J* = 3.0, 17.1 Hz, 1H), 2.59 (q, *J* = 7.6 Hz, 2H), 2.32 (m, 1H), 2.15 (s, 1H), 2.11 (m, 1H), 1.77 (s, 1H), 1.59 (s, 1H), 1.21 (t, *J* = 7.6 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 167.13, 157.98, 146.45, 133.52, 129.87, 129.35, 129.28, 128.84, 126.40, 125.15, 121.42, 121.40, 114.72, 66.69, 53.61, 34.27, 33.15, 29.89, 27.80, 15.54; GC/MS *m/z* 438 (M+); Anal. Calcd. for C₂₉H₃₀N₂O₂: C, 79.42; H, 6.89; N, 6.39; O, 7.30. Found: C, 79.40; H, 6.93; N, 6.38; O, 7.31.

3-Phenoxy-1-phenyl-8'-*p***-fluorophenyl-8'-***a***zaspiro**[**azeti-dine-2,3'-bicyclo**[**3.2.1**]**octan**]**-4-one** (**4c**). Yield: 30%; mp 164–165 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.23 (m, 5H), 6.93 (m, 5H), 6.76 (m, 4H), 4.56 (s, 1H), 4.33 (s, 2H), 4.26 (d, *J* = 8.2 Hz, 2H), 2.55 (dd, *J* = 3.3, 17.2 Hz, 1H), 2.31 (m, 1H), 2.10 (m, 2H), 1.76 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 167.10, 158.32, 155.17, 143.74, 129.88, 129.51, 127.02, 125.65, 121.95, 117.87, 116.28, 115.14, 67.27, 54.33, 34.68, 33.25, 30.27; GC/MS *m*/*z* 428 (M+); Anal. Calcd. for C₂₇H₂₅FN₂O₂C, 75.68; H, 5.88; F, 4.43; N, 6.54; O, 7.47. Found: C, 75.57; H, 5.92; F, 4.49; N, 6.53; O, 7.49.

3-Phenoxy-1-*p*-fluorophenyl-8'-*p*-fluorophenyl-8'-azaspiro[azetidine-2,3'-bicyclo[3.2.1]octan]-4-one (4c'). Yield: 82%; mp 120–121 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.02 (m, 4H), 6.80 (m, 5H), 6.69 (m, 4H), 4.58 (s, 1H), 4.42 (s, 2H), 4.28 (d, *J* = 8.2 Hz, 2H), 2.65 (dd, *J* = 3.3, 17.2 Hz, 1H), 2.35 (m, 1H), 2.13 (m, 2H), 1.79 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 171.16, 158.22, 155.20, 153.52, 143.51, 130.40, 129.55, 123.13, 120.90, 117.97, 116.51, 115.93, 115.47, 67.47, 54.38, 34.95, 33.46, 30.33; GC/MS *m/z* 446 (M+); Anal. Calcd. for C₂₇H₂₄F₂N₂O₂: C, 72.63; H, 5.42; F, 8.51; N, 6.27; O, 7.17. Found: C, 72.58; H, 5.45; F, 8.53; N, 6.20; O, 7.19.

3-Phenoxy-1-phenyl-8'-furfuryl-8'-azaspiro[azetidine-

2,3'-bicyclo[3.2.1]octan]-4-one (4e). Yield: 73%; mp 136–137 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.25 (m, 7H), 7.06 (d, J = 2.17 Hz, 1H), 6.99 (s, 1H), 6.87 (m, 2H), 6.80 (t, 1H), 6.77 (s, 2H), 6.45 (d, J = 7.8 Hz, 2H), 6.32 (m, 1H), 6.23 (d, J = 3.56 Hz, 1H), 4.59 (s, 1H), 4.37 (m, 2H), 4.25 (s, 2H), 3.68 (s, 2H), 2.68 (dd, J = 3.1, 17.1 Hz, 1H), 2.30 (m, 1H), 2.12 (s, 1H), 2.07 (m, 1H), 1.71 (s, 1H), 1.48 (s, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 167.05, 157.59, 145.34, 134.33, 129.79, 129.12, 128.86, 127.96, 126.10, 125.03, 121.42, 113.88, 66.21, 52.42, 33.97, 33.15, 28.37; GC/MS *m/z* 414 (M+); Anal. Calcd. for C₂₆H₂₆N₂O₃: C, 75.34; H, 6.32; N, 6.76; O, 11.58. Found: C, 75.32; H, 6.33; N, 6.79; O, 7.81.

Acknowledgments. This work was supported by the Dong-A University Research Fund (2013).

References

- Fodor, G. Chemistry of the Alkaloids; Pelletier, S. W., Ed.; Van Nostrand-Reinhold: Princeton, New Jersey, 1970; Chapter 15.
- Fodor, G.; Janzso, G.; Dobo, P.; Koczor, J. Abh. Disch. Akad. Wiss. Berlin, Kl. Chem. Geol. Biol. 1964, 187.
- De La Camp, U. O.; Bottini, A. T.; Thut, C. C.; Gal, J.; Bellettini, A. G. J. Org. Chem. 1972, 37, 324.
- 4. Wiger, G. R.; Rettig, M. F. J. Am. Chem. Soc. 1976, 98, 4168.
- 5. Trigo, G. G.; Martines, M. Pharm. Mediaterr. 1974, 10, 643.
- Sacristan, A. G; Illera, M.; Sancher, F. S. Arch. Formcol. Toxicol. 1977, 3, 57.
- 7. Trigo, G. G.; Martinez, E. J. Pharm. Sci. 1981, 70, 87.
- Izquierdo, M. L.; Galvez, E.; Burgos, C.; Florencio, F. J. Heterocyclic Chem. 1988, 25, 419.
- (a) *The Organic Chemistry of β-Lactams*; Georg, G. I., Ed.; VCH: New York, 1993. (b) Nau, R.; Eiffert, A. *Clin. Microb. Rev.* 2002, *15*, 95.
- (a) Palomo, C. In *Recent Progress in the Chemical Synthesis of Antibiotics*; Luckacs, G., Ohno, M., Eds.; Springer: Berlin, 1990.
 (b) Palomo, C.; Aizpurua, J. M.; Ganboa, I.; Oiarbide, M. *Pure Appl. Chem.* 2000, 72, 1763. (c) Palomo, C.; Aizpurua, J. M.; Ganboa, I.; Oiarbide, M. *Synlett* 2001, 1813.
- For selected, recent references, see: (a) Strauss, A.; Otto, H. H. *Helv. Chim. Acta* **1997**, *80*, 1823. (b) Anklam, S.; Liebscher, J. *Tetrahedron* **1998**, *54*, 6369. (c) Dalla-Croce, P.; La Rosa, C. *Tetrahedron: Asymmetry* **1999**, *10*, 1193. (d) Barba, V.; Hernández, C.; Rojas-Lima, S.; Farfan, N.; Santillán, R. Can. J. Chem. **1999**, *77*, 2025. (e) Papillon, J. P. B.; Taylor, R. J. K. Org. Lett. **2000**, *2*, 1987. (f) Arjona, O.; Csákÿ, A. G.; Murcia, C.; Plumet, J. Tetrahedron Lett. **2002**, *43*, 6405.
- For selected, recent references, see: (a) Yamamoto, Y.; Asao, N.; Tsukuda, N. In *Advances in Asymmetric Synthesis*; Hassner, A. H., Ed.; Asymmetric Synthesis of β-Aminoacids and β-Lactam Derivatives *via* Conjugate Addition of Metal Amides. JAI Press: Stamford, CT, 1998; Vol. 3. (b) Muller, T. E.; Beller, M. *Chem. Rev.* **1998**, *98*, 675. (c) Benaglia, M.; Cinquini, M.; Cozzi, F. Eur. J. Org. Chem. **2000**, 563. (d) For solid phase and combinatorial synthesis, see: Mata, E. G. *Curr. Pharm. Des.* **1999**, *5*, 955.
- For a selected recent review, see: Palomo, C.; Aizpurua, J. M.; Ganboa, I.; Oiarbide, M. *Eur. J. Org. Chem.* 1999, 3223 and references cited therein.
- Jung, D. I.; Park, J. H.; Roh, S. A.; Lee, Y. G.; Park, Y. M.; Kim, I. S.; Jeong, I. S.; Park, M. S. J. Korean Chem. Soc. **1997**, *41*, 414.
- Jung, D. I.; Park, J. H.; Park, C. S.; Kim, Y. Y.; Jung, D. H.; Kim, I. S.; Park, M. S. J. Korean Chem. Soc. 1999, 43, 122.
- Jung, D. I.; Park, C. S.; Kim, Y. H.; Lee, D. H.; Lee, Y. G.; Park, Y. M.; Choi, S. K. Synth. Commun. 2001, 31, 3255.
- 17. Jung, D. I.; Song, J. H.; Kim, Y. H.; Lee, D. H.; Song, H. A.; Lee,

Notes

Y. G; Park, Y. M.; Choi, S. K.; Hahn, J. T.; Cho, S. J. Bull. Korean Chem. Soc. 2004, 25, 1932.

- Jung, D. I.; Song, J. H.; Lee, D. H.; Kim, Y. H.; Lee, Y. G.; Hahn, J. T. Bull. Korean Chem. Soc. 2006, 27, 1493.
- Different behavior of alkyl and alkozyacetyl chlorides in the Staudinger reaction has been previously reported. See: Georg, G. L.; He, P.; Kant, J.; Wu, Z. J. Org. Chem. 1993, 58, 5571 and references cited therein.
- 20. The stereochemistry at C-3' of the β -lactam moiety was deduced from NOE measurements. Thus, saturation of H-3' ($\delta = 4.50$ ppm, s) in **6b** gave rise to a 3% NOE in H-3_{endo} ($\delta = 1.60$ ppm, d, ${}^{2}J = 12.5$ Hz). No NOE was observed with H-3_{exo}. The spatial

proximity of H-3' and H-3_{endo} in **6b** (2.4 Å) was deduced from AM-1 calculations. On the other hand, a distande of 3.4 Å was found between H-3' and H-3_{exo}.

- exo-Addition is the characteristic reaction way of the carbonyl group of the derivatives of 7-oxanorbornen-5-one. See: (a) Vogel, P.; Cossy, J.; Plumet, J.; Arjona, O. *Tetrahedron* 1999, *55*, 13521. (b) Vogel, P. *Curr. Org. Chem.* 2000, *4*, 455.
- 22. For the participation of this kind of reaction pathway in β-lactam formation under Staudinger reaction conditions see: Arrieta, A.; Lecea, B.; Cossio, F. P. J. Org. Chem. 1998, 63, 5869 and references cited therein.