



Erratum

Choi, In-Su, So-Young Kim and Byoung-Hee Choi. 2015. A taxonomic revision of *Astragalus* L. (Fabaceae) in Korea. Korean J. Pl. Taxon. 45: 227–238.

Page	Column	Position	For	Read
232	Right	Line 41	Fruit	Fruiting
233		Fig. 2	Photographs of <i>Astragalus uliginosus</i> L. (A–C) and <i>A. schelichowii</i> Turcz. (D–F).	Photographs of <i>Astragalus schelichowii</i> Turcz. (A–C) and <i>A. uliginosus</i> L. (D–F).
234		Table 1	Comparism	Comparison
235	Left	Line 5	7. <i>Astragalus mongholicus</i> Bunge. Mém. Acad. Imp. Sci. Saint Pétersbourg 11: 25, 1868.	7. <i>Astragalus mongholicus</i> Bunge. Mém. Acad. Imp. Sci. Saint Pétersbourg 11 (16): 25, 1868 [and 15 (1): 30, 1869].
235	Left	Line 8	Perennial herb, pubescent with basifixed hairs; stems 50–100 cm long; erect, ascending, or decumbent, sparsely branched, covered with appressed long white hairs. Leaves 8–12 cm long, short petiolate; stipules free, cauline, green, foliaceous, triangular-ovate, subacute, 5–10 mm long, sparsely vested on the outside with white hairs; leaflets sessile, narrowly ovate or narrowly oblong, obtuse to retuse or rarely mucronulate at apex, 8–10 pairs, 0.4–2.0 cm long, 5–8 mm wide. Inflorescences racemes, 4–9 cm long, densely 10 to 15-flowered; flowers unilaterally pendent on inflorescence; peduncles 6–9 cm long; bracts narrowly lanceolate to subulate, as long as or shorter than calyx, 36 mm long, ca. 1 mm wide, sparsely covered with white hairs on the outside; pedicels 2, 3 mm long. Calyx 5–9 mm long, campanulate, strongly oblique, covered with short appressed black hairs; teeth subequal, triangular to narrowly triangular, 0.5–2.0 mm long. Corolla yellowish; standard 15–20 mm long, slightly longer or equal to other petals; the lamina obovate, retuse gradually attenuate, 34 times as long as the claw, reflexed in upper part; wings 17–19 mm long, the claw 1.5 times as long as the oblong obtuse lamina; keel equaling the wings, the claw 1.5 times as long as the acute lamina; ovary with a stipe 5–6 mm, hairy. Legumes pendulous, stipitate ca. 1 cm long, membranous, oblong, apex acute, inflated, 3–4 cm long, ca. 1.5 cm wide, sparsely covered with appressed minute hairs, unilocular, 5–7 seeded. stems erect; leaflets 1.5–2 cm long, both surfaces pubescent.	(delete)

Page	Column	Position	For	Read
236	Left	Line 1	Var. dahuricus (DC.) Podlech, Sendtnera 6: 172, 1999. Type. Russia, Dahuria, without date, <i>Fischer s.n.</i> (syntype, G; isosyntype, P seen as a photo!).	Var. dahuricus (DC.) Podlech, Sendtnera 6: 172, 1999. <i>Phaca alpina</i> L. var. <i>dahurica</i> DC., Prodr. 2: 273, 1825. <i>Astragalus membranaceus</i> Bunge, Mém. Acad. Imp. Sci. Saint Pétersbourg 11 (16): 25, 1868 [and 15 (1): 30, 1869], nom. illeg., non Moench, Methodus P. 167, 1794. Type. Russia, Dahuria, without date, <i>Fischer s.n.</i> (syntype, G; isosyntype, P seen as a photo!).
236	Left	Line 5	Stems erect; leaflets 1.5–2 cm long, both surfaces pubescent.	Perennial herb, pubescent with basifixed hairs; stems 50–100 cm long; erect, sparsely branched, covered with appressed long white hairs. Leaves 8–12 cm long, short petiolate; stipules free, cauline, green, foliaceous, triangular-ovate, subacute, 5–10 mm long, sparsely vested on the outside with white hairs; leaflets subsessile, narrowly ovate or narrowly oblong, obtuse to retuse or rarely mucronulate at apex, 810 pairs, 1.5–2.0 cm long, 5–8 mm wide, both surfaces pubescent. Inflorescences racemes, 49 cm long, densely 10 to 15-flowered; flowers unilaterally pendent on inflorescence; peduncles 6–9 cm long; bracts narrowly lanceolate to subulate, as long as or shorter than calyx, 3–6 mm long, ca. 1 mm wide, sparsely covered with white hairs on the outside; pedicels 2–3 mm long. Calyx 5–9 mm long, campanulate, strongly oblique, covered with short appressed black hairs; teeth subequal, triangular to narrowly triangular, 0.5–2.0 mm long. Corolla yellowish; standard 15–20 mm long, slightly longer or equal to other petals; the lamina obovate, retuse gradually attenuate, 3–4 times as long as the claw, reflexed in upper part; wings 17–19 mm long, the claw 1.5 times as long as the oblong obtuse lamina; keel equaling the wings, the claw 1.5 times as long as the acute lamina; ovary with a stipe 5–6 mm, hairy. Legumes pendulous, stipitate ca. 1 cm long, membranous, oblong, apex acute, inflated, 3–4 cm long, ca. 1.5 cm wide, sparsely covered with appressed minute hairs, unilocular, 5–7 seeded. Flowering July–August. Fruiting August–September.
236	Left	Line 16	Y.N. Lee (1996b)	Y.N. Lee (1996)
236	Left	Line 34	Lee Y., 1996b	Lee Y., 1996
236	Right	Line 37	Var. nakaianus (Y.N. Lee) I.S. Choi & B.H. Choi, Choi et al., Kor. J. Pl. Taxon. 43: 17, 2013. Lee, Kor. J. Bot. 24 (1): 28 (1981); Y.N. Lee, Fl. Kor.: 409 (1996) <i>A. membranaceus</i> var. <i>nakaianus</i> Y.N. Lee, New Fl. of Korea: 738 (2006), nom. illeg. Type. Korea, Jeju Isl., Aug. 9, 1957, <i>Y.N. Lee s.n.</i> , (lectotype, EWH!).	Var. nakaianus (Y.N. Lee) I.S. Choi & B.H. Choi, Choi et al., Kor. J. Pl. Taxon. 43: 17, 2013. <i>A. nakaianus</i> Y.N. Lee, Kor. J. Bot. 24 (1): 28, 1981. <i>A. membranaceus</i> var. <i>nakaianus</i> Y.N. Lee, New Fl. of Korea: 738, 2006., nom. illeg. Type. Korea, Jeju Isl., Aug. 9, 1957, <i>Y.N. Lee s.n.</i> , (lectotype, EWH!).