

The Importance of Esophageal and Gastric Diseases as Causes of Chest Pain

Yong Joo Kim, Eun Jung Shin, Nam Su Kim, Young Ho Lee, and Eun Woo Nam

Department of Pediatrics, Hanyang University College of Medicine, Seoul, Korea

Purpose: Pediatric chest pain is considered to be idiopathic or caused by benign diseases. This study was to find out how much upper gastrointestinal (UGI) diseases are major causes of chest pain in pediatric patients.

Methods: The records of 75 children (42 boys and 33 girls, aged 3-17 years old) who have presented with mainly chest pain from January 1995 to March 2015 were retrospectively reviewed. Chest X-ray and electrocardiography (ECG) were performed in all patients. Further cardiologic and gastrointestinal (GI) evaluations were performed in indicated patients.

Results: Chest pain was most common in the children of 6 and 9 to 14 years old. Esophago-gastric diseases were unexpectedly the most common direct causes of the chest pain, the next are idiopathic, cardiac diseases, chest trauma, respiratory disease, and psychosomatic disease. Even though 21 showed abnormal ECG findings and 7 showed abnormalities on echocardiography, cardiac diseases were determined to be the direct causes only in 9. UGI endoscopy was performed in 57 cases, and esophago-gastric diseases which thereafter were thought to be causative diseases were 48 cases. The mean age of the children with esophago-gastric diseases were different with marginal significance from that of the other children with chest pain not related with esophago-gastric diseases. All the 48 children diagnosed with treated with GI medicines based on the diagnosis, and 37 cases (77.1%) subsequently showed clinical improvement.

Conclusion: Diagnostic approaches to find out esophageal and gastric diseases in children with chest pain are important as well as cardiac and respiratory investigations.

Key Words: Chest pain, Child, Esophageal diseases, Stomach diseases, Endoscopy

INTRODUCTION

Chest pain in the pediatric population is not rare and is mostly a benign symptom. The causes of chest

pain are diverse, and formulating a differential diagnosis is not easy. In most cases pediatric chest pain is either idiopathic or is caused by benign diseases. This is not the case for adults, in whom chest pain is usu-

Received : August 27, 2015, Revised : September 18, 2015, Accepted : September 30, 2015

Corresponding author: Yong Joo Kim, Department of Pediatrics, Hanyang University Seoul Hospital, 222 Wangsimni-ro, Seongdong-gu, Seoul 04763, Korea. Tel: +82-2-2290-8390, Fax: +82-2-2290-8380, E-mail: kyjoo@hanyang.ac.kr

Copyright © 2015 by The Korean Society of Pediatric Gastroenterology, Hepatology and Nutrition

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ally the result of more aggressive diseases [1-3]. But, few clinical studies have addressed the causes and clinical characteristics of pediatric chest pain. The most common causes are musculoskeletal diseases, pulmonary diseases, and idiopathic cases. Less common cause are gastrointestinal diseases and cardiac diseases [4]. The situation in Korea is similar [5].

Sabri et al. [6] suggested that there is a higher proportion of upper gastrointestinal (UGI) disease among children and teenagers with chest pain, and recommended including a clinical examination focusing on GI disease to determine the true source of chest pain.

The aim of this study is to evaluate the causes of chest pain in children and to determine the usefulness of several diagnostic examinations used in formulating the differential diagnosis of pediatric chest pain.

MATERIALS AND METHODS

We retrospectively surveyed 75 children (42 males, 56.0%; 33 females, 44.0%) mainly presenting with chest pain who visited the pediatric clinic of Hanyang University Hospital (Seoul, Korea) from January 1995 to March, 2015 Children 9-14 years of age (57.3%) were most common, followed by 6-year-olds (14.7%) (Fig. 1).




Fig. 1. Age and sex distribution of the 75 children with chest pain.

Data on clinical history, physical examination, chest X-ray, and electrocardiography (ECG) were reviewed for each patient using outpatient records. Thorough cardiovascular examinations (including 24-hour Holter monitor recording, echocardiography, and exercise stress test), and UGI examinations (endoscopy, gastroesophageal scan, 24-hour esophageal pH monitoring) were performed in select patients depending on specific history and clinical status.

We investigated what diseases and organ diseases were the primary causes of chest pain in these children, and how valuable the diagnostic procedures were in terms of differential diagnoses. A simple abnormality in a clinical examination without true correlation was not regarded as the primary cause of the symptom. We made the final diagnoses in the patients according to the symptomatic improvement in 6 months upon specific treatment and whether the symptoms recurred or not afterwards. This study was approved from the institutional review board of Hanyang University Hospital (IRB No. HYUH 2015-07-029-001).

Statistical analysis

General descriptive statistics of patients are presented as frequencies and percentages (%). The prevalence and the proportion were evaluated by the exact or normal approximation test to test the hypothesis of a single population proportion, and the comparison of means was conducted using the Wilcoxon rank-sum test. All the tests are bilateral at a significance level of $\alpha = 0.05$. All statistical analyses were performed using SAS ver. 9.3 (SAS Institute Inc., Cary, NC, USA).

RESULTS

Gastrointestinal endoscopy was performed in 57 patients. Of these, 48 patients showed obvious mucosal lesions. These patients subsequently underwent further GI examinations, gastroesophageal scan, and 24-hour esophageal pH monitoring. The mucosal lesion sites of these patients were mainly

esophageal: reflux esophagitis (n=27), acute erosive esophagitis (n=8), esophageal ulcer (n=7), nodular gastritis (n=3), and gastric ulcer (n=3). Eight (16.7%) among these 48 patients also showed positive *Helicobacter pylori* infections. Accompanied endoscopic findings in addition with the above were chronic superficial gastritis in 13 cases, duodenogastric reflux in 12, chronic duodenitis in 10, and duodenal ulcer in one.

The prevalence rate of *H. pylori* in these children showed statistical significance ($p < 0.001$) compared with the rate in children with chronic abdominal pain (6.7%, 2011-2014) and in the general population of children (5.8%, 2011-2014) (Table 1) [7].

The most common cause of the chest pain was esophago-gastric diseases (64.0%; esophageal 56.0%, gastric 8.0%), followed by idiopathic cases (16.0%), cardiac diseases (12.0%), chest trauma (4.0%), respiratory diseases (2.7%), and psychiatric disease (1.3%). Among the children presenting with chest pain, the proportion finally diagnosed as esophago-gastric diseases was significantly higher than half ($p = 0.015$). The mean age of the children with esophago-gastric diseases was higher than that of the children with other diseases, but not statistically significant ($p = 0.056$) (Table 2).

Table 1. Endoscopic Findings Diagnosed as the Main Cause of Chest Pain (n=75)

Endoscopic finding	Diagnosed as the main cause of chest pain	Cases with positive <i>H. pylori</i>
Reflux esophagitis	27 (36.0)	1 (1.3)
Erosive esophagitis	8 (10.7)	1 (1.3)
Esophageal ulcer	7 (9.3)	1 (1.3)
Nodular gastritis	3 (4.0)	4 (5.3)
Gastric ulcer	3 (4.0)	1 (1.3)
Total	48 (64.0)	8 (10.7)*

Values are presented as number (%).

*The prevalence rate of *Helicobacter pylori* in these children showed statistical significance ($p < 0.001$) compared with that in the children with chronic abdominal pain (6.7%, 2011-2014) and those in general population children (5.8%, 2011-2014) [7]. Accompanied endoscopic findings in addition with the above (number of cases): chronic superficial gastritis, 13 duodenogastric reflux, 12 chronic duodenitis, 10 and duodenal ulcer, 1.

ECG was performed in all patients, of whom 21 (28.0%) showed abnormalities. Sinus arrhythmia was most commonly noted, right bundle branch block (RBBB) in three, with one patients each displaying ST elevation, left axis deviation, PR prolongation, left ventricular hypertrophy, and right axis deviation. Only two cases of sinus arrhythmia showed the same abnormality in 24-hour Holter monitoring; one case with RBBB and one case with PR prolongation were in ventricular septal defect (VSD) status with normal findings in echocardiography. Echocardiography performed in 34 children revealed nine abnormalities including three VSD, one aortic regurgitation, one atrial septal defect, one septal hypertrophy, and one high overload. Twenty four-hour Holter monitoring was performed in four, of whom two showed sinus arrhythmia (Table 3).

Chest X-ray performed in all patients revealed abnormalities in three, consisting of one each of enlargement of pulmonary conus, bronchopneumonia, and cardiomegaly due to VSD. The patient with en-

Table 2. Final Diagnoses of the Children Presenting with Chest Pain

Final diagnose	Case	Mean age (yr)
Esophago-gastric diseases	48 (64.0)*	10 [†]
Reflux esophagitis	27	9
Erosive esophagitis	8	11
Esophageal ulcer	7	11
Nodular gastritis	3	9
Gastric ulcer	3	11
Idiopathic disease	12 (16.0)	8
Cardiac diseases	9 (12.0)	8
Arrhythmia	5	7
Congenital heart disease	4	10
Chest trauma (contusion)	3 (4.0)	13
Respiratory disease	2 (2.7)	8
Bronchopneumonia	1	5
Pulmonary tuberculosis	1	11
Psychiatric disease	1 (1.3)	5
Total	75 (100)	

Values are presented as number (%) or number only.

*Among the children presenting with chest pain, the proportion of children diagnosed as esophago-gastric diseases finally is significantly higher than half ($p = 0.015$). [†]The mean age of the children with esophago-gastric diseases was higher than that of the children with other diseases, but not statistically significant ($p = 0.056$).

largement of pulmonary conus was diagnosed as pulmonary tuberculosis and in the echocardiography. Aortic regurgitation was also noted, and prompted a clinic with complaint of chest pain 5 years later (Table 4).

The patients whose primary diseases were esophageal or gastric had accompanying symptoms that most often were nausea (n=8) and vomiting (n=6). Patients whose primary diseases were not UGI diseases showed few GI symptoms. There was no distinct difference concerning general symptoms and thoracic symptoms between children with and without primary UGI diseases (Table 5).

After the final diagnoses of chest pain, patients were followed-up for at least 6 months to assess

whether symptoms disappeared with treatment, persisted, or recurred after initial improvement. Chest pain was improved with GI medicines in 37 patients (49.3%). Of these 37, 13 (35.1%) showed abnormalities in their cardiac examinations, which eventually showed little clinical correlation as a cause of chest pain. We continued to observe 24 patients (32.0%) who were not treated and did not display symptom recurrence; eight showed abnormal findings in their cardiac examination but did not require specific treatment. Three improved but the symptoms recurred, three (4.0%) suffered from sustained chest pain despite GI medicine, in whom abnormal cardiac examination results were already reported and might be the main reason for the persistence of

Table 3. Cardiologic Evaluation Findings of the Children with Chest Pain

Variable	Data
Electrocardiography	
Sinus arrhythmia	12 (16.0)*
RBBB [†]	3 (4.0)
ST-elevation	1 (1.3)
LAD	1 (1.3)
PR prolongation [‡]	1 (1.3)
LVH	1 (1.3)
RAD	1 (1.3)
Sinus bradyarrhythmia	1 (1.3)
Total	21 (28.0)
Echocardiography	
VSD	3 (4.0)
ASD	1 (1.3)
AR	1 (1.3)
Septal hypertrophy	1 (1.3)
High overload	1 (1.3)
Total	7 (9.3)
24-hr Holter monitoring	
Sinus arrhythmia	2 (2.7)
Total	2 (2.7)

Values are presented as number (%).

RBBB: right bundle branch block, LAD: left axis deviation, LVH: left ventricular hypertrophy, RAD: right axis deviation, VSD: ventricular septal defect, ASD: atrial septal defect, AR: aortic regurgitation, ECG: electrocardiography.

*Case number 12 and 25 showed sinus arrhythmia both in ECG and in 24-hr Holter monitoring. [†]Case number 33 showed RBBB in ECG and postop status of VSD in echocardiography. [‡]Case number 7 showed PR prolongation in ECG and postop status of VSD.

Table 4. Chest Radiologic Findings and the Related Cardiologic Problem of the Children

Chest Radiologic Findings	Case (n)	Cardiologic problem
Enlargement of left pulmonary conus (pulmonary tuberculosis)	1	Aortic regurgitation
Bronchopneumonia	1	None
Cardiomegaly	1	VSD

VSD: ventricular septal defect.

Table 5. Symptoms Accompanying Chest Pain

Accompanied symptom	UGI diseases as causes (n=48)	Non-UGI diseases as causes (n=27)
General symptoms	2 (2.7)	3 (4)
Dizziness and/or fatigue	0	2
Headache	1	1
Anorexia	1	0
Gastrointestinal symptoms	15 (20.0)	2 (2)
Vomiting	8	0
Epigastric pain	6	1
Chronic abdominal pain	1	1
Respiratory symptoms	4 (5.3)	5 (6.7)
Dyspnea	4	3
Cough	0	2
Total	21	10

Values are presented as number (percentage in 75 patients) or number only.

UGI, upper gastrointestinal.

Table 6. Post-diagnosis Clinical Courses of the Changes of Chest Pain of the Children

Clinical courses	No. of cases (%)	No. of cases with abnormal results of cardiopulmonary evaluations in the patients belonging to each clinical course
Improved with GI medicine	37 (49.3)	13 (35.1)
Observation, no treatment	24 (32.0)	8 (33.3)
Improved with GI medicine →Recurred	3 (4.0)	0 (0)
Sustained pain despite GI medicine	3 (4.0)	3 (100.0)
Spontaneously improved	3 (4.0)	1 (33.3)
Lost follow-up	3 (4.0)	2 (66.7)
Psychiatric consult	2 (2.7)	0 (0)
Total	75 (100.0)	27

GI: gastrointestinal.

symptoms. In three patients chest pain spontaneously resolved without any treatment (Table 6).

DISCUSSION

Chest pain in children is not a frequent symptom, as in adults. Pediatric chest pain usually present around 10-years-of-age [1,8-10], and females present more commonly than males. In this study, the majority of patients were 9 to 14-years-of-age (57.3%) and 6-years-of-age (14.7%). The male and female ratio was about 4:3.

Children with chest pain are usually referred to pediatric cardiologists, but chest pain is not an important or frequent symptom in children with cardiac diseases [4,11,12]. Careful history-taking, physical examination, and diverse laboratory investigations are required to determine the etiology of chest pain [13-15]. Musculoskeletal diseases (trauma, severe muscle sprain), respiratory diseases (pneumonia, pleuritis, asthma, chronic cough), UGI diseases (esophagitis, gastrointestinal reflux disease, foreign body, esophageal spasm, peptic ulcer disease), cardiovascular diseases (cardiomyopathy, endocarditis, valvular diseases, myocardial ischemia), and idiopathic causes are included in the differential diagnosis [4]. Idiopathic chest pain is most common in girls around 10-years-of-age [16].

Rowe et al. [17] studied clinical characteristics and causes of chest pain in 325 children visiting the

emergency room. The most common cause was idiopathic (33%), followed by chest wall pain (28%), respiratory disease (19%), trauma (15%), and psychogenic (5%). Very few patients (2%) were hospitalized and there were no patients with cardiovascular disease. There have been several studies regarding the causes of pediatric chest pain; in most of these UGI endoscopic examinations were seldom done so detailed explanations concerning the primary problems are lacking.

Chest pain caused by cardiovascular disease is not common in children with an incidence of 4% to 6% [18]. Mitral regurgitation is a major cardiac cause of chest pain in children, and is accompanied by vertigo, dyspnea, anxiety, and palpitations [19]. In this study, diverse examinations were performed to investigate cardiovascular disease, but clinically pathologic cardiac disease was not a major cause.

Several recent studies have suggested that UGI disease should be included in the differential diagnosis of chest pain [6,20-23]. Hsia et al. [21] performed UGI endoscopy and esophageal manometry in 100 patients with non-cardiac chest pain and diagnosed 24 cases of esophagitis, 18 cases of gastritis or duodenitis, 14 cases of esophageal hernia, and six cases of peptic ulcer disease, and recommended specific examinations for UGI disease in non-cardiac chest pain. Berezin et al. [22] diagnosed UGI diseases in 78% of 27 patients whose causes were specific; these patients showed clinical improvement after

the treatment of the UGI diseases. In this study, UGI disease was suspected in 18 children, and evaluations specific for the UGI diseases were performed. Specific UGI diseases were diagnosed in all these children and the symptoms improved after proper treatment. Lipsitz et al. [24] reported that psychosomatic anxiety disorders could be major causes of non-cardiac chest pain in children; however, further evaluation for UGI disease was not done. The authors reported the diagnostic efficacies of several examinations for pediatric chest pains in 2003 [25]. In the former study, the evaluation of UGI diseases were performed only in 18% of patients of the overall patient study group, and idiopathic cases were most common. Performing more examinations to determine whether UGI disease is a cause of cardiac pain, as UGI diseases were shown to be major causes of chest pain in children in this study. That is, the idiopathic causes of chest pain were less than 20%, while UGI disease caused chest pain in 40% of all cases in this study.

In our study, the incidence of *H. pylori* among the children with UGI diseases was higher than that among those with chronic abdominal pain in Korean children. We could not find our similar study performed in Korean children, but it was already reported that *H. pylori* infection have a important role in non-cardiac chest pain from Asia-Pacific survey [26].

Study limitations include the retrospectively design, lack of complete laboratory examinations in some patients, and failure of a specific treatment for certain organ disease to alleviate patient suffering. More differential diagnostic plans are needed, and should be investigated using a well-designed protocol.

In conclusion, careful history taking, physical examination, and proper clinical interventions are usually required to determine the diverse causes of chest pain. GI evaluation, including UGI endoscopic examination, is recommended to diagnose esophageal and gastric diseases in patients with non-cardiac chest pain.

REFERENCES

1. Driscoll DJ, Glicklich LB, Gallen WJ. Chest pain in children: a prospective study. *Pediatrics* 1976;57:648-51.
2. Fyfe DA, Moodie DS. Chest pain in pediatric patients presenting to a cardiac clinic. *Clin Pediatr (Phila)* 1984;23:321-4.
3. Pantell RH, Goodman BW Jr. Adolescent chest pain: a prospective study. *Pediatrics* 1983;71:881-7.
4. Bernstein D. History and physical examination. In: Kliegman RM, Stanton BF, St Geme JW, Schor NF, eds. *Nelson textbook of pediatrics*. 20th ed. Philadelphia: Elsevier Saunders, 2015:2163-70.
5. Yoon KL. Chest pain in children and adolescents. *J Korean Med Assoc* 2010;53:407-14.
6. Sabri MR, Ghavanini AA, Haghighat M, Imanieh MH. Chest pain in children and adolescents: epigastric tenderness as a guide to reduce unnecessary work-up. *Pediatr Cardiol* 2003;24:3-5.
7. Jang KM, Choe BH, Choe JY, Hong SJ, Park HJ, Chu MA, et al. Changing prevalence of helicobacter pylori infections in korean children with recurrent abdominal pain. *Pediatr Gastroenterol Hepatol Nutr* 2015;18:10-6.
8. Asnes RS, Santulli R, Bemporad JR. Psychogenic chest pain in children. *Clin Pediatr (Phila)* 1981;20:788-91.
9. Raiola G, Galati MC, De Sanctis V, Salerno D, Arcuri VM, Mussari A. Chest pain in adolescents. *Minerva Pediatr* 2002;54:623-30.
10. Selbst SM. Chest pain in children. *Pediatrics* 1985;75:1068-70.
11. Swenson JM, Fischer DR, Miller SA, Boyle GJ, Ettetdgui JA, Beerman LB. Are chest radiographs and electrocardiograms still valuable in evaluating new pediatric patients with heart murmurs or chest pain? *Pediatrics* 1997;99:1-3.
12. Kim JH, Moon HK, Jun JG. Clinical study of chest pain in children. *J Korean Pediatr Soc* 1990;33:1526-32.
13. Leung AK, Robson WL, Cho H. Chest pain in children. *Can Fam Physician* 1996;42:1156-60, 1163-4.
14. Selbst SM. Consultation with the specialist. Chest pain in children. *Pediatr Rev* 1997;18:169-73.
15. Evangelista JA, Parsons M, Renneburg AK. Chest pain in children: diagnosis through history and physical examination. *J Pediatr Health Care* 2000;14:3-8.
16. Selbst SM, Ruddy RM, Clark BJ, Henretig FM, Santulli T Jr. Pediatric chest pain: a prospective study. *Pediatrics* 1988;82:319-23.
17. Rowe BH, Dulberg CS, Peterson RG, Vlad P, Li MM. Characteristics of children presenting with chest pain

- to a pediatric emergency department. *CMAJ* 1990;143:388-94.
18. Kocis KC. Chest pain in pediatrics. *Pediatr Clin North Am* 1999;46:189-203.
 19. Woolf PK, Gewitz MH, Berezin S, Medow MS, Stewart JM, Fish BG, et al. Noncardiac chest pain in adolescents and children with mitral valve prolapse. *J Adolesc Health* 1991;12:247-50.
 20. Berezin S, Medow MS, Glassman MS, Newman LJ. Esophageal chest pain in children with asthma. *J Pediatr Gastroenterol Nutr* 1991;12:52-5.
 21. Hsia PC, Maher KA, Lewis JH, Cattau EL Jr, Fleischer DE, Benjamin SB. Utility of upper endoscopy in the evaluation of noncardiac chest pain. *Gastrointest Endosc* 1991;37:22-6.
 22. Berezin S, Medow MS, Glassman MS, Newman LJ. Chest pain of gastrointestinal origin. *Arch Dis Child* 1988;63:1457-60.
 23. Katz PO. Approach to the patient with unexplained chest pain. *Semin Gastrointest Dis* 2001;12:38-45.
 24. Lipsitz JD, Masia C, Apfel H, Marans Z, Gur M, Dent H, et al. Noncardiac chest pain and psychopathology in children and adolescents. *J Psychosom Res* 2005;59:185-8.
 25. Shin SA, Kim YJ, Lee JW, Kim NS, Moon SJ. Clinical evaluation and diagnosis of children with chest pain. *J Korean Pediatr Soc* 2003;46:1248-52.
 26. Cheung TK, Lim PW, Wong BC. Noncardiac chest pain--an Asia-Pacific survey on the views of primary care physicians. *Dig Dis Sci* 2007;52:3043-8.