

MULTI-PARAMETERIZED SCHWARZ ALTERNATING METHOD FOR 3D-PROBLEM[†]

SANG-BAE KIM

ABSTRACT. The convergence rate of a numerical procedure based on Schwarz Alternating Method(SAM) for solving elliptic boundary value problems depends on the selection of the interface conditions applied on the interior boundaries of the overlapping subdomains. It has been observed that the *Robin condition* (mixed interface condition), controlled by a parameter, can optimize SAM's convergence rate. In [7], one formulated the *multi-parameterized SAM* and determined the optimal values of the multi-parameters to produce the best convergence rate for one-dimensional elliptic boundary value problems. Two-dimensional implementation was presented in [8]. In this paper, we present an implementation for three-dimensional problem.

AMS Mathematics Subject Classification : 65N35, 65N05, 65F10.

Key words and phrases : elliptic partial differential equations, Schwarz alternating method, Jacobi iterative methods.

1. Introduction

Schwarz-type alternating methods have become some of the most important approaches in domain decomposition techniques for solution of the boundary value problems (BVP's). These methods are based on a decomposition of the BVP domain into overlapping subdomains. The original BVP is reduced to a set of *smaller* BVP's on a number of subdomains with appropriate *interface conditions* on the interior boundaries of the overlapping areas, whose solutions are coupled through some iterative scheme to produce an approximation of the solution of the original BVP. It is known [1], [6] that under certain conditions the sequence of the solutions of the subproblems converges to the solution of the original problem.

One of the objectives of this research is to study a class of Schwarz alternating methods (SAM's) whose interface conditions are parameterized and estimate the

Received September 10, 2014. Revised October 30, 2014. Accepted November 5, 2014.

[†]This work was supported by Hannam University Research Fund 2012

© 2015 Korean SIGCAM and KSCAM.

values of the parameters involved that speed up the convergence of these methods for a class of BVP's. In the context of elliptic BVP's the most commonly used *interface conditions* are of Dirichlet type. For this class of numerical SAM, there are several studies about the convergence, which include [10], [12], [15], [16], [13], [2], [11], [17]. The effect of parameterized mixed interface conditions has been considered by a number of researchers [3], [14], [5], [18]. Among them, Tang proposed a generalized Schwarz splitting [18]. The main part of his approach to the solution of a BVP is to use the mixed boundary condition, known as *Robin condition*,

$$B(u) = \omega u + (1 - \omega) \frac{\partial u}{\partial n} \quad (1)$$

on the artificial boundaries. In [5], a multi-parameter SAM is formulated in which the mixed boundary conditions

$$B_i(u) = \omega_i u + (1 - \omega_i) \frac{\partial u}{\partial n} \quad (2)$$

are controlled by a distinct parameter ω_i for the i -th overlapping area. Fourier analysis is applied to determine the values of ω_i parameters that make the convergence factor of SAM be zero.

In [7], one formulated a multi-parameter SAM at the matrix level where the parameters α_i are used to impose mixed interface conditions. The relation between the parameters α_i and ω_i is given by

$$\alpha_i = \frac{1 - \omega_i}{1 - \omega_i + \omega_i h} \quad (3)$$

(Refer to [7]), where h is the grid size. One determined analytically the optimal values of α_i 's for one-dimensional(1-dim) boundary value problems, which minimize the spectral radius of the block Jacobi iteration matrix associated with the SAM matrix.

For two-dimensional (2-dim) boundary value problems [8], we used distinct multi-parameter $\alpha_{i,j}$ for each j -th grid point of the i -th interfaces of the subdomains to get the best convergence, while we used a fixed parameter α_i along the i -th interfaces of the subdomains in the previous paper [7].

In this paper, we consider the case of three-dimensional (3-dim) problem. Here we also use distinct multi-parameter $\alpha_{i,j,k}$ for each (j,k) -th grid point of the i -th interfaces of the subdomains to get successfully the best convergence.

In section 2, we summarize the result of the multi-parameterized SAM on 1-dim problem, which has been presented in [7] and is necessary for notations in the following section. In section 3, we formulate the multi-parameterized SAM on 3-dim problem where we impose distinct parameters on each grid point on the interfaces of the subdomains. We show that the 3-dim case can be reduced to the one-dimensional ones and obtain the optimal values of the multi-parameters which minimize the spectral radius of the block Jacobi iteration matrix associated with the SAM matrix of 3-dim problem.




FIGURE 1. An example of the κ -way decomposition of the domain of one dimensional boundary value problem (4).

2. Multi-Parameterized Schwarz Splitting for 1-dim problem

We consider the two-point boundary value problem:

$$\begin{aligned} -u''(t) + q u(t) &= f(t), \quad t \in (0, 1) \\ u(0) &= a_0, \quad u(1) = a_1, \end{aligned} \quad (4)$$

with $q \geq 0$ being a constant. We will formulate a numerical instance of SAM based on a κ -way decomposition (i.e. the number of subdomains is κ) of the problem domain. An example of κ -way decomposition is depicted in Figure 1.

Let $T_j(x, y, z)$ be a $j \times j$ tridiagonal matrix such that

$$T_j(x, y, z) = \begin{bmatrix} x & -1 & 0 & \cdots & 0 \\ -1 & y & -1 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & -1 & y & -1 \\ 0 & \cdots & 0 & -1 & z \end{bmatrix} \quad (5)$$

and let

$$T_j(x) \equiv T_j(x, x, x). \quad (6)$$

If we discretize the problem (4) by a second order central divided difference discretization scheme with a uniform grid of mesh size $h = \frac{1}{n+1}$, we obtain a linear system

$$Ax = f \quad (7)$$

where $A = T_n(\beta)$ with $\beta = 2 + qh^2$.

If we consider 3-way ($\kappa = 3$) decomposition, then $Ax = f$ has three overlapping diagonal blocks as follows.

$$\begin{bmatrix} \boxed{T_{m-l} \quad -F} & 0 & 0 & 0 \\ -E & \boxed{T_l} & -F & 0 \\ 0 & -E & T_{m-2l} & -F \\ 0 & 0 & -E & \boxed{T_l} & -F \\ 0 & 0 & 0 & -E & T_{m-l} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} f_1 \\ f_2 \\ f_3 \\ f_4 \\ f_5 \end{bmatrix} \quad (8)$$

where $T_j = T_j(\beta, \beta, \beta)$ in (5) and m and l are the numbers of nodes in each subdomain and the overlapping regions, respectively, such that $l < \frac{m-1}{2}$. In (8), the matrix E have zero elements everywhere except for a 1 at the rightmost top position and the matrix F have zero elements everywhere except for a 1 at the

leftmost bottom position. So the matrices E and F have compatible sizes with the following forms.

$$E = \begin{bmatrix} 0 & \cdots & 0 & 1 \\ 0 & \cdots & 0 & 0 \\ \vdots & & \vdots & \vdots \\ 0 & \cdots & 0 & 0 \end{bmatrix}, F = \begin{bmatrix} 0 & 0 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & 0 \\ 1 & 0 & \cdots & 0 \end{bmatrix} \quad (9)$$

The numerical version of SAM [17] for the problem (4) is equivalent to a *block Gauss-Seidel iteration procedure* for a new linear system, called *Schwarz Enhanced Matrix Equation*,

$$\tilde{A}\tilde{x} = \tilde{f} \quad (10)$$

where

$$\tilde{A} = \tilde{A}(\beta) = \begin{bmatrix} T_{m-l} & -F & 0 & 0 & 0 & 0 & 0 \\ -E & T_l & 0 & -F & 0 & 0 & 0 \\ -E & 0 & T_l & -F & 0 & 0 & 0 \\ 0 & 0 & -E & T_{m-2l} & -F & 0 & 0 \\ 0 & 0 & 0 & -E & T_l & 0 & -F \\ 0 & 0 & 0 & -E & 0 & T_l & -F \\ 0 & 0 & 0 & 0 & 0 & -E & T_{m-l} \end{bmatrix}, \tilde{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_2' \\ x_3 \\ x_4 \\ x_4' \\ x_5 \end{bmatrix}, \tilde{f} = \begin{bmatrix} f_1 \\ f_2 \\ f_2' \\ f_3 \\ f_4 \\ f_4' \\ f_5 \end{bmatrix}. \quad (11)$$

$\tilde{A}(\beta)$ means that \tilde{A} is a function of β . Note that the solution x of (7) is obtained from the solution \tilde{x} of (10), vice versa. In [18], it is shown that a good choice of the splitting of T_l 's can significantly improve the convergence of SAM. Applying for some splittings of T_l 's into \tilde{A} in (11), we have a new equation

$$A'\tilde{x} = \tilde{f} \quad (12)$$

with

$$A' = A'(\beta, \alpha_1, \alpha_2) = \begin{bmatrix} T_{m-l} & -F & 0 & 0 & 0 & 0 & 0 \\ -E & B_1 & C_1 & -F & 0 & 0 & 0 \\ -E & C_1' & B_1' & -F & 0 & 0 & 0 \\ 0 & 0 & -E & T_{m-2l} & -F & 0 & 0 \\ 0 & 0 & 0 & -E & B_2 & C_2 & -F \\ 0 & 0 & 0 & -E & C_2' & B_2' & -F \\ 0 & 0 & 0 & 0 & 0 & -E & T_{m-l} \end{bmatrix} \quad (13)$$

where $B_i, C_i', i = 1, 2$ are some matrices such that $(B_i - C_i')$ is non-singular and

$$T_l = B_i + C_i = B_i' + C_i', \quad i = 1, 2. \quad (14)$$

Note that two linear system (10) and (12) are equivalent in the sense that they have the same solutions. If C_i' and C_i are chosen such that they are the $l \times l$ matrices with all zero entries except for an α_i in the positions $(1, 1)$ and (l, l) , respectively, as follows,

$$C_i' = \begin{bmatrix} \alpha_i & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix}, C_i = \begin{bmatrix} 0 & \cdots & 0 & 0 \\ \vdots & & \vdots & \vdots \\ 0 & \cdots & 0 & 0 \\ 0 & \cdots & 0 & \alpha_i \end{bmatrix}, \quad (15)$$

the resulting matrix A' is given as follows

$$A' = A'(\beta, \alpha_1, \alpha_2) = \begin{bmatrix} T_m(\beta, \beta, \beta - \alpha_1) & -F_1' & 0 \\ -E_1' & T_m(\beta - \alpha_1, \beta, \beta - \alpha_2) & -F_2' \\ 0 & -E_2' & T_m(\beta - \alpha_2, \beta, \beta) \end{bmatrix} \quad (16)$$

where $T_m(x, y, z)$'s are $m \times m$ matrices defined in (5) and E_i' 's are the $m \times m$ matrices with zero elements everywhere except that

$$\begin{aligned} (1, m-l) \text{-th entry} &= 1, \\ (1, m-l+1) \text{-th entry} &= -\alpha_i \end{aligned}$$

and F_i' 's are the $m \times m$ matrices with zero elements everywhere except that

$$\begin{aligned} (m, l+1) \text{-th entry} &= 1, \\ (m, l) \text{-th entry} &= -\alpha_i. \end{aligned}$$

If the number of subdomains κ is more than 3, the matrix A' is a block $\kappa \times \kappa$ matrix of the form

$$A' = A'(\beta, \mathbf{a}) = \begin{bmatrix} G_1 & -F_1' & 0 & 0 & \cdots & 0 \\ -E_1' & G_2 & -F_2' & 0 & \cdots & 0 \\ \vdots & & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & -E_{\kappa-2}' & G_{\kappa-1} & -F_{\kappa-1}' \\ 0 & \cdots & 0 & 0 & -E_{\kappa-1}' & G_{\kappa} \end{bmatrix} \quad (17)$$

where $\mathbf{a} = (\alpha_0, \alpha_1, \alpha_2, \dots, \alpha_{\kappa})$ with $\alpha_0 = \alpha_{\kappa} = 0$ and G_i 's are defined as

$$G_i = T_m(\beta - \alpha_{i-1}, \beta, \beta - \alpha_i), \quad i = 1, 2, \dots, \kappa. \quad (18)$$

We call the matrix A' as *Multi-Parameterized Enhanced Matrix*. If we define

$$M = M(\beta, \mathbf{a}) = \begin{bmatrix} G_1 & 0 & \cdots & 0 \\ 0 & G_2 & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \cdots & 0 & G_{\kappa} \end{bmatrix} \quad \text{and} \quad N = N(\beta, \mathbf{a}) = \begin{bmatrix} 0 & F_1' & 0 & 0 & \cdots & 0 \\ E_1' & 0 & F_2' & 0 & \cdots & 0 \\ \vdots & & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & E_{\kappa-2}' & 0 & F_{\kappa-1}' \\ 0 & \cdots & 0 & 0 & E_{\kappa-1}' & 0 \end{bmatrix} \quad (19)$$

with $\mathbf{a} = (\alpha_0, \alpha_1, \alpha_2, \dots, \alpha_{\kappa})$, then we can write the multi-parameterized enhanced matrix A' as


$$A' = M - N \quad (20)$$

which is called a *Multi-Parameterized Schwarz Splitting (MPSS)*.

The convergence behavior of MPSS depends on the spectral radius of the following block Jacobi matrix

$$J = M^{-1}N = \begin{bmatrix} 0 & G_1^{-1}F_1' & 0 & 0 & \cdots & 0 \\ G_2^{-1}E_1' & 0 & G_2^{-1}F_2' & 0 & \cdots & 0 \\ 0 & G_3^{-1}E_2' & 0 & G_3^{-1}F_3' & \cdots & 0 \\ \vdots & & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & G_{\kappa-1}^{-1}E_{\kappa-2}' & 0 & G_{\kappa-1}^{-1}F_{\kappa-1}' \\ 0 & \cdots & 0 & 0 & G_{\kappa}^{-1}E_{\kappa-1}' & 0 \end{bmatrix}. \quad (21)$$

Note that J is a function of the parameters α_i 's, which correspond to the parameters ω_i 's in the mixed interface condition (2), respectively. The convergence

FIGURE 2. A 3-way splitting of the unit cube Ω .

rate of SAM can be optimized by controlling these parameters α_i 's. In [7], one determined the optimal values of the multi-parameter α_i 's that make the spectral radius of the block Jacobi matrix J in (21) to be zero. The result of [7] is presented in the following theorem.

Theorem 2.1. *Let $\theta = \cosh^{-1}(\frac{\beta}{2})$ with $\beta = 2 + qh^2$ and let $p \in \{1, 2, \dots, \kappa - 1\}$ and let*

$$\Theta(x) \equiv \begin{cases} \sinh(x\theta), & \theta > 0 \\ x, & \theta = 0. \end{cases}$$

If the values α_i , $i = 0, 1, \dots, \kappa$, are given by

$$\alpha_0 = \alpha_\kappa = 0$$

$$\alpha_i = \frac{\Theta(m-l) - \alpha_{i-1} \Theta(m-l-1)}{\Theta(m-l+1) - \alpha_{i-1} \Theta(m-l)}, \quad i = 1, 2, \dots, p,$$

$$\alpha_i = \frac{\Theta(m-l) - \alpha_{i+1} \Theta(m-l-1)}{\Theta(m-l+1) - \alpha_{i+1} \Theta(m-l)}, \quad i = p + 1, \dots, \kappa - 1,$$

then the spectral radius of the block Jacobi matrix J in (21) is zero.

3. Multi-Parameterized Schwarz Splitting for 3-dim Problem

Consider the three-dimensional boundary value problem

$$\begin{aligned} -\nabla^2 u(x) + q u(x) &= f(x), & x \in \Omega, \\ u(x) &= g(x), & x \in \Gamma \end{aligned} \quad (22)$$

where Γ is the boundary of $\Omega \equiv (0, 1) \times (0, 1) \times (0, 1)$ and $q \geq 0$ is a constant. We formulate a SAM based on a κ -way splitting of the domain Ω , i.e., we decompose our domain into κ overlapping subdomains Ω_i along the x_1 -axis and make a strip-type decomposition of the cube domain Ω (for instance, see Figure 2). Next we apply the interface conditions on the two interior boundaries between subdomains Ω_i and Ω_{i+1} . Let ℓ be the length of the overlap in x_1 -direction and

η be the length of each subdomain in the same direction. Figure 2 depicts such a 3-way splitting of the unit cube Ω .

To begin our analysis we use a 7-point finite difference discretization scheme with uniform grid of mesh size $h = \frac{1}{n+1}$ on all of x_1 -, x_2 - and x_3 -axes and discretize the BVP in (22) to obtain a linear system of the form

$$Bx = f. \quad (23)$$

The natural ordering of the nodes is adopted starting from the origin and going in the x_3 -direction first, and then x_2 - and x_1 -direction in turn, so that the resulting matrix A can be partitioned into block matrices corresponding to the subdomains, respectively. Using tensor product notation \otimes (See [4], and [9] in which tensor products in connection with BVP's were introduced.), the matrix B in (23) can be written as

$$B = T_n(\beta) \otimes I_n \otimes I_n + I_n \otimes T_n(2) \otimes I_n + I_n \otimes I_n \otimes T_n(2) \quad (24)$$

where $T_j(x)$ is defined in (6) and $\beta = 2 + qh^2$.

Define $l+1 = \frac{\ell}{h}$ and $m+1 = \frac{\eta}{h}$ such that $n = \kappa m - l(\kappa - 1)$ and $l < \frac{m-1}{2}$. The numerical version of SAM for the problem (22) is equivalent to a *block Gauss-Seidel iteration procedure* for a new linear system, called the *Schwarz Enhanced Matrix Equation*,

$$\tilde{B}\tilde{x} = \tilde{f} \quad (25)$$

with

$$\tilde{B} = \tilde{A}(\beta) \otimes I_n \otimes I_n + I_{\kappa m} \otimes T_n(2) \otimes I_n + I_{\kappa m} \otimes I_n \otimes T_n(2) \quad (26)$$

where $I_{\kappa m}$ is the $\kappa m \times \kappa m$ identity matrix and $\tilde{A}(\beta)$ is the $\kappa \times \kappa$ block matrix as that defined in (11), which is the case of $\kappa = 3$. Note that each diagonal block in $\tilde{A}(\beta)$ is $m \times m$ matrix.

Let X_n be the $n \times n$ orthogonal matrix whose columns are the eigenvectors of the matrix $T_n(2)$. Since the eigenvalues of the matrix $T_n(2)$ are known to be $\gamma_i = 2 + 2 \cos(\frac{i\pi}{n+1})$, $i = 1, 2, \dots, n$, we can write

$$X_n^T T_n(2) X_n = D_n \equiv \text{diag}(\gamma_1, \gamma_2, \dots, \gamma_n). \quad (27)$$

Let $X = I_{\kappa m} \otimes I_n \otimes X_n$, then its inverse is given by $X^{-1} = I_{\kappa m} \otimes I_n \otimes X_n^T$, so we have

$$\begin{aligned} X^{-1} \tilde{B} X &= (I_{\kappa m} \otimes I_n \otimes X_n^T) (\tilde{A}(\beta) \otimes I_n \otimes I_n) (I_{\kappa m} \otimes I_n \otimes X_n) \\ &\quad + (I_{\kappa m} \otimes I_n \otimes X_n^T) (I_{\kappa m} \otimes T_n(2) \otimes I_n) (I_{\kappa m} \otimes I_n \otimes X_n) \\ &\quad + (I_{\kappa m} \otimes I_n \otimes X_n^T) (I_{\kappa m} \otimes I_n \otimes T_n(2)) (I_{\kappa m} \otimes I_n \otimes X_n) \\ &= (I_{\kappa m} \tilde{A}(\beta) I_{\kappa m}) \otimes I_n \otimes (X_n^T I_n X_n) \\ &\quad + I_{\kappa m} \otimes T_n(2) \otimes (X_n^T I_n X_n) \\ &\quad + I_{\kappa m} \otimes I_n \otimes (X_n^T T_n(2) X_n) \\ &= \tilde{A}(\beta) \otimes I_n \otimes I_n + I_{\kappa m} \otimes T_n(2) \otimes I_n + I_{\kappa m} \otimes I_n \otimes D_n. \end{aligned}$$

If P is the permutation matrix that maps

$$\begin{array}{l} \text{row} \quad (i-1)n^2 + (j-1)n + k \\ \text{into} \\ \text{row} \quad (k-1)n\kappa m + (i-1)n + j \end{array}$$

for $i = 1, 2, \dots, \kappa m$, $j = 1, 2, \dots, n$, $k = 1, 2, \dots, n$, then we have

$$\begin{aligned} \check{B} &= P^{-1}X^{-1}\check{B}XP \\ &= P^{-1}(\check{A}(\beta) \otimes I_n \otimes I_n)P + P^{-1}(I_{\kappa m} \otimes T_n(2) \otimes I_n)P \\ &\quad + P^{-1}(I_{\kappa m} \otimes I_n \otimes D_n)P \\ &= I_n \otimes \check{A}(\beta) \otimes I_n + I_n \otimes I_{\kappa m} \otimes T_n(2) + D_n \otimes I_{\kappa m} \otimes I_n \end{aligned} \quad (28)$$

Note that the solution \check{x} of linear system (25) is obtained by $\check{x} = XP\check{x}$ if we solve the linear system

$$\check{B}\check{x} = \check{f} \quad (29)$$

where $\check{f} = P^{-1}X^{-1}\tilde{f}$ with \tilde{f} in (25).

Likewise, using $Y = I_n \otimes I_{\kappa m} \otimes X_n$, we have

$$\begin{aligned} Y^{-1}\check{B}Y &= (I_n \otimes I_{\kappa m} \otimes X_n^T)(I_n \otimes \check{A}(\beta) \otimes I_n)(I_n \otimes I_{\kappa m} \otimes X_n) \\ &\quad + (I_n \otimes I_{\kappa m} \otimes X_n^T)(I_n \otimes I_{\kappa m} \otimes T_n(2))(I_n \otimes I_{\kappa m} \otimes X_n) \\ &\quad + (I_n \otimes I_{\kappa m} \otimes X_n^T)(D_n \otimes I_{\kappa m} \otimes I_n)(I_n \otimes I_{\kappa m} \otimes X_n) \\ &= I_n \otimes \check{A}(\beta) \otimes (X_n^T I_n X_n) + I_n \otimes I_{\kappa m} \otimes (X_n^T T_n(2) X_n) \\ &\quad + D_n \otimes I_{\kappa m} \otimes (X_n^T I_n X_n) \\ &= I_n \otimes \check{A}(\beta) \otimes I_n + I_n \otimes I_{\kappa m} \otimes D_n + D_n \otimes I_{\kappa m} \otimes I_n. \end{aligned}$$

If Q is the permutation matrix that maps

$$\begin{array}{l} \text{row} \quad (k-1)n\kappa m + (i-1)n + j \\ \text{into} \\ \text{row} \quad (k-1)n\kappa m + (j-1)\kappa m + i \end{array}$$

for $i = 1, 2, \dots, \kappa m$, $j = 1, 2, \dots, n$, $k = 1, 2, \dots, n$, then we have

$$\begin{aligned} \hat{B} &= Q^{-1}X^{-1}\check{B}XQ \\ &= Q^{-1}(I_n \otimes \check{A}(\beta) \otimes I_n)Q + Q^{-1}(I_n \otimes I_{\kappa m} \otimes D_n)Q \\ &\quad + Q^{-1}(D_n \otimes I_{\kappa m} \otimes I_n)Q \\ &= I_n \otimes I_n \otimes \check{A}(\beta) + I_n \otimes D_n \otimes I_{\kappa m} + D_n \otimes I_n \otimes I_{\kappa m} \\ &= \Phi[\check{A}(\zeta_{j,k})]_{\substack{j=1,2,\dots,n \\ k=1,2,\dots,n}} \end{aligned} \quad (30)$$

where $\zeta_{j,k} = \beta + \gamma_j + \gamma_k$ for $j = 1, 2, \dots, n$ and $k = 1, 2, \dots, n$ and

$$\begin{aligned} &\Phi[s_{(j,k)}]_{\substack{j=1,2,\dots,n \\ k=1,2,\dots,n}} \\ &= \text{diag}(\text{diag}(s_{(1,1)}, s_{(2,1)}, \dots, s_{(n,1)}), \\ &\quad \text{diag}(s_{(1,2)}, s_{(2,2)}, \dots, s_{(n,2)}), \\ &\quad \dots \\ &\quad \text{diag}(s_{(1,n)}, s_{(2,n)}, \dots, s_{(n,n)}) \\ & \quad) \end{aligned} \quad (31)$$

which represents a diagonal matrix of diagonal matrices of entries $s_{(j,k)}$'s.

Note that the solution \check{x} of linear system (29) is obtained by $\check{x} = YQ\hat{x}$ if we solve the linear system

$$\hat{B}\hat{x} = \hat{f} \quad (32)$$

where

$$\hat{f} = Q^{-1}Y^{-1}\check{f}$$

with \check{f} in (29), so that

$$\begin{aligned} \hat{f} &= Q^{-1}Y^{-1}\check{f} \\ &= Q^{-1}Y^{-1}(P^{-1}X^{-1}\tilde{f}) \end{aligned}$$

with \tilde{f} in (25). Finally \tilde{x} in (25) is computed by

$$\tilde{x} = XP\check{x} = XP(YQ\hat{x})$$

with \hat{x} in (32).

From (30) and (32), we know that the three-dimensional problem (25) is reduced to n^2 number of one dimensional problems

$$\tilde{A}(\zeta_{j,k}) = \hat{f}_{j,k}, \quad j = 1, 2, \dots, n, \quad k = 1, 2, \dots, n,$$

where $\hat{f}_{j,k}$ is the corresponding sub-vector of \hat{f} .

The *Multi-Parameterized Schwarz Enhanced Matrix* for \hat{B} in (30) is defined as

$$B' = \Phi[A'(\zeta_{j,k}, \mathbf{a})]_{\substack{j = 1, 2, \dots, n \\ k = 1, 2, \dots, n}} \quad (33)$$

where $A'(x, \mathbf{a})$ is defined in (17). If we let

$$\begin{aligned} M &= \Phi[M(\zeta_{j,k}, \mathbf{a})]_{\substack{j = 1, 2, \dots, n \\ k = 1, 2, \dots, n}} \\ N &= \Phi[N(\zeta_{j,k}, \mathbf{a})]_{\substack{j = 1, 2, \dots, n \\ k = 1, 2, \dots, n}} \end{aligned} \quad (34)$$

where $M(x, \mathbf{a})$ and $N(x, \mathbf{a})$ are defined in (19), then we can write the multi-parameterized enhanced matrix B' in (33) as

$$B' = M - N \quad (35)$$

which is called a *Multi-Parameterized Schwarz Splitting (MPSS)*. The convergence behavior of MPSS depends on the spectral radius of the following block Jacobi matrix

$$J = M^{-1}N = \Phi[L_{j,k}(\mathbf{a})]_{\substack{j = 1, 2, \dots, n \\ k = 1, 2, \dots, n}} \quad (36)$$

where

$$L_{j,k}(\mathbf{a}) = M(\zeta_{j,k}, \mathbf{a})^{-1} N(\zeta_{j,k}, \mathbf{a})$$

for $j = 1, 2, \dots, n$ and for $k = 1, 2, \dots, n$.

In [7], one failed to determine a parameter vector \mathbf{a} such that the spectral radius of the block Jacobi matrix J in (36) is zero because it is not possible to find such a parameter vector \mathbf{a} that makes all of the spectral radii of the diagonal blocks $L_{j,k}(\mathbf{a})$'s in (36) zero simultaneously.

So instead of fixed parameter \mathbf{a} , we adopt multi-parameter vector $\mathbf{a}_{j,k}$ for each diagonal block as follows

$$J = M^{-1}N = \Phi[L_{j,k}(\mathbf{a}_{j,k})] \quad \begin{matrix} j = 1, 2, \dots, n \\ k = 1, 2, \dots, n \end{matrix} \quad (37)$$

where

$$\mathbf{a}_{j,k} = (\alpha_{0,j,k}, \alpha_{1,j,k}, \alpha_{2,j,k}, \dots, \alpha_{\kappa,j,k}),$$

for $j = 1, 2, \dots, n$ and $k = 1, 2, \dots, n$. Note these triple indices (i, j, k) in multi-parameter $\alpha_{i,j,k}$ are related to the idea that one adopts variable parameter $\omega_i(y, z)$ instead of constant parameter ω_i in (2) along the i -th interface, i.e., we have

$$B_i(u) = \omega_i(y, z)u + (1 - \omega_i(y, z))\frac{\partial u}{\partial n} \quad (38)$$

as the mixed interface condition on the i -th interface boundary.

Now, using these triple-index multi-parameter $\alpha_{i,j,k}$'s, we have the following theorem for three-dimensional multi-parameterized Schwarz splitting $B' = M - N$ in (35).

Theorem 3.1. For $j = 1, 2, \dots, n$ and $k = 1, 2, \dots, n$, let $\theta_{j,k} = \cosh^{-1}(\frac{\zeta_{j,k}}{2})$ with $\zeta_{j,k}$ in (30) and let $p \in \{1, 2, \dots, \kappa - 1\}$ and let

$$\Theta_{j,k}(x) \equiv \begin{cases} \sinh(x\theta_{j,k}), & \theta > 0 \\ x, & \theta = 0. \end{cases}$$

If the values $\alpha_{i,j,k}$ for each $j = 1, 2, \dots, n$ and each $k = 1, 2, \dots, n$ and each $i = 0, 1, \dots, \kappa$ are given by

$$\begin{aligned} \alpha_{0,j,k} &= \alpha_{\kappa,j,k} = 0 \\ \alpha_{i,j,k} &= \frac{\Theta_{j,k}(m-l) - \alpha_{i-1,j,k}\Theta_{j,k}(m-l-1)}{\Theta_{j,k}(m-l+1) - \alpha_{i-1,j,k}\Theta_{j,k}(m-l)}, \quad i = 1, 2, \dots, p, \\ \alpha_{i,j,k} &= \frac{\Theta_{j,k}(m-l) - \alpha_{i+1,j,k}\Theta_{j,k}(m-l-1)}{\Theta_{j,k}(m-l+1) - \alpha_{i+1,j,k}\Theta_{j,k}(m-l)}, \quad i = p+1, \dots, \kappa-1, \end{aligned}$$

then the spectral radius of the block Jacobi matrix J in (37) is zero.

4. Numerical Experiments

In this section, we present a numerical experiment to prove the result of the previous section. we will compare the results of *Multi-Parameterized SAM (MP-SAM)* with those of the *Classical SAM*. Consider the following model problem

$$\begin{aligned} -\nabla^2 u(x, y, z) &= 0, \quad (x, y, z) \in \Omega = (0, 1) \times (0, 1) \times (0, 1), \\ u(x, y, z) &= f(x, y, z), \quad (x, y, z) \in \Gamma, \end{aligned} \quad (39)$$

where Γ is the boundary of Ω , with solution

$$u(x, y, z) = \sin(2\pi x) \cos(2\pi y) \sin(2\pi z).$$

In all the experiments, the vector with all its components 0.0 was used as initial guess of the solution vector. The *relative residual* r_κ is computed as the

TABLE 1. The classical *SAM* is applied to the BVP (39).

m	l	<i>relative residual</i> r_κ (<i>max(err)</i>)		
		$\kappa = 2$	$\kappa = 4$	$\kappa = 8$
8	1	8.4817E-02 (7.7862E-02)	6.3237E-02 (1.1697E-01)	5.8736E-02 (1.9661E-01)
8	4	1.6126E-02 (3.3959E-02)	7.6499E-04 (8.6187E-03)	2.4438E-03 (1.9430E-03)

TABLE 2. The *MPSAM* is applied to the BVP (39).

m	l	<i>relative residual</i> r_κ (<i>max(err)</i>)		
		$\kappa = 2$	$\kappa = 4$	$\kappa = 8$
8	1	2.7744E-16 (1.3282E-02)	4.1060E-16 (3.7107E-03)	6.0494E-16 (9.9877E-04)
8	4	3.4301E-16 (1.9391E-02)	3.2078E-16 (7.5565E-03)	5.0200E-16 (2.4514E-03)

ratio of ℓ_2 -norms of the residuals of the corresponding system of equations after κ iterations, i.e.,

$$r_\kappa = \frac{\|B'x^{(\kappa)} - \hat{f}\|_2}{\|B'x^{(0)} - \hat{f}\|_2}.$$

Table 1 shows the relative residuals of *SAM* computed after κ iterations for each number of subdomains ($\kappa = 2, 4, 8$), and number of local grids ($m = 8$) and minimum and half overlaps ($l = 1, 4$). Table 2 shows the performance of *MPSAM* under the same conditions. It shows the optimal convergence. Indeed, the relative residuals by *MPSAM* are less than 5.02×10^{-15} after κ iterations for the case of κ subdomains.

The convergence rate is very sensitive to the computed optimal value of parameter $\alpha_{i,j,k}$'s and the symmetric choice of them (i.e. Take $p = \lceil \kappa/2 \rceil$ in Theorem (3.1)) reduces the error propagation when we compute the optimal value of parameters $\alpha_{i,j,k}$'s.

REFERENCES

1. R. Courant and D. Hilbert, *Methods of mathematical physics, vol 2*, Willey, New York, 1962.
2. M. Dryja, *An additive Schwarz algorithm for two- and three-dimensional finite element elliptic problems*, Domain Decomposition Methods (T. Chan, R. Glowinski, J. Periaux, and O. Widlund, eds.), SIAM, 1989, pp. 168–172.
3. D.J. Evans, L.-S. Kang, Y.-P. Chen, and J.-P. Shao, *The convergence rate of the Schwarz alternating procedure (iv) : With pseudo-boundary relaxation factor*, Intern. J. Computer Math. **21** (1987), 185–203.
4. P.R. Halmos, *Finite-dimensional Vector Spaces*, Van Nostrand, Princeton, N.J., 1958.

5. L.-S. Kang, *Domain decomposition methods and parallel algorithms*, Second International Symposium on Domain Decomposition Methods for Partial Differential Equations (Philadelphia, PA) (Tony F. Chan, Roland Glowinski, Jacques Périaux, and Olof B. Widlund, eds.), SIAM, 1989, pp. 207–218.
6. L.V. Kantorovich and V.I. Krylov, *Approximate methods of higher analysis*, 4th ed., P. Noordhoff Ltd, Groningen, The Netherlands, 1958.
7. S.-B. Kim, A. Hadjidimos, E.N. Houstis, and J.R. Rice, *Multi-parametrized schwarz splittings for elliptic boundary value problems*, Math. and Comp. in Simulation **42** (1996), 47–76.
8. S.-B. Kim, *Two-dimensional Muti-Parameterized Schwarz Alternating Method*, J. Appl. Math. & Informatics **29** (2011), 161–171
9. R.E. Lynch, J.R. Rice, and D.H. Thomas, *Direct solution of partial difference equations by tensor products*, Numer. Math. **6** (1964), 185–189.
10. K. Miller, *Numerical analogs to the Schwarz alternating procedure*, Numer. Math. **7** (1965), 91–103.
11. J. Olinger, W. Skamarock, and W.P. Tang, *Schwarz alternating methods and its SOR accelerations*, Tech. report, Department of Computer Science, Stanford University, 1986.
12. G. Rodrigue, *Inner/outer iterative methods and numerical Schwarz algorithms*, J. Parallel Computing **2** (1985), 205–218.
13. G. Rodrigue and P. Saylor, *Inner/outer iterative methods and numerical Schwarz algorithms-ii*, Proceedings of the IBM Conference on Vector and Parallel Computations for Scientific Computing, IBM, 1985.
14. G. Rodrigue and S. Shah, *Pseudo-boundary conditions to accelerate parallel Schwarz methods*, Parallel Supercomputing : Methods, Algorithms, and Applications (New York) (G. Carey, ed.), Wiley, 1989, pp. 77–87.
15. G. Rodrigue and J. Simon, *A generalization of the numerical Schwarz algorithm*, Computing Methods in Applied Sciences and Engineering VI (Amsterdam, New York, Oxford) (R. Glowinski and J. Lions, eds.), North-Holland, 1984, pp. 273–281.
16. G. Rodrigue and J. Simon, *Jacobi splitting and the method of overlapping domains for solving elliptic PDE's*, Advances in Computer Methods for Partial Differential Equations V (R. Vichnevetsky and R. Stepleman, eds.), IMACS, 1984, pp. 383–386.
17. W.P. Tang, *Schwarz Splitting and Template Operators*, Ph.D. thesis, Department of Computer Science, Stanford University, 1987.
18. W.P. Tang, *Generalized Schwarz splittings*, SIAM J. Sci. Stat. Comput. **13** (1992), 573–595.

Sang-Bae Kim received a Ph.D from Purdue University, USA, in 1993. He is a professor of Department of Mathematics, Hannam University, Korea. His research interests are numerical linear algebra, scientific computing.

Department of Mathematics, Hannam University, Daejeon 306-791, Korea
e-mail: sbk@hnu.kr