

Rehabilitative Effect of Intramuscular Electrostimulation after Reconstruction of Medial Patellar Luxation in Small Sized Dog

Shinho Lee, Joo-Myoung Lee, Hyunjung Park, Sangcheol Won and Jongtae Cheong¹

College of Veterinary Medicine · Veterinary Medical Research Institute, Jeju National University, Jeju 690-756, Korea

(Accepted: January 28, 2015)

Abstract : Medial patellar luxation (MPL) of dog is one of the most common joint diseases. This study performed to know rehabilitative effect after reconstruction as application with intramuscular electrostimulation (IMES) on the cranial part of sartorius muscle. In this study, the 12 dogs with operated medial luxation divided into electrostimulation group (n = 8) and no electrostimulation group (n = 4) measured affected sided range of motion, muscle mass, lameness score and weight bearing for hospitalization 2 weeks. IMES group showed significant results after operation on 5th day ($p < 0.01$). Except lameness score, IMES group showed significant results on 10th day ($p < 0.05$). As results, in small sized dog after reconstruction of MPL, IMES on the cranial part of sartorius muscle considered a great help in the initial rehabilitation.

Key words : intramuscular electrostimulation (IMES), sartorius muscle, patellar luxation, dog.

Introduction

The most common diagnosis of patellar luxation is congenital or developmental medial patellar luxation (MPL) (20,24, 25). In small-breed dogs, luxation was medial in 99% of cases, and was lateral in only 1%; among the affected dogs, 35% were males and 65% females (1). Most patients with patellar luxation have associated musculoskeletal abnormalities, such as medial displacement of the quadriceps muscle group, lateral torsion of the distal femur, a lateral bowing of the distal one third of the femur, femoral epiphyseal dysplasia, rotational instability of the stifle joint, or tibial deformity (6). Femoropatellar instability, anteversion, angle of femur and inclination leading to patellar luxation are common causes of lameness in dogs (6,17,27).

The sartorius muscles may pay an important role in the medial tensile force which caused the patella to luxate medially. Morphometric analyses of the pelvis showed that the origin of the cranial head of the sartorius muscle lies significantly more medially in dogs with patellar luxation than in normal dogs (11,12,18). This theory supported transplantation of the cranial part of the sartorius muscle for the treatment of patellar luxation (10).

In case observed intermittent episodes of a toe-touching lameness of the left pelvic limb, physical therapy was performed, including underwater treadmill exercises and electrical stimulation of the vastus lateralis muscle every 3 to 4 days for 1 month. Three months postoperatively, no lameness was seen at a walk, but occasional flexing of the limb occurred at a trot (3). In another study, After reconstruction of MPL, the animal is restricted to short leash walks for 4 to 6 weeks, and

the owner is encouraged to provide current post-operative rehabilitation protocols such as massage, passive range of motion (ROM) and swimming to regain the strength of the muscle mass (6,24). There were other reports for rehabilitation of MPL but there was no rehabilitation of shorten cranial part of sartorius muscle.

Electroacupuncture was first used in China during 1930s (22). It was getting the usual, especially in the area of physical rehabilitation (21). Electrostimulation in animals had clearly established growth, repair and remodeling of hard and soft tissues and often assumed to augment regeneration of various tissues (5). Stimulation in any pain of neurological origin is suppression of signals through a reflex arc. In addition to functioning as an alternative method of pain relief, there is evidence that electrical stimulation encourages muscle contractions and increased blood flow to the targeted tissues (7,8). Lower tetanic frequencies in the range of 35~50 Hz reduce muscle fatigue while still providing strong muscle contractions (16).

This study hypothesized that the dogs applied on shorten cranial part of sartorius muscle with intramuscular stimulation (IMES) help to return to activity of daily living quickly. The purpose of this study is to ascertain the rehabilitative effect of IMES treatment in small sized dog with reconstructed MPL.

Materials and Methods

Animals

The dogs were belonged to different breed, sex and weight, ranging from 9 months to 12 years old. The breeds were Maltese (n = 6), Poodle (n = 1), crossbreed (n = 1), Pomeranian (n = 1), Yorkshire Terriers (n = 2) and Shih Tzu (n = 1). There were 6 males and 6 females, 6 of which were sexually intact;

¹Corresponding author.
E-mail : cjt123@jejunu.ac.kr

Table 1. Description of Dogs with Medial Patellar Luxation

Group	No.	Breed	Sex*	Age**	B. W. (kg)	Surgery lesion (grade)
IMES	1	Maltese	F	10 m	2.7	Left (II)
	2	Maltese	S	5 y	3.8	Right (II)
	3	Poodle	C	9 m	2.7	Right (I)
	4	Crossbreed	F	5 y	6.8	Right (II)
	5	Yorkshire terrier	M	10 m	3.0	Right (III)
	6	Maltese	M	5 y	4.4	Right (III)
	7	Yorkshire terrier	S	12 y	4.2	Right (III)
	8	Maltese	F	5 y	4.0	Right (III)
Control	9	Maltese	C	9 y	4.8	Left (III)
	10	Shih Tzu	C	3 y	5.5	Left (III)
	11	Maltese	C	3 y	4.2	Right (III)
	12	Pomeranian	F	2 y	6.6	Left (III)

*F: Female, M: Male, C: Castrated, S: Spayed

**m: month, y: year

Table 2. Grading System for Medial Patellar Luxation²³

Grade I	The patella is positioned normally but can be luxated with slight manual pressure.
Grade II	Spontaneous luxation occurs; however, it can reduce spontaneously or can be replaced manually
Grade III	The patella is luxated most of the time; however it can be replaced manually
Grade IV	The patella cannot be reduced manually

of the remainder, 4 were neutered and 2 were spayed (Table 1). All the patients were symptomatic and they presented various degrees of unilateral lameness, sometimes difficulty and reluctance to walk, jump and go upstairs. None of the dogs had a previous history of trauma, and all had clinical signs associated with lameness. The animals were excluded from this study if they had any previous orthopedic disorder, including bone fracture, hip dysplasia, hip luxation and cranial cruciate ligament rupture.

Classifying

The clinical signs associated with medial patellar luxation vary with the degree or grade of luxation (Table 2) (23). Palpation of the stifle joint was performed on both standing animals and dogs in lateral recumbency. The patella is best located by first palpating the tibial tuberosity, then continuing in a proximal direction over the patellar ligament. One hand was placed over the patella, and the other hand was used to pick up the tibia and place the joint through range of motion. With the stifle joint in extension, the patella is isolated between the thumb and index fingers and pushed medially and laterally. The limb should be flexed and extended during examination, and the stability of the patella should be tested in all positions of the limb to determine the degree of luxation.

Surgical technique

Anesthetic induction was used with medetomidine (Domitor[®], Pfizer, USA, 4 µg/kg), tiletamine and zolazepam (Zoletil50[®], Virbac, France, 0.5 mg/kg), and tramadol (Doranjin[®], Samsungpharm, Korea, 0.3 mg/kg) intravenously. And

the anesthesia was maintained with isoflurane (Ifran[®], Hanapharm, Korea).

The patient was placed in dorsal recumbency. The surgical procedure was selected for the treatment of patellar luxation depends on the degree of luxation, the animal's age and physical evaluation of the patella both preoperative and intraoperative. The approach of the stifle joint was cranio-lateral incision. Lateral imbrications, antirotational suture and wedge recession sulcoplasty were performed on grade I case. Lateral imbrications, wedge recession and tibial tuberosity transposition using k-wire were accomplished in the remainder of dogs. After transposition of the tibial tuberosity and before joint capsule closure, the limb should be manipulated extensively several times through the whole range of motion with both internal and external rotation. The transposition is considered adequate if the patella cannot be displaced by manipulation of the limb alone.

Postoperative care

All dogs were rested during 14 days after operation without a Robert-Jones bandage. In IMES group, during two weeks hospitalization, the patients had intramuscular stimulation at the cranial part of sartorius muscle twice a day. Cefazolin was administered intravenously after the operation 1 day and enrofloxacin (Baytril[®], Bayer, 10 mg/kg) at a dose of 10 mg/kg was administered SC every 12 hours for 4 days. Povidone iodine topical solution (10%) was used daily for application on the wound, and the skin sutures were removed on 14 days after surgery. The patients were discharged without problem.

Electrical stimulation

Dry needling was performed using stainless steel acupuncture needles (Dongbang acupuncture[®], Dongbang acupuncture Inc, 0.25 mm * 30 mm) and electrical acupuncture stimulator was used for electrostimulation applications (Suzuki[®], pulse generator pg-306, JAPAN). By adopting the asymmetrical biphasic spike pulse, the tissues were prevented about burns and injuries to the needle. The region was cleaned with alcohol swab and the needle was placed in the appropriate depth until the skin reaction was observed. The frequency of stimulation was 30 Hz and the intensity was set at a level sufficient to include muscle contraction. The current was increased from the occurrence of local muscle contractions to the peak of pain endurance for the 15~20 minutes, twice a day for hospitalization 2 weeks. At the end of the treatment, the electrostimulation was discontinued and the needles were removed. After 30~60 minutes, the animals were measured for degree of stifle joint, muscle mass, lameness score and weight bearing.

Measurement of the degree of stifle joint

The quantity of joint flexion motion is measured by using a goniometer. Measuring maximum angles may involve some discomfort. An animal experiencing discomfort is unlikely to use the limb at those angles while ambulating. Therefore measuring the comfortable ROM may be more clinically applicable. Flexion of stifle joint is measured as the angles formed by the long axis of the tibial shaft and the line joining the lateral femoral epicondyle and greater trochanter. To measure the comfortable ROM, the joint is slowly flexed until the first indication of discomfort, such as tensing the muscles, pulling the limb away, or turning the head slightly, is noted. A transparent plastic goniometer was used to measure the joint angles directly on skin.

Muscle mass

A measuring tape with a spring tension device is useful in measuring limb circumference to improve consistent placement of tension on the tape when making measurements. Thigh length was determined by measuring from the tip of the greater trochanter to the distal aspect of the lateral fabella. Measurements of the sizes of the sartorius muscle were measured at the proximal edge of the patella (16).

Lameness score

The dogs were examined during movement, by two veterinarians who were blind to the grade of patellar luxation of the

Table 3. Lameness Evaluation at Walking¹⁶

Score	State of Ambulation
0	Walks normally
1	Slight lameness
2	Obvious weight-bearing lameness
3	Severe weight-bearing
4	Intermittent non-weight-bearing lameness
5	Continuous non-weight-bearing lameness

dogs. The dogs walked three times to evaluate lameness during movement. The lameness was evaluated by the method of Millis *et al.* (16) (Table 3).

Measurement of weight bearing

Four body weight scales (SW-1s, CAS[®]) were used to evaluate each weight bearing of four limbs at the same time. The scales were placed side by side. Dogs' four limbs were on the each scale. When the dogs took front, instantaneous readings were taken from the digital displays of each scale by as assistant on the "hold" command. The dogs were tried three times to step on and off the scales in the same manner. A 10 seconds interval was allowed between each weight bearing trial on the scales. Three recorded value were averaged, rounded off to a whole number.

Statistical analysis

Data are presented as means \pm SD. A Student's t-test was used for statistical analysis. Differences were seemed to be statistically significant when $P < 0.01$. All analyses were performed using the SPSS software (SPSS ver.19.0, IBM, USA).

Results

ROM of stifle joint

Preoperatively, the result of passive ROM was $81.5 \pm 8.03^\circ$ in IMES group. Passive ROMs of the stifle joint in IMES group were identified improvably on 5th day ($94.2 \pm 4.83^\circ$), 10th day ($102.7 \pm 5.73^\circ$), compared with the control group. The ROM in IMES group improved as $110.9 \pm 3.83^\circ$ on 14 days (Fig 1).

Muscle mass

For sartorius muscle atrophy that occurs with MPL, muscle masses to determine the effects of IMES were measured on 5th, 10th and 14th day after reconstruction of MPL. On 5th day after operation, size of the muscle in IMES group increased to $2.4 \pm 0.18\%$, but increased to $1.2 \pm 0.69\%$ in control group. On 10th day, the size of the muscle mass showed an increase of $3.8 \pm 0.33\%$ in IMES group, whereas an increase of $1.87 \pm 0.78\%$ in control group. On 14th day, the size of muscle in IMES group was $4.75 \pm 0.55\%$, and in




Fig 1. Comparison of the degree of stifle joint between IMES group and control group. The graph is shown significant results on 5th and 10th day. * $p < 0.05$, ** $p < 0.01$.




Fig 2. Comparison of the muscle mass between IMES group and control group. The graph is shown significant results on 5th, 10th and 14th day. ** $p < 0.01$.




Fig 4. Comparison of the affected hindlimb weight bearing between IMES group and control group. The graph is shown significant result on 5th and 10th day. * $p < 0.05$, ** $p < 0.01$.




Fig 3. Comparison of the lameness score between IMES group and control group. The graph is shown significant result on 5th day. ** $p < 0.01$.




Fig 5. Comparison of the non-affected hindlimb weight bearing between IMES group and control group. The graph is shown significant results on 5th and 10th day.

control group was $3.69 \pm 0.50\%$ (Fig 2).

Lameness score

Preoperative lameness observed in the IMES group was 3.5 ± 0.53 and in control group was 4.0 ± 0.00 . On 5th day, unusual changes were identified in IMES group. It showed a significant recovery of the lameness score to 2.6 ± 0.51 . On 10th day, the lameness score in control group was seen to 3.3 ± 0.95 , but in IMES group was seen significant recovery to 2.1 ± 0.35 . Two groups were shown close to normal recovery on 14th day (Fig 3).

Weight bearing

In order to compare the weight bearing correlation of IMES group and control group, weights of the each limb were measured in both affected and non-affected hindlimb. In the IMES treatment group of affected hindlimb, the amount of support on 5th day had decreased to $14.4 \pm 7.44\%$, but in the control group the amount of supported had decreased to $40.4 \pm 4.03\%$ in. On 10th day, weight bearing was increased to $54.7 \pm 7.33\%$ in the IMES group, but was increased $42.8 \pm 6.94\%$ in the control group. On 14th day, there was no significant difference between the two groups (Fig 4). In the IMES treatment group of non-affected hindlimb, the amount of support on 5th day had decreased to $12.4 \pm$

11.66% , but in the control group the amount of support had decreased to $2.3 \pm 2.87\%$. On 10th day, weight bearing was increased to $18.2 \pm 26.98\%$ in the IMES group, but $0 \pm 24.08\%$ in the control group. On 14th day, there was no significant difference between the two groups (Fig 5).

Discussion

Physical agents are various forms of energy, which are often called modalities and their use forms the basis of the practice for human and animal health care. Of all the physical stimuli used for therapy, that produced by a needle is the most efficacious as it is particularly capable of releasing contractures (9). In human, IMES has immediate and mid-term efficacy in pain reduction for myofascial trigger points. ROM of cervical and shoulder was increased objectively but in visual analog scale ROM was significantly decreased subjectively (14). Dogs with limited range of motion in their arthritic joints tend to perceive more pain because these joints are functioning near extreme flexion (e.g., with elbow dysplasia) or extension (e.g., with hip dysplasia) (2). In this study, IMES group indicated improvement of ROM comparing with control group. Like other reports, IMES reduced the shortening of sartorius muscle because spasm of muscle was de-

creased by electrical stimulation. In IMES group, ROM in affected limb showed the significant improvement of the up to 10 days compared in control group. After operation, there were swelling and pain around stifle joint. Swelling and pain around stifle joint as well as shortened sartorius muscle were reduced in IMES group compared in control group.

The physical effects of neuromuscular electrostimulation were to increase in muscle strength and muscle mass (4,19). The application of electrostimulation on the site of injury were shown not only the gross improvement on muscle mass, but also the significant improvement of muscular power in the semitendinosus and semimembranosus muscle, 45 days after surgery (15). In this study, there is an increment of muscle mass in IMES group compared in control group at all recorded days. By the increment of the muscle mass, it is supposed IMES could induce the muscle hypertrophy by contracting muscle. If IMES apply on the sartorius muscle preoperatively, it would prevent further injury to joints and other soft tissues. In this study, better results may be obtained. However, use of a tape measure with a spring tension device may be biased by the expectations of observers. The thickness of the skin and subcutaneous tissues might be another factor affecting thigh circumference measurements.

Lameness score is a vital clinical outcome assessment tool (16). German shepherd observed in the left stifle lameness showed the decreased flexion and extension of all joints, and the right hind leg was placed further forward than normal. He had ruptured cranial cruciate ligament (13). Electrostimulation was felt to be able to produce the same excitatory characteristics within the motor nerve and muscle as does exercise (9,26). In previous researches, electrostimulation affected to the muscular re-education. In this study, the lameness score in IMES group indicated significant result on 5th day. This result means IMES group obtained more quickly muscular function than control group. If IMES is applied with exercise, the score may be improved more rapidly.

In this study, the weight bearing was improved significantly. The animals were compared to weight bearing on affected hindlimb in Fig 4. IMES group had a positive effect on weight shifting into affected hindlimb on 5th and 10th day. Two groups showed the decrease of weight bearing on 5th day. It was thought that the decreases were occurred in pain and swelling of stifle joint after operation. However, there was a significant difference in decreasing of weight bearing between two groups. The weight bearing in IMES group ($14.4 \pm 7.44\%$) was reduced much less than in control group ($40.4 \pm 4.03\%$). This difference means IMES group was more effective than control group in functional and rehabilitative aspects. On 10th day, it showed actual acceleration of weight bearing as observed in Fig 4. Because of the reducing of pain and swelling of stifle joint, sartorius muscle was gradually released and the degree of stifle joint was improved. Also, affected weight bearing was thought to be gradually improved. Before performing this study, although IMES was applied, the animals could not contact on the scale until 4th day in preliminary experiment. Not until 5th day, the dogs did not contact on the scale. The weight was measured from 5th day.

In Fig 5, there was no significant result in non-affected

hindlimb weight bearing. However, on 5th day, weight bearing of IMES group ($12.4 \pm 11.66\%$) was more decreased than that of control group ($2.3 \pm 2.87\%$). As affected hindlimb recovered, it was considered weight bearing transfer non-affected to affected hindlimb. On 10th day, weight bearing of control group (0 ± 24.08) more decreased than that of IMES group (18.2 ± 26.98). It was thought that IMES was helpful for the early rehabilitation.

As results, this study showed the improvement of ROM of stifle joint, femoral muscle mass, lameness score, and weight bearing of hind limb in IMES group. Recovery of ROM and muscle strength after operation was more important than ever. As rehabilitative modality, IMES might provide the novel evidence for the effect of electrostimulation on regaining muscular ability after operation. Therefore IMES can be recommended during regaining muscular ability and re-education in other orthopedic disease. Based on the finding of improved ROM, muscle mass, lameness score, and weight bearing in this study, IMES is expected to play an important role in small animal physical therapy and rehabilitation.

Conclusion

This study performed to know rehabilitative effect after reconstruction as application with intramuscular electrostimulation (IMES) on the cranial part of sartorius muscle. In this study, the 12 dogs with operated medial luxation divided into electrostimulation group ($n=8$) and no electrostimulation group ($n=4$) measured affected sided range of motion, muscle mass, lameness score and weight bearing for hospitalization 2 weeks. IMES group showed significant results after operation on 5th day ($p < 0.01$). Except lameness score, IMES group showed significant results on 10th day ($p < 0.05$).

As results, in small sized dog after reconstruction of MPL, IMES on the cranial part of sartorius muscle considered a great help in the initial rehabilitation.

Acknowledgement

This research was supported by the 2014 scientific promotion program funded by Jeju National University.

References

1. Alam MR, Lee HB, Hea SY, Shin SH, Kim JH, Lee JI, Lee CH, Lee KC, Choi IH, Kim NS. A retrospective study of distribution and surgical management of patellar luxation in the dog: 126 cases (2000 to 2005). *J Vet Clin* 2005; 22: 313.
2. Alam MR, Lee HB, Kim MS, Kim NS. Surgical model of osteoarthritis secondary to medial patellar luxation in dogs. *Veterinari Medicina* 2011; 56: 123-130.
3. Bevan JM, Taylor RA. Arthroscopic release of the medial femoropatellar ligament for canine medial patellar luxation. *J Am Anim Hosp Assoc* 2004; 40: 321-330.
4. Bigard AX, Lienhard F, Merino D, Serrurier B, Guezennec CY. Effects of surface electrostimulation on the structure and metabolic properties in monkey skeletal muscle. *Med Sci Sports Exerc* 1993; 25: 355-362.
5. Black J. Electrical stimulation of hard and soft tissue in

- animal models. Clin Plast Surg 1985; 12: 243-257.
6. Fossum TW. Diseases of the Joints. In: Small animal surgery, 4th ed. St. Louis, Missouri: Mosby Elsevier. 2013: 1353-1360.
 7. Gunn CC. Neuropathic pain: a new theory for chronic pain of intrinsic origin. Ann R Coll Physicians Surg Can 1989; 22: 327-330.
 8. Gunn CC, Milbrandt WE, Little AS, Mason KE. Dry needling of muscle motor points for chronic low back pain. A randomized clinical trial with long-term follow-up. Spine 1980; 5: 279-291.
 9. Hainaut K, Duchateau J. Neuromuscular electrical stimulation and voluntary exercise. Sports Medicine 1992; 14: 100-113.
 10. Horne RD. Transplantation of the cranial head of the sartorius muscle for correction of medial patellar luxations. J Am Anim Hosp Assoc 1979; 15: 561-565.
 11. L'Eplattenier H, Montavon P. Patellar luxations in dog and cats: pathogenesis and diagnosis. Compend Contin Educ Pract Vet 2002; 24: 234-241.
 12. L'Eplattenier H, Montavon P. Patellar luxation in dogs and cats: management and prevention. Compend Contin Educ Pract Vet 2002; 24: 292-300.
 13. Leach D, Sumner-Smith G, Dagg AI. Diagnosis of lameness in dogs: a preliminary study. Can Vet J 1997; 18: 58-63.
 14. Lee SH, Chen CC, Lee CS, Lin TC, Chan RC. Effects of needle electrical intramuscular stimulation on shoulder and cervical myofascial pain syndrome and microcirculation. J Chin Med Assoc 2008; 71: 200-206.
 15. Mert T, Daglioglu YK, Gunay I, Gocmen CI. Changes in electrophysiological properties of regenerating rat peripheral nerves after crush injury. Neurosci Lett 2004; 363: 212-217.
 16. Millis DL, Levine D, Taylor RA. Assessing and measuring outcomes. In: Canine rehabilitation & physical therapy. St Louis: WB Saunders. 2004; 12: 211-227.
 17. Mortari AC, Rahal SC, Vulcano LC, da Silva VC, Volpi RS. Use of radiographic measurements in the evaluation of dogs with medial patellar luxation. Can Vet J 2009; 50: 1064-1068.
 18. Nagaoka K. Advanced surgical techniques for the adult dogs with patellar luxation. Proceeding of WSAVA. Bangkok, Thailand. 2003; 609-611.
 19. Nessler JP, Mass DP. Direct-current electrical stimulation of tendon healing in vitro. Clin Orthop Relat Res 1987; 217: 303-312.
 20. Roush JK. Canine patellar luxation. Vet Clin North Am Small Anim Pract 1993; 23: 855-868.
 21. Sandra SL. Electroacupuncture and stimulatory frequencies in analgesia. In: Acupuncture-concepts and physiology, 1st ed. InTech 2011; 5: 69-90.
 22. Schoen AM. Acupuncture for musculoskeletal disorders. Probl Vet Med 1992: 88-97.
 23. Singleton WB. The surgical correction of stifle deformities in the dog. J Small Anim Pract 1969; 10: 59-69.
 24. Slatter D. Stifle Joint. In: Textbook of small animal surgery, 3rd ed. Philadelphia PA: Saunders Elsevier Science. 2003; 2122-2133.
 25. Tobias KM, Johnston SA. Stifle Joint. In: Veterinary surgery small animal, 1st ed. St.Louis Missouri: Saunders Elsevier. 2012; 973-982.
 26. Toma K, Conatser RR, Gilders RM, Hagerman FC. The effects of acupuncture needle stimulation on skeletal muscle activity and performance. J Strength Cond Res 1998; 12: 253-257.
 27. Towle HA, Griffon DJ, Thomas MW, Siegel AM, Dunning D, Johnson A. Pre- and postoperative radiographic an computed tomographic evaluation of dogs with medial patellar luxation. Vet Surg 2005; 34: 265-272.

소형견의 내측 슬개골 탈구 정복술 후 근육 내 전기자극의 재활효과

이신호 · 이주명 · 박현정 · 원상철 · 정종태¹

제주대학교 수의과대학 · 수의과학연구소

요 약 : 개의 내측 슬개골 탈구는 가장 많이 발생하는 관절 질환 중 하나로 정복 수술 후 넙다리빗근에 전기 자극을 실시하여 재활효과를 관찰하기 위하여 본 연구를 수행하였다. 본 연구는 내측 슬개골 탈구가 있는 소형견 12두를 대상으로 슬개골 탈구 정복술 후에 입원 2주 동안 수술한 후지의 앞쪽 넙다리빗근에 근육 내 전기자극 치료를 적용한 군(n=8)과 대조군(n=4)으로 나누어 관절가동범위, 근육 둘레, 파행 및 체중지지 정도를 측정하였다. 근육 내 전기자극을 실시한 군은 대조군에 비하여 관절가동범위, 근육 둘레, 파행점수 및 체중지지 정도 등 네 가지 사항 모두 수술 후 5일에는 유의성을 나타냈고(p < 0.01), 수술 후 10일에는 파행점수를 제외한 관절가동범위, 근육둘레 및 체중지지 정도의 항목에서 유의성을 보였다(p < 0.05). 이상의 결과로 볼 때, 소형견의 내측 슬개골 탈구 정복술을 실시한 후, 넙다리빗근에 전기자극을 실시하면 초기 재활치료에 도움이 많은 것으로 사료된다.

주요어 : 근육, 전기자극, 넙다리빗근, 슬개골, 탈구, 개