

Case Report

Transcranial Direct Middle Meningeal Artery Puncture for the Onyx Embolization of Dural Arteriovenous Fistula Involving the Superior Sagittal Sinus

Jae-Sang Oh, M.D., Seok-Mann Yoon, M.D., Jai-Joon Shim, M.D., Hack-Gun Bae, M.D.

Department of Neurosurgery, Soonchunhyang University Cheonan Hospital, Cheonan, Korea

A 66-year-old woman presented with intermittent paraparesis and generalized tonic-clonic seizure. Cerebral angiography demonstrated dural arteriovenous fistula (AVF) involving superior sagittal sinus (SSS), which was associated with SSS occlusion on the posterior one third. The dural AVF was fed by bilateral middle meningeal arteries (MMAs), superficial temporal arteries (STAs) and occipital arteries with marked retrograde cortical venous reflux. Transfemoral arterial Onyx embolization was performed through right MMA and STA, but it was not successful, which resulted in partial obliteration of dural AVF because of tortuous MMA preventing the microcatheter from reaching the fistula closely enough. Second procedure was performed through left MMA accessed by direct MMA puncture following small decortications of cranium overlying the MMA using diamond drill one week later. Microcatheter could be located far distally to the fistula through 5 F sheath placed into the MMA and complete obliteration of dural AVF was achieved using 3.9 cc of Onyx.

Key Words : Dural arteriovenous fistula · Superior sagittal sinus · Transcranial · Middle meningeal artery · Onyx embolization.

INTRODUCTION

Dural arteriovenous fistulas (dAVFs) of the superior sagittal sinus (SSS) account for 8–11% of intracranial dural fistulas^{1,4)}. Treatment is often challenging because of multiple and bilateral arterial feeders, most commonly the middle meningeal arteries (MMAs). Transarterial embolization and surgical coagulation of draining vein are preferred treatment options for SSS fistulas, especially in the anterior third of SSS. However, due to the penetration capability of Onyx, transarterial Onyx embolization can result in complete obliteration of dAVF in post-Onyx era. It is important to locate the tip of microcatheter near the fistula as close as possible to obliterate fistula successfully with Onyx. However, superselection of MMA feeders is not always possible when proximal part of MMA is tortuous.

Recently, a combination of surgical and endovascular techniques, including direct-puncture transvenous embolization during surgery or transvenous embolization alone, has been described. And we introduced a case of DAVF within the wall of SSS associated with cortical venous reflux, which was embolized with direct puncture technique after failed initial conventional

transarterial endovascular therapy.

CASE REPORT

Case presentation

A 66-year-old woman was transferred to our hospital for the evaluation of intermittent paraparesis. She had a history of head injury, which was acute subdural hematoma managed conservatively 3 years ago in our hospital. She has been doing well without sequelae even though she complained of intermittent headache for 2 years.

On admission, she complained of gait ataxia because of intermittent paraparesis, which was more severe in the left lower extremity, and left hand tremor for 2 weeks. However, her weakness was not definite on neurologic examination. Whole spine magnetic resonance image (MRI) demonstrated no abnormal findings, but brain MRI revealed multiple signal void lesions on both fronto-parietal convexity suggesting vascular lesion (Fig. 1), which was not visible on previous MRI performed 3 years ago. Before the planned cerebral angiography was done, sudden generalized tonic-clonic seizure was developed.

• Received : February 18, 2014 • Revised : March 13, 2014 • Accepted : March 19, 2014

• Address for reprints : Seok-Mann Yoon, M.D.

Department of Neurosurgery, Soonchunhyang University Cheonan Hospital, 31 Suncheonhyang 6-gil, Dongnam-gu, Cheonan 330-930, Korea
Tel : +82-41-570-3649, Fax : +82-41-572-9297, E-mail : smyoon@schmc.ac.kr

• This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Angiography depicted a dAVFs on the SSS fed by both MMAs, superficial temporal arteries (STAs) and occipital arteries (OAs) and drained into the occluded SSS retrograde fashion, then marked reflux to bilateral superficial cortical veins. There was no feeder from both internal carotid arteries. Because the SSS was occluded at the posterior one third, most cerebral venous outflow was

drained into deep venous system through internal cerebral vein and straight sinus to transverse sigmoid sinus (Fig. 2). These dAVFs were classified as type IIa+IIb according to Cognard et al.¹⁾

Endovascular embolization

1st embolization

Under the general anesthesia, bilateral femoral arterial punctures were done. Through right femoral artery, a 6 F Envoy (Cordis, Miami, FL, USA) guiding catheter was advanced to the right external carotid artery (ECA). And a 5 F diagnostic catheter was placed to the left common carotid artery to evaluate the degree of dAVFs obliteration during the procedures. Initially, Echelon-10 (ev3 Neurovascular, Irvine, CA, USA) microcatheter was navigated through the 6 F guiding catheter, but it was very difficult to navigate the tortuous turning point beyond the foramen spinosum level (Fig. 3A). Thus, the microcatheter was changed to Marathon (ev3 Neurovascular, Irvine, CA, USA) catheter, however, it was impossible to get access the fistula portion. Thus Onyx in-




Fig. 1. Brain magnetic resonance image showing multiple microbleeding (arrowheads) along the sulci of cerebral convexities on gradient echo image (A) and multiple vascular signals (arrows) on both centrum semiovale on gadolinium enhanced T1-weighted image (B).




Fig. 2. Angiography depicting dural arteriovenous fistulas on superior sagittal sinus (SSS) fed by both middle meningeal arteries (arrowheads), superficial temporal arteries (arrows) and occipital arteries (A) and draining into the occluded SSS retrograde fashion, then marked reflux to bilateral superficial cortical veins (B). There was no feeder from both internal carotid arteries and most cerebral venous outflow was drained into deep venous system through internal cerebral vein and straight sinus to transverse sigmoid sinus (C).




Fig. 3. Radiologic images obtained during first embolization session. Lateral view of roadmap image showing the tortuous turning point (arrow) beyond the foramen spinosum level (A). Postoperative skull radiographs showing the Onyx cast after incomplete embolization (B). Postoperative angiogram showing the incomplete obliteration of dAVFs (C). dAVFs : dural arteriovenous fistulas.

jection was done from the proximal frontal branch of MMA, but resulted in the incomplete obliteration.

So, we performed Onyx embolization from the petrosal branch of MMA and the parietal branch of STA. Total of 4 cc Onyx was




Fig. 4. Roadmap image showing the drilling point of MMA (A). Intraoperative photograph demonstrating decortication of temporal bone using diamond drill and dilated MMA after verapamil injection (B). MMA : middle meningeal artery.




Fig. 5. Photograph showing the placement of 5 F micropuncture sheath in the MMA and fixation with black silk tie to prevent accidental removal during the procedures (A). Microcatheter angiogram showing the dAVFs clearly (B). MMA : middle meningeal artery, dAVFs : dural arteriovenous fistulas.



Fig. 6. Postoperative angiograms showing the complete obliteration of dAVFs without cortical venous reflux (A and B). Postoperative skull radiographs showing the Onyx cast after complete obliteration of dAVFs (C and D). dAVFs : dural arteriovenous fistulas.

injected during the first session through right ECA branches. However, the infiltration of Onyx into the fistula was not successful from these vessels (Fig. 3B, C).

2nd embolization

Even though her paraparesis was improved a lot after the first incomplete embolization of dural AVF, 2nd procedure was inevitable to prevent fatal hemorrhage, or seizure.

2nd embolization was composed of direct MMA puncture through the small decortication on the left frontal bone overlying the MMA beyond the turning point. After one week, left transcranial arterial embolization was performed under the general anesthesia in the angiosuite. She was placed on supine position and the head was rotated 60 degree to the opposite side. A 5 F diagnostic catheter was placed to left ECA with continuous heparinized saline irrigation. Under the roadmap guidance, exact location of MMA to be punctured was marked with drill tip (Fig. 4A). After a small semilunar shaped scalp incision was made, decortication of temporal bone using diamond drill was performed on the distal part of turning point of MMA (Fig. 4B). A 10 mg of verapamil was infused through the catheter to dilate the MMA because it was not large enough to puncture successfully.

Direct puncture of MMA using 20 G angioneedle was done without difficulty. Hairwire was inserted into the MMA and 5 F micropuncture sheath was placed over the wire. This sheath was fixed with black silk tie to prevent accidental removal during the procedures (Fig. 5). Through the sheath Echelon-10 (ev3 Neurovascular, Irvine, CA, USA) microcatheter was introduced and advanced to the fistula portion near the SSS. Microangiogram demonstrated dAVF clearly. Onyx injection (total 3.9 cc) from this single artery resulted in complete obliteration of AVF without remnant (Fig. 6). And on post operative angiogram, AVF from both STAs and OAs was also completely obliteration.

Postoperatively, she recovered well and her paraparesis was improved to normal. Diffusion weighted image showed no abnormal high signal lesion on brain parenchyma. Computerized tomography (CT) scan showed Onyx cast within the fistula and arterialized cortical veins.

DISCUSSION

DAVFs involving the SSS are rare but have aggressive clinical

symptom and intracranial hemorrhage due to the venous hypertension. The resultant encephalopathy can be clinically demonstrated on dementia, gait ataxia, seizure, myelopathy, cerebral edema, ischemia, subarachnoid hemorrhage, or any combined of several these signs and symptom¹²). If the cortical venous reflux persists, annual risk of hemorrhage or non hemorrhagic neurological deficits is known to be 15% along with an annual mortality rate of 10.4%. Cognard et al.¹¹ noted intracranial hemorrhage in 10% of patients with type II, 40% with type III and 65% with type IV DAVFs.

Recently, there are many multimodality treatment modalities including compression, transvenous, transarterial embolization, surgery, and radiosurgery^{3,6}). In addition, especially on DAVFs involving the SSS, a combination of surgical and endovascular techniques, including direct-puncture transvenous embolization during surgery or transvenous alone has been described^{4,8,10}). And, stent placement for occluded sinus is an additional treatment option for DAVFs involving the SSS¹³).

In this case, we considered many endovascular treatment options including transvenous coil embolization, SSS stenting, and transarterial Onyx embolization^{5,7,15}). Of these, transvenous coil embolization required surgical exposure of SSS, which had a disadvantage of excessive bleeding. Although SSS stenting could convert aggressive type to benign type, it was considered difficult because the length of occlusion was long¹³).

Therefore, we decide to try transarterial embolization with Onyx. Onyx has an advantage of nonadhesive nature and penetration capability over n-butyl-2-cyanoacrylate. There have been many successful reports on transarterial Onyx embolization of dAVFs^{2,8}).

We tried transarterial Onyx embolization using MMA, however, it was not successful because of proximal tortuosity of MMA. If we can get access to the fistula through MMA, there is a high likelihood of complete obliteration of dAVF with Onyx. To overcome the proximal tortuosity of MMA, we performed direct MMA puncture.

Transcranial combined approach using direct puncture technique is sometimes very useful when a traditional transfemoral transarterial approach was failed. Many neurosurgeons has successfully treated for dAVFs using transcranial approach for venous embolization^{5,11}). We treated this case using transcranial approach for arterial embolization. Koh et al.⁹) reported transcranial transarterial approach with craniectomy for occlusion of dAVFs combined with SSS. In our case we used diamond drill to decoricate the temporal bone to prevent accidental damage to MMA^{5,11,14}). Decorication is more beneficial in cosmetic aspect and more safe for preserving the MMA, instead of burrhole or craniectomy. Then verapamil injection through the catheter can dilate MMA and prevent the vasospasm.

CONCLUSION

Transcranial combines approach using direct puncture tech-

nique may be a alternative way to occluded DAVFs with a patent sinus, or when usual transfemoral transarterial approach have failed.

References

- Cognard C, Gobin YP, Pierot L, Bailly AL, Houdart E, Casasco A, et al. : Cerebral dural arteriovenous fistulas : clinical and angiographic correlation with a revised classification of venous drainage. *Radiology* **194** : 671-680, 1995
- Fukai J, Terada T, Kuwata T, Hyotani G, Raimura M, Nakagawa M, et al. : Transarterial intravenous coil embolization of dural arteriovenous fistula involving the superior sagittal sinus. *Surg Neurol* **55** : 353-358, 2001
- Gross BA, Ropper AE, Popp AJ, Du R : Stereotactic radiosurgery for cerebral dural arteriovenous fistulas. *Neurosurg Focus* **32** : E18, 2012
- Halbach VV, Higashida RT, Hieshima GB, Rosenblum M, Cahan L : Treatment of dural arteriovenous malformations involving the superior sagittal sinus. *AJNR Am J Neuroradiol* **9** : 337-343, 1998
- Houdart E, Saint-Maurice JP, Chapot R, Ditchfield A, Blanquet A, Lot G, et al. : Transcranial approach for venous embolization of dural arteriovenous fistulas. *J Neurosurg* **97** : 280-286, 2002
- Ihn YK, Kim MJ, Shin YS, Kim BS : Dural arteriovenous fistula involving an isolated sinus treated using transarterial onyx embolization. *J Korean Neurosurg Soc* **52** : 480-483, 2012
- Kakarla UK, Deshmukh VR, Zabramski JM, Albuquerque FC, McDougall CG, Spetzler RF : Surgical treatment of high-risk intracranial dural arteriovenous fistulae : clinical outcomes and avoidance of complications. *Neurosurgery* **61** : 447-457; discussion 457-459, 2007
- Kiyosue H, Hori Y, Okahara M, Tanoue S, Sagara Y, Matsumoto S, et al. : Treatment of intracranial dural arteriovenous fistulas : current strategies based on location and hemodynamics, and alternative techniques of transcatheter embolization. *Radiographics* **24** : 1637-1653, 2004
- Koh JS, Ryu CW, Bang JS, Lee SW : Transcranial Approach for Arterial Embolization of Dural Arteriovenous Fistula Within the Wall of the Superior Sagittal Sinus. A Case Report. *Neurointervention* **2** : 117-121, 2007
- Kurl S, Saari T, Vanninen R, Hernesniemi J : Dural arteriovenous fistulas of superior sagittal sinus : case report and review of literature. *Surg Neurol* **45** : 250-255, 1996
- Luo CB, Chang FC, Wu HM, Chung WY : Transcranial embolization of a transverse-sigmoid sinus dural arteriovenous fistula carried out through a decompressive craniectomy. *Acta Neurochir (Wien)* **149** : 197-200; discussion 200, 2007
- Oh JT, Chung SY, Lanzino G, Park KS, Kim SM, Park MS, et al. : Intracranial dural arteriovenous fistulas : clinical characteristics and management based on location and hemodynamics. *J Cerebrovasc Endovasc Neurosurg* **14** : 192-202, 2012
- Ohara N, Toyota S, Kobayashi M, Wakayama A : Superior sagittal sinus dural arteriovenous fistulas treated by stent placement for an occluded sinus and transarterial embolization. A case report. *Interv Neuroradiol* **18** : 333-340, 2012
- Pierot L, Visot A, Boulin A, Dupuy M : Combined neurosurgical and neuroradiological treatment of a complex superior sagittal sinus dural fistula : technical note. *Neurosurgery* **42** : 194-197, 1998
- Toyota S, Fujimoto Y, Wakayama A, Yoshimine T : Complete Cure of Superior Sagittal Sinus Dural Arteriovenous Fistulas by Transvenous Embolization through the Thrombosed Sinus in a Single Therapeutic Session. A Case Report. *Interv Neuroradiol* **14** : 319-324, 2008