

Single-dose Intramuscular Injection Toxicology of Danggui Pharmacopuncture (DGP) in Sprague-Dawley Rats

SeungHo Sun¹, JongJin Jeong¹, Sunju Park², KwangHo Lee³, JunSang Yu⁴, Hyung-Sik Seo⁵, KiRok Kwon^{6*}

¹Department of Internal Medicine, College of Korean Medicine, Sangji University, Wonju, Korea

²Department of Preventive Medicine, College of Korean Medicine, Daejeon University, Daejeon, Korea

³Department of Acupuncture & Moxibustion Medicine, College of Korean Medicine, Sangji University, Wonju, Korea

⁴Department of Sasang Constitutional Medicine, College of Korean Medicine, Sangji University, Wonju, Korea

⁵Department of Ophthalmology, Otolaryngology, and Dermatology, Korean Medicine Hospital, Pusan National University, Yangsan, Korea

⁶Research Center of the Korean Pharmacopuncture Institute, Seoul, Korea

Key Words

acupuncture, *Angelica gigas nakai*, *Angelica gigantis radix*, herbal medicine, pharmacopuncture, single-dose toxicity test

Abstract

Objectives: The purpose of the study is to assess both the approximate lethal dose and the single dose intramuscular injection toxicity of Danggui (*Angelica gigantis radix*) pharmacopuncture (DGP) in Sprague-Dawley (SD) rats.

Methods: The experiments were conducted at the good laboratory practice (GLP) laboratory, Biototech Co., which is a laboratory approved by the ministry of food and drug safety (MFDS). The study was performed according to the GLP regulation and the toxicity test guidelines of the MFDS (2009) after approval of the institutional animal care and use committee of Biototech. Single doses of DGP were injected intramuscularly into the rats in three test groups of 6 week old SD rats (5 male and 5 female rats per groups) in the amounts of 0.1, 0.5, and 1.0 mL/animal for groups 2, 3, and 4, respectively, and normal saline solution in the amount of 1.0 mL/animal was injected intramuscularly into the

rats (5 male and 5 female rats) in the control group. Observations of the general symptoms and weight measurements were performed during the 14 day observation period after the injection. Hematologic and serum biochemical examination, necropsy, and a local tolerance test at the injection site were done after the observation period.

Results: No death was observed in three test groups (0.1, 0.5 and 1.0 mL/animal group). In addition, the injection of DGP had no effect on general symptoms, weights, hematologic and serum biochemical examination, and necropsy. The results from the local tolerance tests at injection site showed no treatment related effects in the SD rats.

Conclusion: The results of single dose intramuscular injection of DGP suggest that the approximate lethal dose is above 1.0 mL/animal for both male and female SD rats and that intramuscular injection of DGP may be safe.

1. Introduction

Pharmacopuncture is a unique acupuncture treatment of traditional Korean medicine (TKM) which combines acupuncture based on the meridian theory

Received: Nov 17, 2014 Reviewed: Dec 02, 2014 Accepted: Dec 15, 2014

© This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

© This paper meets the requirements of KS X ISO 9706, ISO 9706-1994 and ANSI/NISO Z39.48-1992 (Permanence of Paper).

*Corresponding Author

Kirok Kwon. Korean Pharmacopuncture Institute, 4F, Association of Korean Oriental Medicine B/D 26-27, Gayang-dong, Gangseo-gu, Seoul 157-200, Korea.
Tel: +82-33-744-9304 Fax: +82-33-744-9305
E-mail: drkwon5031@daum.net

and herbal medicine based on Qi and flavor theory. Specific amounts of herbal extracts are injected at acupoints according to the differential diagnosis of a TKM doctor [1]. Danggui (DG) is commonly refers to *Angelica* plants that belong to the family of Apiaceae. The types of DG are *Angelica sinensis* and *Chinese angelica* in China and *Angelica gigas* in Korea. *Angelica gigas radix* (AGR) is a root of *Angelica gigas*, which is called Cham Dang Gui (Korean Danggui), and is used as medicinal herb in Korea. The effects of AGR have been reported to tonification and activation of blood (補血活血), regulation of menstruation (調經), relief of pain (止痛), moistening of the intestines (潤腸), and relaxation of the bowels (通便) [2, 3]. DG has been named 'female ginseng' and has been widely used to treat blood deficiency and various diseases in women [2, 3].

Danggui pharmacopuncture (DGP), namely AGR pharmacopuncture, is an acupuncture method in which an AGR extract solution is injected at an acupoint to treat disease. Experimental studies have reportedly shown that DGP has analgesic, antioxidant and potential chemopreventive effects [4-6] and that it has a beneficial effect on immune function, recovery anemia due to blood loss, liver protection against acute drug induced hepatitis and suppression of neuron cell regeneration in ischemic stroke or neural diseases [7-11]. Although research on the efficacy of DGP is ongoing, safety studies of DGP have not been reported. The toxicity of DGP administered by using intramuscular injection, which is the main treatment method of pharmacopuncture, needs to be determined. The purpose of this study is to assess the single dose intramuscular injection toxicity of DGP in Sprague-Dawley (SD) rats and to evaluate its approximate lethal dose.

2. Materials and Methods

One kg of ground *Angelicae gigantis radix* was mixed with 5 L of 70% ethanol (EtOH) for 6 hours at 80 rpm and 60 °C. The process was repeated once. The extract was produced after filtering. Five kg of the extract (extract + solvent) was concentrated, and 370 g of *Angelicae gigantis radix* extract, including decursin and decursinol angelate, were obtained. Appropriate amounts of decursin extract, emulsifying agent, and the water for injection (WFI, Fine FA, Korea) were added to a pyrex glass to produce a water solution after mixing with a high speed agitator. Sodium hydroxide (NaOH) was used to obtain a pH of 7.3 for the water solution. An emulsifying solution was transferred with N₂ gas from the mixing tank to the filling tank through a filter (1 μm, Satorius). This was fired in a sterilized bottle in an autoclave at 121 °C for 30 minutes and was then kept under refrigeration at 4 °C.

SD rats (Orientbio Inc., Korea), which are widely used for safety tests, were used in this study. Basic visual examinations were done when the animals were received, after which their weights were measured with an electronic scale (CP3202S, Sartorius, Germany). The animals were then moved to animal chambers. The general symptoms were observed daily during the 7-day stabilization period. The body weights, general basic symptoms, and weight changes were measured on the last day of stabilization in

order to confirm the health status of the rats. At the time of intramuscular (IM) injection, the weight range of the 6 6-week-old male rats (n = 20) was 191.5 — 217.2 g and that of the 6-week-old female rats (n = 20) was 145.5 — 173.4 g.

The animals were housed in a room maintained at a temperature of 21.0 — 23.2 °C and a relative humidity of 40.9% — 59.4%, with artificial lighting from 07:00 to 19:00 (150 — 300 Lux) and a ventilation rate 10 — 15 times/hour. During the stabilization period, the animals (three during the stabilization period and one during the observation period) were housed in stainless steel wire mesh cages 260 mm (w) × 350 mm (d) × 210 mm (h) during the single dose intramuscular injection toxicity test. Animals were allowed free access to irradiation sterilized pellet feed (Teklad Certified Irradiated Global 18% Protein Rodent Diet, 2918C, Harlan Laboratories, Inc., USA). This study was conducted with the approval of the Institutional Animal Care and Use Committee of Biototech Co. (Approval No: 110156) and was performed in compliance with the good laboratory practice (GLP) regulation and the toxicity test guidelines of the ministry of food and drug safety (MFDS). Grouping was done on the last day of stabilization. Twenty animals of each sex with the mean weight were selected and randomly placed into 4 groups with 5 male rats and 5 female

Table 1 Grouping of the animals

Group	Injection dose (mL/animal)	Number of animals	
		Male	female
G1: Control group	1.0 (Saline)	5	5
G2: Low-dose group	0.1 (DGP)	5	5
G3: Mid-dose group	0.5 (DGP)	5	5
G4: High-dose group	1.0 (DGP)	5	5

DGP, Danggui pharmacopuncture.

rats for each group so as to have the same average weight per group (Table 1).

DGP was injected intramuscularly by using a method applicable to clinics. Dosages for the control and the high-dose groups were 1.0 mL/animal of saline and of DGP, respectively, and those for the low dose and the mid dose groups were 0.1 and 0.5 mL/animal, respectively. A single dose of DGP was injected into the left thigh muscle of the SD rats in the low and the mid dose groups, and 0.5 mL of DGP was injected into each thigh muscle of the SD rats in the control and the high dose groups. The injections were done using disposable syringes (1 mL, 26 gauge). The expected clinical dose of DGP is 1.0 mL/time. The results of a pilot test (Biototech Study No.: B12875P) showed no death for a single dose intramuscular injection of 1.0 mL/animal in both male and female rats. As a result, 1.0 mL/animal was selected as the high dose, and 0.5 mL and 0.1 mL were set as the mid and the low doses. Normal saline (Choongwae Pharma Corp., Korea) was injected for control group. At the day of injection (day 0), clinical signs of toxicity, such as the onset time, the recovery time, etc.,

and any deaths were noted at 30 minutes, 1 hour, and 2 hours after the first injection, and at 30 minutes, 1, 2, 4 and 6 hours after the second injection. Clinical signs were observed daily for 14 day after injection. The weights of the SD rats were measured on the day of the injection (before injection), and on the third, seventh, and fourteenth (necropsy day) days after injection. Necropsy and blood collection were performed after fasting for more than 18 hours. Approximately 1 mL of blood was taken from the abdominal aorta by using a syringe needle under isoflurane anesthesia. The blood samples of all rats were collected in a bottle containing ethylenediaminetetraacetic acid (EDTA) and were analyzed using a blood counting analyzer (ADVIA 120, Siemens, Germany) to measure the red blood cell count (RBC), hemoglobin concentration (HGB), hematocrit (HCT), mean corpuscular cell volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular cell hemoglobin concentration (MCHC), platelet count (PLT), white blood cell count (WBC), WBC differential count (neutrophils, lymphocytes, monocytes, eosinophils, and basophils), and reticulocyte count. About 2.0 mL of the blood taken during necropsy was placed into a tube containing 3.2% sodium citrate, and plasma was obtained by using centrifugation (3000 rpm) for 10 minutes. The prothrombin time (PT) and the active partial thromboplastin time (APTT) were measured using an automated coagulation analyzer (Coapresta 2000, Sekisui, Japan).

Blood samples taken from the abdominal aorta were centrifuged at 3,000 rpm for 10 minutes and analyzed using an automatic analyzer (7180, Hitachi, Japan) and an electrolyte analyzer (AVL9181, Roche, Germany). Serum biochemistry parameters, including blood urea nitrogen (BUN), creatinine (Crea), aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP), gamma glutamyl transpeptidase (GGT), total protein (TP), albumin (Alb) and albumin/globulin ratio (A/G ratio), total bilirubin (T-Bili), total cholesterol (T-Chol), triglycerides (TGs), phosphorus (P), glucose (Glu), calcium (Ca), sodium (Na), potassium (K) and chloride (Cl), were measured.

After the observation period, necropsy was conducted after all surviving animals had been euthanized by abdominal aorta bloodletting with carbon dioxide (CO₂) gas anesthesia. Detailed visual inspections of the organs and tissues of all surviving animals were done. A local tolerance test was conducted on the tissue at the injection site for all animals. The tissue from the injection site was fixed with 10% neutral buffered formalin solution. The fixed tissues were routinely processed, embedded in paraffin, and sectioned. The sections were stained with hematoxylin and eosin stain for microscopic examination.

Weight, hematology and serum biochemistry data were analyzed by using statistical analysis system (version 9.3, SAS Institute Inc., USA). Data are presented as mean \pm standard deviation. The test for equality variance was the Bartlett test ($P < 0.05$). If the variance of the data was assumed to be normally distributed, the data were analyzed by using the one way analysis of variance (ANOVA). If a significant difference was observed between the groups, the data were tested by using Dunnett's *t*-test for a post hoc analysis ($P < 0.05$, $P < 0.01$). If the variance was not homo-

geneous, the data were analyzed using the Kruskal-Wallis test ($P < 0.05$).

3. Results

No deaths of both male and female rats were observed in any of the four groups during the observation period (Table 2). In addition, no clinical signs were observed in either male or female SD rats in the control group or in the three test groups during the observation period (Tables 3, 4). Furthermore, no statistically significant changes in weight were observed for either male or female SD rats in the test groups compared with those in the control group during the observation period (Figs. 1, 2).

No significant changes in any of the hematological parameters, such as RBC, HGB, HCT, MCV, MCH, MCHC, PLT, WBC, WBC differential count (neutrophils, lymphocytes, monocytes, eosinophils, and basophils), and reticulocyte count, were observed in any of the groups. In addition, no meaningful changes in serum biochemistry, such as BUN, Crea, ALT, ALP, GGT, TP, Alb, A/G ratio, T-Bili, T-Chol, TGs, P, Glu, Ca, Na, K, and Cl, were found in any of the groups. Significant changes of parameters among the test groups were minimal. No dose dependent toxicological signification was observed in any of the groups, and no macroscopic abnormalities were observed at necropsy in either both male and female SD rats in the control and the test groups during the observation period (Table 5).

Local tolerance tests at the injection sites of all animals showed no treatment-related effects. Moderate linear infiltration of inflammatory cells at the injection site was found in a male rat from the 1.0 mL/animal dose test group. Minimal infiltration was observed in a female rat from both the 0.5 mL and 1.0 mL/animal dose test groups. All sites of infiltration were confined to biceps femoris region (Table 6).

4. Discussion

DG contains essential oil and water-soluble materials. The main constituents of essential oil are ligustilide (45%), flavonoids, carvacrol, and *n*-butylidene phthalide. The water soluble materials are composed of mainly ferulic acid (0.094%) with nicotinic acid, uracil, adenine, and polysaccharide. In terms of pharmacological aspects, organic acids and phthalide are active components of DG. Phthalide is a lactone that has a cyclic structure with a ketone body, and various forms of phthalides exist in DG, such as *n*-butylidene phthalide, ligustilide, and *n*-butylphthalide. Phthalide is a volatile oil, and the odor of DG results from ligustilide and *n*-butylidene phthalide. In addition, coumarin derivatives such as oxypeucedanin, osthole, imperatorin, psoralen, and bergapten are included [2-3, 12].

The characteristics of DGP are as follows: As the medicinal effects of DGP depend on the essential oils, such as ligustilide and the water soluble constituent, the extracted essential oil components of DGP are micro pneumatized to be water soluble to penetrate easily throughout the

Table 2 Summary of mortality

Sex	Group / Dose (mL/animal)	No. of animals	Days after dosing															Mortality
			0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	
Male	G1 1.0 (Saline)	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0/5
	G2 0.1 (DGP)	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0/5
	G3 0.5 (DGP)	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0/5
	G4 1.0 (DGP)	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0/5
Fe- male	G1 1.0 (DGP)	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0/5
	G2 0.1 (DGP)	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0/5
	G3 0.5 (DGP)	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0/5
	G4 1.0 (DGP)	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0/5

DGP, Dangui pharmacopuncture.

Table 3 Summary of clinical signs by hour after dosing

Sex	Group / Dose (mL/animal)	No. of animals	Clinical sign	Hours (Day 0) after dosing				
				0.5	1	2	4	6
Male	G1 1.0 (Saline)	5	NOA	5	5	5	5	5
	G2 0.1 (DGP)	5	NOA	5	5	5	5	5
	G3 0.5 (DGP)	5	NOA	5	5	5	5	5
	G4 1.0 (DGP)	5	NOA	5	5	5	5	5
Fe- male	G1 1.0 (Saline)	5	NOA	5	5	5	5	5
	G2 0.1 (DGP)	5	NOA	5	5	5	5	5
	G3 0.5 (DGP)	5	NOA	5	5	5	5	5
	G4 1.0 (DGP)	5	NOA	5	5	5	5	5

NOA, no observable abnormality; DGP, Dangui pharmacopuncture.

Table 4 Summary of clinical signs by day after dosing

Sex	Group / Dose (mL/animal)	No. of animals	Clinical sign	Days after dosing													
				1	2	3	4	5	6	7	8	9	10	11	12	13	14
Male	G1 1.0 (Sa-line)	5	NOA	5	5	5	5	5	5	5	5	5	5	5	5	5	
	G2 0.1 (DGP)	5	NOA	5	5	5	5	5	5	5	5	5	5	5	5	5	
	G3 0.5 (DGP)	5	NOA	5	5	5	5	5	5	5	5	5	5	5	5	5	
	G4 1.0 (DGP)	5	NOA	5	5	5	5	5	5	5	5	5	5	5	5	5	
Fe-male	G1 1.0 (Sa-line)	5	NOA	5	5	5	5	5	5	5	5	5	5	5	5	5	
	G2 0.1 (DGP)	5	NOA	5	5	5	5	5	5	5	5	5	5	5	5	5	
	G3 0.5 (DGP)	5	NOA	5	5	5	5	5	5	5	5	5	5	5	5	5	
	G4 1.0 (DGP)	5	NOA	5	5	5	5	5	5	5	5	5	5	5	5	5	

NOA, no observable abnormality; DGP, Danggui pharmacopuncture.




Figure 1 Body weights in male SD rats.

body. Therefore, this enables DGP to contain as much of both the water soluble materials and the essential oils as possible to maximize its efficacy.

DG is a representative medicinal herb for treating blood deficiency. The main effects of DG are blood related actions, such as inhibition of platelet aggregation and the antithrombotic effect. It also has hematogenetic effects that stimulate hemopoietic stem cells and colony form-

ing unit-granulocyte monocyte (CFU-GM) [3, 8]. DG also contains components that both stimulate and inhibit the smooth muscles of the uterus through bi-modal functions that mediate uterine diseases, such as dysmenorrhea, irregular menstruation, oxytocic uterine contractions, and uterine hemorrhage [3, 13]. The water extract of DG and ferulic acid release myocardial ischemia, and ligustilide and n-butylidenephthalide of DG vasodilate and lowers




Figure 2 Body weights in female SD rats.

Table 5 Summary of necropsy findings

Sex	Male				Female				
	Group	G1	G2	G3	G4	G1	G2	G3	G4
Dose of DGP (mL/animal)		0	0.1	0.5	1.0	0	0.1	0.5	1.0
No. of animals		5	5	5	5	5	5	5	5
Unremarkable findings		5	5	5	5	5	5	5	5
No. examined		5	5	5	5	5	5	5	5

DGP, Danguigui pharmacopuncture. External surfaces and all organs in the body cavity were unremarkable.

Table 6 Summary of histopathological findings.

Sex	Male				
	Group	G1	G2	G3	G4
Organ / Findings	Dose of DGP (mL/animal)	0	0.1	0.5	1.0
	No. of animals	5	5	5	5
Injection site	-Infiltration, inflammatory cells	± 0	0	0	1
	No. examined	5	5	5	5

Sex	Female				
	Group	G1	G2	G3	G4
Organ / Findings	Dose of DGP (mL/animal)	0	0.1	0.5	1.0
	No. of animals	5	5	5	5
Injection site	-Infiltration, inflammatory cells	± 0	0	1	1
	No. examined	5	5	5	5

Grade- ±, minimal. DGP; Danguigui pharmacopuncture.

blood pressure, which may have an effect when used to treat cardiovascular diseases [3, 13]. Based on the effects of DG described above, DGP can be used to treat gynecological diseases, such as dysmenorrhea, as well as cardiovascular diseases and hematologic diseases, such as stroke, arteriosclerosis, and cerebral thrombosis [10, 14-15].

Ligustilide of DG is a potent, toxic metabolite, as well as an active component. Furthermore, caffeic acid is contained not only in DG but also in other various plants. Despite its pharmacological effects, such as anti-carcinogenic, anti-oxidant, immune regulating, and anti-inflammation effects, carcinogenic properties of active metabolites have been reported, so the possibility of toxicity cannot be ruled out [16, 17]. However, no toxic results were observed in our single dose intramuscular injection toxicity tests. Moderate linear infiltration of inflammatory cells at the injection site was found on the histopathologic test, but seemed to have been the result of a needle injury rather than DGP injection. On the basis of our test results, a single dose intramuscular injection of DGP is safe. However, repeated toxicity tests are required to increase confidence in these safety results.

5. Conclusion

The results of single dose intramuscular injection of DGP suggest that the approximate lethal dose is higher than 1.0 mL/animal for both male and female SD rats and that intramuscular injection of DGP may be safe.

Acknowledgment

The authors have no financial interest related to the material of this manuscript.

Conflict of interest

The authors declare that there are no the conflicts of interest.

References

1. Korean Pharmacopuncture Institute. [Pharmacopunctureology: principles and clinical application]. Seoul: Elsevier Korea LLC; 2012. Chapter 1, Definition and history. p. 3-4. Korean.
2. Park YC, Lee JS, Kim MH, Kim DY, Lee SD. [Pharmacological action and toxicity of *Angelica sinensis*]. Korean J Orient Med Prescr. 2011;19(2):93-108. Korean.
3. Han JH, Kim KY. [Korean traditional pharmacology]. Seoul: Euseongdang publishing INC; 2004. Chapter 20, Deficiency-tonifying medicinal. p. 479-83. Korean.
4. Kim YK, Cho KH, Shon YH, Choi HK, Kim SY, Lim JK, et al. [Chemopreventive potential of *Angelicae gigantis radix* aqua-acupuncturesolution]. Yakhak Hoeji. 2000;44(3):283-92. Korean.
5. Park KH, Kang SK. [Studies on the analgesic effect of aqua-acupuncture with angelicae radix extract solution to mice]. KyungHee Univ Orient Med J. 1984;7:261-71. Korean.
6. Ahn JC, Moon JY, Lim JK. [Antioxidant effect of AGR (*Angelicae girantis radix*) aqua-acupuncture solution *in vitro*]. The acupuncture. 1996;13(2):254-62. Korean.
7. Hwang HS, Ahn BC, Park DS. [The effects of *Angelicae gigantis radix* aqua - acupuncture by density on immune response induced by radiation in mice]. The acupuncture. 1994;11(1):113-29. Korean.
8. Lee SJ, Koh HK, Kim CH. [Effects of aqua-acupuncture with saiviae radix and angelicae radix extract solution on the acute anemia induced by loss of blood in rats]. KyungHee Univ Orient Med J. 1991;14:381-36. Korean.
9. Park SH, Kang JH, Jung YS, Kim GW, Koo BS. [The effect of an essential oil fragrance from radix *Angelica sinensis* on differentiation and proliferation of neural stem cells of rat]. J of Oriental Neuropsychiatry. 2009;20(4):63-77. Korean.
10. Song BK, Jeon YC, Kim SA, Shim AN, Seong KM, Lee EJ. [The effect of intravenous injection of the water extract of *Angelica gigas nakai* on gliosis in the middle cerebral artery occlusion rats]. J Pharmacopuncture. 2011;14(3):5-17. Korean.
11. Han SG, Lee BR. [Effects of *Angelica gigas nakai* herbal acupuncture into hyolhae (SP10) of brain ischemic injury induced by intraluminal filament insertion in the rats]. The acupuncture. 2004;21(2):1-20. Korean.
12. Sun Y, Tang J, Gu X, Li D. Water-soluble polysaccharides from *Angelica sinensis* (oliv.) diels: preparation, characterization and bioactivity. Int J Biol Macromol. 2005;36(5):283-9.
13. Kim SJ, Song BK, Lee EJ, Kim HK, Kim JK. [Effects of radix *Angelicae gigantis* and resina ferulae on the relaxation of smooth muscle and expression of iNOS]. J Korean Oriental Med. 2000;20(2):60-7. Korean.
14. Kim YJ, Hwang CW. [An experimental study on brain damage and cardiovascular system effect of *Angelicae gigantis radix* extract]. J Korean Oriental Med. 2000;21(4):37-47. Korean.
15. Kim KS, Song YJ. [Effects of aqua-acupuncture with radix *Angelicae gigantis* and radix astragali on the blood in the liver damage rats with CCL4]. The acupuncture. 1996;13(1):1-10. Korean.
16. Yan R, Ko NL, Li S, Tam YK, Lin G. Pharmacokinetics and metabolism of ligustilide, a major bioactive component in rhizoma chuanxiong, in the rat. Drug Metab Dispos. 2008;36(2):400-8.
17. Barlow S, Schlatter J. Risk assessment of carcinogens in food. Toxicol Appl Pharmacol. 2010;243(2):180-90.