

## Pediatric Issue

# Pediatric Stroke

Goun Jeong, M.D.,<sup>1</sup> Byung Chan Lim, M.D.,<sup>2</sup> Jong-Hee Chae, M.D., Ph.D.<sup>2</sup>

Department of Pediatrics,<sup>1</sup> Cheil General Hospital and Women's Healthcare Center, Dankook University College of Medicine, Cheonan, Korea

Department of Pediatrics,<sup>2</sup> Seoul National University Children's Hospital, Seoul National University College of Medicine, Seoul, Korea

Pediatric stroke is relatively rare but may lead to significant morbidity and mortality. Along with the advance of brain imaging technology and clinical awareness, diagnosis of pediatric stroke is increasing worldwide. Pediatric stroke differs from adults in variable risk factor/etiologies, diverse and nonspecific clinical presentation depending on ages. This review will be discussed pediatric stroke focusing on their clinical presentations, diagnosis and etiologies/risk factors.

**Key Words :** Pediatric · Stroke · Clinical · Risk factors · Etiology · Diagnosis.

## INTRODUCTION

Stroke (focal cerebral injury with an underlying vascular basis) is typically classified into ischemic or hemorrhagic subtypes. Ischemic stroke (IS) is further subdivided into arterial ischemic stroke (AIS) and cerebral sinovenous thrombosis (CSVT). Stroke is increasingly recognized as a serious neurologic disorders of children and awareness about pediatric stroke is increasing. In western countries, the incidence of pediatric AIS is estimated at 2–3/100000 children/year during the first 5 years and 8–13/100000 from 5 to 14 years<sup>8)</sup>. Many cases occur during the perinatal or neonatal period, during which the incidence is approximately 1/4000 live birth<sup>14)</sup>.

Stroke is also commonly encountered disease in the practice of pediatric neurology and among the top 10 causes of death in children, with highest rates in the 1st year of life in USA<sup>2)</sup>. Therefore, the timely appropriate diagnosis and detection of associated risk factors is very important to improve the management and modify the prognosis. However, even when stroke presents acutely, delays in diagnosis are common and appropriate treatments are difficult because pediatric stroke is relatively rare and the etiologies and associated risk factors are remarkably variable depending on ages and different from those of adults. The International Pediatric Stroke Study (IPSS) and other studies of childhood stroke reported various underlying systemic factors associated with pediatric stroke. In previous studies, preexisting risk factors have been reported in 77–79% of childhood stroke and

varied depending on their ages and ethnics<sup>11)</sup>. The associated conditions are known to be congenital heart disease, vasculopathies, infection, metabolic disease, prothrombotic disorders and trauma. However, most of the previous studies have been established in European and western and few reported in Asian countries.

Here, etiology and risk factors including our center experience, diagnosis of pediatric stroke particularly in arterial ischemic stroke will be focused in this review.

## CLINICAL FEATURES AND DIAGNOSIS

Diagnosis of stroke in children is frequently delayed or missed, which reflect a combination of rareness, lack of awareness and complex differential diagnosis. Many studies have been reported the diagnostic delay beyond 24 hours from symptom onset<sup>16)</sup>. A high clinical suspicion, conjunction with a systematic diagnostic approach is required for timely diagnosis. The acute onset of focal neurologic deficit in children is considered to be stroke until otherwise proven.

A couple of studies have reported how strokes present in children (Table 1). AIS usually presents as a focal neurologic deficit. Hemiparesis is most common focal neurologic deficit but may be misdiagnosed as seizures or Todd's paralysis, meningitis/encephalitis, migraine, or demyelination, which are mimic strokes. Additional common focal neurologic deficits are diplopia, dysarthria, vertigo and ataxia. Seizures at the onset of stroke are

• Received : April 21, 2015 • Revised : May 26, 2015 • Accepted : May 29, 2015

• Address for reprints : Jong-Hee Chae, M.D., Ph.D.

Department of Pediatrics, Seoul National University Children's Hospital, Seoul National University College of Medicine, 101 Daehak-ro, Jongno-gu, Seoul 110-744, Korea  
Tel : +82-2-743-3455, Fax : +82-2-2072-3622, E-mail : chaeped1@snu.ac.kr

• This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Table 1.** Clinical presentation of pediatric stroke

	Ischemic		Hemorrhagic	
	Earley et al. <sup>4)</sup>	deVeber et al. <sup>3)</sup>	Earley et al. <sup>4)</sup>	Meyer- Heim and Boltshauser <sup>12)</sup>
Hemiparesis or focal CNS deficits	94%	51%	21%	16%
Change in mental status	28%		88%	52%
Headache	22%		59%	76%
Seizure	16%	48%	29%	28%
Speech Disorder inc. aphasia		17%		8%
Vomiting				48%
Nausea				20%
Somnolence				12%
Visual Impairment				12%
Neck pain				8%
Fever/Prodrome	35–40%		35–40%	

CNS : central nervous system

common both in ischemic and hemorrhagic stroke, compared to adult. They occur in up to 50% of children, which are not restricted to any age group<sup>24)</sup>.

In addition to focal neurologic deficits, non-focal and diffuse neurologic symptoms such as headache, confusion, irritability and behavioral changes are frequent especially in younger children<sup>5,13,23)</sup>.

Clinical presentation can be significantly different based on the child's age. The younger the child, more non-specific for their symptoms. Neonates usually present with seizures and symptoms of encephalopathy (irritability, lethargy, excessive sleeps, or poor feeding) or apnea. Focal neurologic deficits occur in less than 25% of all newborns. Toddlers can also present with nonspecific symptoms such as increased crying, drowsiness, irritability, poor feeding, vomiting and sepsis-like features. With increasing ages, stroke manifests with a clinical presentation similar to that in adults. The other typical presentation in children is fluctuation course of neurologic deficits, which are frequently brief/recurrent and may resolve as quickly as within a few hours. This should alert the clinician to the possibility of an underlying vasculopathy or thromboembolic causes.

There are many other diseases, which may mimic a stroke. Complicated migraine can cause focal neurologic deficits which typically resolve within 24 hours and usually having family history of hemiplegic migraine<sup>23)</sup>. Partial seizures can result in transient postictal paresis (Todd's palsy). Brain tumor as well as intracranial infection such as meningitis, encephalitis, and brain abscess are also considered as a differential diagnosis. Rarely alternating hemiplegia, metabolic disorders like Mitochondrial Encephalopathy, Lactic Acidosis and Stroke like episodes (MELAS) can also cause stroke like symptoms.

## ETIOLOGIES/RISK FACTORS

Etiologies/risk factors in childhood stroke differ from those in adult stroke<sup>11)</sup>. The etiologies/risk factors identified in adult cases are primarily related to arrhythmias and obstructive ath-

erosclerotic arteriopathies; in contrast, these are rarely thought to be related to childhood stroke. The International Pediatric Stroke Study (IPSS) and other studies of childhood stroke reported various underlying systemic factors. Approximately 50–80% children with AIS have at least one identifiable risk factor for stroke and multiple risk factors often present in as many as 25% of children with stroke<sup>3,7,11)</sup>. According to the largest study by the IPSS of 493 children with stroke in 10 countries, risk factor variability can be influenced by geography, ethnicity, age at presentation and the availability of medical resources<sup>11)</sup>. In neonatal stroke, dehydration, infection and congenital heart disease are more commonly associated compared to older children<sup>13)</sup>. Arteriopathies and congenital heart disease (CHD) are commonly associated from 20–50% of AIS in children in developed countries<sup>1,10)</sup>. Studies in Saudi Arabia and Northern India reported infectious etiologies to be the most common risk factor for stroke in tertiary centers, with similar findings in Beijing, China, and Turkey. In referral centers located in areas with a large African or Mediterranean population, hemoglobinopathies including sickle cell disease (SCD) may be the most common cause<sup>15,17–20)</sup>. In East Asian countries, MMD is the most common etiologies in pediatric stroke<sup>9)</sup>. Inherited metabolic diseases such as mitochondrial encephalopathies, Fabry's disease, Menke's disease, organic acidemia, urea cycle defects and homocystinuria can result in acute focal neurological deficits and stroke-like episodes. In cases of radiological evidence of infarction that is not confined to a known vascular territory, metabolic causes may be a consideration<sup>6,21,22)</sup>.

For assessment of etiologies and risk factors, there are no clearly established guidelines in pediatric stroke. Laboratory assessment including a variety of nonspecific blood tests and more specific tests searching for specific causes of stroke, such as coagulopathies, hematologic disorders or vasculitides are suggested (Table 2).

## DATA FROM TERTIARY CENTERS IN KOREA

We reviewed our cohort of 185 patients with pediatric stroke,

**Table 2.** Laboratory and diagnostic investigation for pediatric stroke

Laboratory tests to consider	Additional tests to consider
Liver function test	Brain MRI
CBC with ESR	Brain MRA
CRP	Intracranial vessels
ANA	Extracranial great vessels (neck)
Lupus anticoagulant/Anticardiolipin antibody	MR venography
Factor V Leiden mutation	Diffusion weighted image
Protein C. Protein S function	CT angiogram
Antithrombin III	Cerebral angiogram
Prothrombin gene mutation	Transesophageal echo
Homocysteine level	EEG
Methyltetrahydrofolate reductase gene mutation ( <i>MTHFR</i> )	Lumbar puncture
Fibrinogen/Plasminogen activator inhibitor	Holter monitoring
Factor VII/VIII elevation, Factor XII deficiency	Transcranial Doppler
Plasma aminoacid/urine aminoacid, organic acid analysis	
Serum and CSF lactate/Pyruvate	
TG/cholesterol/Lipoprotein	
Serum/CSF varicella titer	

CBC : complete blood cell count, ESR : erythrocyte sedimentation rate, CRP : C-reactive protein, ANA : antinuclear antibody, CSF : cerebrospinal fluid, TG : triglycerides, MRA : magnetic resonance angiography, MR : magnetic resonance

who was confirmed as stroke by magnetic resonance imaging at our center, between January 2002 and June 2013, to analyze the risk factors/etiologies and their clinical outcomes based on stroke type and ages. Cases of neonatal and perinatal stroke were excluded. Other childhood cerebrovascular disorders that were excluded included transient ischemic attack without infarction, hypotensive watershed injury, reversible hypertensive leukoencephalopathy, diffuse hypoxic encephalopathies, and asymptomatic infarction.

Stroke was classified into AIS, CSVT, metabolic stroke (MS), and hemorrhagic stroke. MS results from energy failure caused by metabolic conditions such as mitochondrial disease, organic acidemias, and single-gene diseases. Risk factors for AIS were subdivided into four groups : arteriopathy, cardiac disease, prothrombotic condition, and hematologic disease.

Among 185 patients, male dominance was more prominent in adolescence (sex ratio, 3.5 : 1) than it was in preadolescence. The mean age at diagnosis was 6.5 years.

Most patients (n=149, 81%) had two of more symptoms at onset. One hundred seventy-four patients (94%) presented with focal neurological signs : paresis (n=132, 71%), seizure (n=71, 38%), speech impairment, paresthesia, and other focal neurological deficits. Diffuse signs were documented in 102 patients (55%) : headache (n=50, 27%), altered mentality (n=34, 18%), lethargy, nausea/vomiting, dizziness, and other diffuse signs. Ninety-one patients (49%) had both focal and diffuse signs (mixed signs).

There were 156 (156/185, 84.3%) patients with AIS, nine patients with CSVT, nine patients with MS, and 11 patients with HS. Presumptive risk factors were identified in 154 patients (83%).

In AIS, most common risk factors were arteriopathy, followed by cardiac disease, prothrombotic condition, and hematologic disease. Arteriopathy (n=83) includes MMD; CNS infection; inflammatory vasculopathy; systemic vasculitis; arterial dissection; other noninflammatory vasculopathies. Twenty-three patients had underlying cardiac diseases; congenital heart disease, infective endocarditis, arrhythmia, myocarditis, cardiomyopathy, and left atrial myxoma. Prothrombotic condition (n=14) includes methylenetetrahydrofolate reductase (*MTHFR*) gene mutation, antiphospholipid syndrome (APS) and chemotherapy. Hematologic disorders were detected on a single case basis in six patients; leukemia, iron-deficiency anemia, hemophagocytic lymphohistiocytosis, hemolytic uremic syndrome, thrombotic thrombocytopenic purpura, and suprasellar craniopharyngioma. CSVT was identified in nine patients : severe dehydration (n=3); infection (n=3); prothrombotic condition (n=3) such as APS, *MTHFR* gene mutation, and L-asparaginase. Nine patients were newly diagnosed as having mitochondrial disease : eight patients had mitochondrial encephalopathy with lactic acidosis and stroke-like episodes (MELAS), and were confirmed to have the mitochondrial DNA 3243 point mutation. Hemorrhagic stroke affected 11 patients. There were arteriovenous malformation, MMD, arterial aneurysm, arterial dissection, mycotic embolism, *MTHFR* gene mutation, and Evans syndrome.

Table 3 summarizes the risk factor by stroke types, including previous publication by Lee et al.<sup>9)</sup>. The most common single etiology was MMD in both centers.

The most common etiology/risk factors are different according to ages in pediatric stroke. Stroke in children under 1 year of age is more likely to be associated with CNS infection, whereas older children were more likely to have vascular diseases in-

**Table 3.** Risk factors according to stroke type

Risk Factors	n (%) (n : 185, SNUCH)	n (%) (n : 62, Lee et al. <sup>9)</sup> )
AIS	156 (84)	62
Arteriopathy	83 (44.9)	31 (50)
MMD	52	19
CNS infection	11	9
Inflammatory vasculopathy	10	
Systemic vasculitis	4	1
Arterial dissection	3	
Other vasculopathy	3	2
Cardiac disease	23	12 (19.3)
Prothrombotic condition	14	1
Hematologic disease	6	
Undetermined	30	10
CSVT	9 (5)	
Severe dehydration	3	
Infection	3	
Prothrombotic condition	3	
MS	9 (5)	8 (14.5)
MELAS	8	8
Mitochondrial disease	1	
HS	11 (6)	
AVM	3	
MMD	2	
Arterial aneurysm	1	
Arterial dissection	1	
CNS infection	1	
Hematologic disease	1	
Prothrombotic condition	1	
Undetermined	1	

AIS : arterial ischemic stroke, CSVT : cerebral sinovenous thrombosis, MS : metabolic stroke, HS : hemorrhagic stroke, MMD : moyamoya disease, MELAS : mitochondrial encephalopathy with lactic acidosis and stroke-like episodes, AVM : arteriovenous malformation, SNUCH : Seoul National University Children's Hospital

**Table 4.** Most common risk factors according to age group\*

Age group	Most common risk factor	n (%)
1-11 months	CNS infection	8 (24)
	Cardiac disease	7 (21)
	Severe dehydration	3 (9)
1-5 years	MMD	31 (51)
	Cardiac disease	5 (8)
	Inflammatory vasculopathy	4 (7)
6-11 years	MMD	18 (33)
	Prothrombotic condition	6 (11)
	Metabolic disease	5 (9)
≥12 years	Cardiac disease	8 (22)
	Prothrombotic condition	6 (18)
	Metabolic disease	3 (8)

\*Data from Seoul National University Children's Hospital. MMD : moyamoya disease

cluding MMD. Cardiac diseases are relatively associated with all age groups (Table 4).

## CONCLUSION

Pediatric stroke is an importance cause of disability and mortality. However, since the incidence is still low, risk factors are variable according to ethnic groups and ages and etiologies are different, the symptoms are more non-specific compare to adults, it remains a challenge to early diagnosis and appropriate management.

Therefore, high suspicion is important for pediatric neurologists and early recognition, diagnosis, and appropriate treatment based on etiologies/risk factor can significantly improve the clinical outcome and quality of life.

## References

- Bowers KJ, Deveber GA, Ferriero DM, Roach ES, Vexler ZS, Maria BL : Cerebrovascular disease in children : recent advances in diagnosis and management. *J Child Neurol* 26 : 1074-1100, 2011
- Cárdenas JF, Rho JM, Kirton A : Pediatric stroke. *Childs Nerv Syst* 27 : 1375-1390, 2011
- deVeber G, Roach ES, Riela AR, Wiznitzer M : Stroke in children : recognition, treatment, and future directions. *Semin Pediatr Neurol* 7 : 309-317, 2000
- Earley CJ, Kittner SJ, Feeser BR, Gardner J, Epstein A, Wozniak MA, et al. : Stroke in children and sickle-cell disease : Baltimore-Washington Cooperative Young Stroke Study. *Neurology* 51 : 169-176, 1998
- Freundlich CL, Cervantes-Arslanian AM, Dorfman DH : Pediatric stroke. *Emerg Med Clin North Am* 30 : 805-828, 2012
- Fullerton HJ, Wu YW, Sidney S, Johnston SC : Risk of recurrent childhood arterial ischemic stroke in a population-based cohort : the importance of cerebrovascular imaging. *Pediatrics* 119 : 495-501, 2007
- Ganesan V, Prengler M, McShane MA, Wade AM, Kirkham FJ : Investigation of risk factors in children with arterial ischemic stroke. *Ann Neurol* 53 : 167-173, 2003
- Kissela BM, Khoury JC, Alwell K, Moomaw CJ, Woo D, Adeyoye O, et al. : Age at stroke : temporal trends in stroke incidence in a large, biracial population. *Neurology* 79 : 1781-1787, 2012
- Lee EH, Yum MS, Ko TS : Risk factors and clinical outcomes of childhood ischemic stroke in a single Korean tertiary care center. *J Child Neurol* 27 : 485-491, 2012
- Lynch JK, Hirtz DG, DeVeber G, Nelson KB : Report of the National Institute of Neurological Disorders and Stroke workshop on perinatal and childhood stroke. *Pediatrics* 109 : 116-123, 2002
- Mackay MT, Wiznitzer M, Benedict SL, Lee KJ, Deveber GA, Ganesan V; International Pediatric Stroke Study Group : Arterial ischemic stroke risk factors : the International Pediatric Stroke Study. *Ann Neurol* 69 : 130-140, 2011
- Meyer-Heim AD, Boltshauser E : Spontaneous intracranial haemorrhage in children : aetiology, presentation and outcome. *Brain Dev* 25 : 416-421, 2003
- Moharir M, Deveber G : Pediatric arterial ischemic stroke. *Continuum (Minneapolis)* 20 (2 Cerebrovascular Disease) : 370-386, 2014
- Nelson KB, Lynch JK : Stroke in newborn infants. *Lancet Neurol* 3 : 150-158, 2004
- Per H, Unal E, Poyrazoglu HG, Ozdemir MA, Donmez H, Gumus H, et al. : Childhood stroke : results of 130 children from a reference center in Central Anatolia, Turkey. *Pediatr Neurol* 50 : 595-600, 2014

16. Rafay MF, Pontigon AM, Chiang J, Adams M, Jarvis DA, Silver F, et al. : Delay to diagnosis in acute pediatric arterial ischemic stroke. *Stroke* 40 : 58-64, 2009
17. Salih MA, Abdel-Gader AG, Al-Jarallah AA, Kentab AY, Gadelrab MO, Alorainy IA, et al. : Infectious and inflammatory disorders of the circulatory system as risk factors for stroke in Saudi children. *Saudi Med J* 27 Suppl 1 : S41-S52, 2006
18. Salih MA, Al-Jarallah AS, Abdel-Gader AG, Al-Jarallah AA, Al-Saadi MM, Kentab AY, et al. : Cardiac diseases as a risk factor for stroke in Saudi children. *Saudi Med J* 27 Suppl 1 : S61-S68, 2006
19. Salih MA, Murshid WR, Zahraa JN, Abdel-Gader AG, Al-Jarallah AA, Kentab AY, et al. : Congenital and genetic cerebrovascular anomalies as risk factors for stroke in Saudi children. *Saudi Med J* 27 Suppl 1 : S53-S60, 2006
20. Shi KL, Wang JJ, Li JW, Jiang LQ, Mix E, Fang F, et al. : Arterial ischemic stroke : experience in Chinese children. *Pediatr Neurol* 38 : 186-190, 2008
21. Testai FD, Gorelick PB : Inherited metabolic disorders and stroke part 1 : Fabry disease and mitochondrial myopathy, encephalopathy, lactic acidosis, and strokelike episodes. *Arch Neurol* 67 : 19-24, 2010
22. Testai FD, Gorelick PB : Inherited metabolic disorders and stroke part 2 : homocystinuria, organic acidurias, and urea cycle disorders. *Arch Neurol* 67 : 148-153, 2010
23. Tsze DS, Valente JH : Pediatric stroke : a review. *Emerg Med Int* 2011 : 734506, 2011
24. Younkin DP : Diagnosis and treatment of ischemic pediatric stroke. *Curr Neurol Neurosci Rep* 2 : 18-24, 2002