

Predictive Factors of Postoperative Pain and Postoperative Anxiety in Children Undergoing Elective Circumcision: A Prospective Cohort Study

³rd Department of Surgery, General University Hospital "ATTIKON", Athens,

*Penteli General Children's Hospital, Palaia Penteli, Athens,

[†]General Hospital of Lamia, Lamia,

[‡]¹st Department of General Surgery, General University Hospital "LAIKO", Athens, Greece

Nick Zavras, Stella Tsamoudaki*, Vasileia Ntomi, Ioannis Yiannopoulos[†],
Efstratios Christianakis*, and Emmanuel Pikoulis[‡]

Background: Although circumcision for phimosis in children is a minor surgical procedure, it is followed by pain and carries the risk of increased postoperative anxiety. This study examined predictive factors of postoperative pain and anxiety in children undergoing circumcision.

Methods: We conducted a prospective cohort study of children scheduled for elective circumcision. Circumcision was performed applying one of the following surgical techniques: sutureless prepuceplasty (SP), preputial plasty technique (PP), and conventional circumcision (CC). Demographics and base-line clinical characteristics were collected, and assessment of the level of preoperative anxiety was performed. Subsequently, a statistical model was designed in order to examine predictive factors of postoperative pain and postoperative anxiety. Assessment of postoperative pain was performed using the Faces Pain Scale (FPS). The Post Hospitalization Behavior Questionnaire study was used to assess negative behavioral manifestations.

Results: A total of 301 children with a mean age of 7.56 ± 2.61 years were included in the study. Predictive factors of postoperative pain measured with the FPS included a) the type of surgical technique, b) the absence of siblings, and c) the presence of postoperative complications. Predictive factors of postoperative anxiety included a) the type of surgical technique, b) the level of education of mothers, c) the presence of preoperative anxiety, and d) a history of previous surgery.

Conclusions: Although our study was not without its limitations, it expands current knowledge by adding new predictive factors of postoperative pain and postoperative anxiety. Clearly, further randomized controlled studies are needed to confirm its results. (Korean J Pain 2015; 28: 244-253)

Key Words: Anxiety, Child; Circumcision; Pain measurement; Phimosis; Postoperative pain; Predictive value of tests; Questionnaires.

Received June 10, 2015. Revised August 20, 2015. Accepted August 31, 2015.

Correspondence to: Nick Zavras

³rd Department of Surgery, General University Hospital "ATTIKON", 32 Terpsitheas str, Agia Paraskevi 15341, Athens, Greece

Tel: +30-2106549325, Fax: +30-2105326411, E-mail: nzavras@med.uoa.gr

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © The Korean Pain Society, 2015

INTRODUCTION

Circumcision for phimosis is one of the most frequent interventions in pediatric surgery [1]. Although regarded as minor and routine, it is a painful procedure [2] which, compounded by general anesthesia and hospitalization, can prove to be a particularly distressing event for children [3]. Furthermore, these disturbing factors may cause such anxiety as to influence pre- and postoperative recovery. It has been shown that patients with lower preoperative anxiety have a better recovery and show fewer emotional and behavioral problems after discharge, while those displaying heightened anxiety have a greater likelihood of suffering a number of negative psychological effects such as anxiety, depression, irritability, aggressiveness and disturbances in their relationship with caregivers [2]. From a medical point of view, perioperative anxiety may affect the immune system by increasing sensitivity to infections [4] and subsequent consumption of analgesics postoperatively [5].

In the present study, we investigated risk factors that may influence the presence of postoperative anxiety and pain in children undergoing elective surgery for phimosis using three different surgical techniques.

MATERIALS AND METHODS

We conducted a prospective observational cohort study which took place from January 2010 to August 2014. Children included in the study were those with American Society of Anesthesiology (ASA) physical status I–II and aged 5 to 14 years that were scheduled for elective outpatient circumcision under general anesthesia. Exclusion criteria included any degree of cognitive and communicative impairment, bleeding disturbances, poor speaking and reading of the Greek language, refusal of parents to participate in the study, or loss to follow-up.

1. Objectives of the study

1) To determine the impact of demographics and base-line clinical characteristics on the degree of postoperative pain.

2) To examine the association between postoperative negative behavior manifestations (NBMs) and demographics, and base-line clinical characteristics.

2. Ethical considerations

The study was approved by the Institutional Review Board and informed consent was obtained from the parents.

3. Definitions

1) Pain: defined as an unpleasant sensory experience associated with tissue damage [6].

2) Anxiety: defined as a response of the brain to stimuli that the organism will attempt to avoid [7].

Table 1. Demographics and Base-line Clinical Characteristics

	Number (%)
Age (years)	
5–7	107 (35.6)
8–10	171 (56.8)
11–14	23 (7.6)
Nationality	
Greek	222 (73.8)
Other	79 (26.2)
Familial status	
Married	274 (91.0)
Divorced	27 (9.0)
Father's education level	
Basic education	173 (57.5)
High education	128 (42.5)
Mother's education level	
Basic education	185 (61.5)
High education	116 (38.5)
Birth order	
1st	122 (40.5)
≥ 2nd	179 (59.5)
Siblings	
Yes	269 (89.4)
No	32 (10.6)
Health problems	
No	292 (97.0)
Yes	9 (3.0)
History of previous surgery	
No	190 (63.1)
Yes	111 (36.9)
Scheduled surgery	
No	6 (2.0)
Yes	295 (98.0)
Preoperative preparation	
Not at all/a little	24 (8.0)
Moderate	99 (32.9)
Satisfactory	178 (59.1)
Type of fimosis	
Congenital	285 (94.7)
Acquired	16 (5.3)

4. Preoperative protocol

All children and their parents attended a preparation program before surgery (7–10 days before admission to the hospital), which included a visit to the ward and operating room and receiving detailed information concerning the surgical procedure, type of anesthesia, duration of the operation and possible complications. Data collected at this time included age, nationality, familial status, educational level of parents, birth order, health problems, previous surgical experience, schedule of surgery, preparation, and type of phimosis (congenital or acquired) (Table 1).

On the day of surgery, after separating the children from their parents, preoperative anxiety was assessed in the operating room by an experienced anesthesiologist by using the modified Yale preoperative anxiety scale (m-YPAS) [8] (Table 2). This is a validated observational tool for assessing children's anxiety that focuses on five items: activity, emotional expressivity, state of arousal, vocalization and use of parents. Each item comprises four categories, with the exception of vocalization that has six categories. A partial score is allocated to each item, and the sum is divided by the number of categories within that item. The score of each item is added to the others and the sum is multiplied by 20. Children with a score 23.5–30 are classified as not suffering from anxiety, while a score > 30 denotes severe anxiety [9].

5. Intraoperative protocol

No premedication was given to the children, and the presence of parents at their child's anesthesia induction was not allowed. General anesthesia was delivered with atropine 0.01 mg/kg, propofol 3 mg/kg and fentanyl 1

μg/kg intravenously. When needed, rocuronium 0.8 mg/kg was used to enable a laryngeal mask of appropriate size to be put in place. Anesthesia was maintained with sevoflurane and O₂/N₂O. Mean arterial pressure (MAP), heart rate (HR), peripheral oxygen saturation (SPO₂), and capnography were monitored during anesthesia. Intraoperative analgesia was estimated on the basis of gross movements or changes in HR and MAP. Increases ≥ 20% of the initial values were recorded and considered as signs of inadequate analgesia.

Following induction of anesthesia and before the start of surgery, a subcutaneous ring block of 0.25% levobupivacaine (0.5% levobupivacaine diluted in normal saline, Chirocaine[®], Abbott Laboratories Ltd) was injected around the base of the penis at a dose of 0.1 ml/kg (total dose 0.5 mg/kg). The duration of each surgical procedure (from induction to completion of surgery) was defined as low (10–15 minutes), moderate (15–25 minutes) or increased (25–30 minutes).

Circumcision was performed according to the preference of surgeons. The following surgical techniques were used: conventional circumcision (CC), sutureless prepuceplasty (SP) [10] and preputial plasty (PP) [11]. The complications of the surgical procedures were rated.

6. Postoperative evaluation

On the ward, the doctor responsible (NZ) documented the type of surgical technique and any complication due to anesthetic or surgical interventions. An experienced nurse blinded to the surgical technique assessed postoperative pain according to Bieri's Faces Pain Scale [12]. FPS scores ≥ 3, were considered as an indication of pain.

Table 2. Univariate and Multivariate Logistic Regression Analysis Demonstrating the Most Significant Factors Related to Preoperative Increased Anxiety

	Univariate		Multivariate	
	Odds ratio (95% CI)	P	Odds ratio (95% CI)	P
Nationality				
Greek vs non-Greek	2.79 (1.823–8.248)	0.004	3.004 (1.474–6.124)	0.002
Familial status				
Both parents vs divorced	2.38 (0.98–5.82)	0.05	5.587 (2.624–11.890)	<0.001
Number of siblings				
None vs ≥1	1.39 (1.03–1.88)	0.03	2.728 (0.762–9.758)	0.123
Type of phimosis				
Congenital vs acquired	1.16 (1.00–1.35)	0.05	6.129 (0.782–48.020)	0.084

Paracetamol (20 mg/kg) per os or rectum was given when indicated.

7. After discharge

1) Children's NBMs were evaluated according to observations made at home on the 3rd postoperative day. We selected the 3rd postoperative day to avoid the influence of significant pain on NBMs. Studies have shown that pain decreases by the 3rd postoperative day [13]. Vernon's Post Hospitalization Behavior Questionnaire (PHBQ) [14] was chosen to estimate NBMs in children and was offered to the parents in a typed form. PHBQ consists of a list of 27 possible PB traits that children could exhibit. The following domains of anxiety are included in PHBQ: general anxiety, separation anxiety, sleep anxiety, eating disturbances, aggression against authority, and apathy/withdrawal. The children are rated by parents as exhibiting NBM according to the following: (1) much less than before surgery, (2) less than before surgery, (3) the same as before surgery, (4) more than before surgery, and (5) much more than before. The total score ranges from 27 to 135. A score from 27–80 represents fewer NBMs, 81 represents the same as before, and 81–135 represents more. PHBQ has been shown to demonstrate reliability and prediction of changes due to preoperative interventions [15].

2) At home, parents rated their child's pain according to the total doses of analgesics given on the 1st, 2nd, and 3rd postoperative day.

Parents were encouraged to visit the hospital on the 8th postoperative day to allow a second look at the wound, submit the written answers to the PHBQ and provide the recorded total of analgesics given. Alternatively, two doctors (NZ and VN) contacted families by phone.

8. Statistical analysis

The association of each one of the demographics and base-line clinical characteristics with m-YPAS scale and FPS were evaluated using a univariate logistic regression model of analysis. Subsequently, a multivariate logistic regression analysis was performed for those parameters that were proven to be significant in the univariate analysis in order to evaluate the independency of each association, and the statistical significant differences were recorded accordingly. Children were evaluated for NBMs on the 3rd postoperative day and were divided into two groups according to PHBQ scores: those with NBMs and those with-

out NBMs. These groups were further subdivided into two groups according to age: 5–7 years and 8–14 years respectively. Pearson's chi-square test, Fisher's exact test, and Mann-Whitney test were used for the comparison of continuous variables between the two groups. The impact of different factors was compiled and further analysed in a multiple logistic regression analysis in order to find independent factors associated with NBMs. Adjusted odds ratios (OR) with 95% confidence intervals (95% CI) were computed from the results of the logistic regression analyses. All reported *P* values were two-tailed. Statistical significance was set at *P* < 0.05 and analyses were conducted using SPSS statistical software (version 19.0).

RESULTS

After the exclusion of 23 children for various reasons, a total of 301 children (mean age: 7.56 ± 2.61 years) were enrolled in the study. A hundred and seven (35.5%) children were aged 5–7 years, 171 (56.8%) were aged 8–10 years, and 23 (7.6%) were aged 11–14 years. Sample demographics and base-line clinical characteristics are presented in **Table 1**.

The majority of children (73.8%) were of Greek nationality. In 9% of the sample, the parents were divorced. A separate health problem was reported in nine (3.0%) children, and 111 (36.9%) patients had a history of previous surgery. Surgery was scheduled in almost all cases (98%). In 178 (59.1%) of cases, the preoperative preparation was considered as satisfactory. Preoperative increased anxiety was recorded in 185 (61.5%) children. The univariate logistic regression analysis showed that m-YPAS was increased in children of non-Greek origin and children without siblings (*P*: 0.004, and *P*: 0.03, respectively). The same analysis revealed a borderline increase of m-YPAS for children of divorced parents (*P*: 0.05) and those having acquired phimosis (*P*: 0.05) (**Table 2**). However, the multivariate regression analysis showed that preoperative increased anxiety was observed only in children of divorced parents [OR: 5.587 (CI: 2.624–11.890)], *P*: < 0.001, and in those of non-Greek nationality [OR: 3.004 (95% CI: 1.474–6.124)], *P*: 0.002 (**Table 2**).

The SP technique was performed on 99 children (32.9%), PP on 93 (30.9%), and CC on 109 (36.2%) (**Table 3**). Regarding the duration of surgery, the mean time for the SP technique was significantly less (10.25 ± 1.1 minutes)

Table 3. Intraoperative and Postoperative Findings. Results are Expressed as Mean \pm SD and %

	SP	PP	CC	P
Number Of patients	99	93	109	
Duration of operation (min)	10.3 \pm 1.1	17.3 \pm 2.7	23.6 \pm 2.5	<0.001
FPS score	3.12 \pm 0.95	3.56 \pm 1.09	4.89 \pm 1.32	<0.001
Postoperative complications (no of pts)				
Oedema	20	30	47	<0.001
Hematoma	0	22	32	<0.001
Consumption of analgesics (in doses)				
1 st post. day	9	9	52	<0.001
2 nd post. day	6	5	16	NS
3 rd post. day	0	4	10	0.007
Total: 111	15	18	78	

Table 4. Multivariate Logistic Regression Analysis Regarding Demographics, Base-line Clinical Characteristics and FACES Scale

	Odds Ratio	SE	z	P	95% Conf, Interval of OR	
Surgical technique						
PP technique vs SP	10,245	7,232	3,300	0,001	2,568	40,870
CC technique vs SP	69,410	59,182	4,970	<0,001	13,051	369,147
Number of siblings						
None vs \geq 1	0,354	0,182	-2,020	0,043	0,130	0,968
Complications						
None vs yes	4,625	1,686	4,200	<0,001	2,264	9,450

as compared with PP (17.31 \pm 2.71 minutes) and CC (23.58 \pm 2.48 minutes) ($P < 0.001$). The mean FPS score was 4.37 \pm 1.48 (range 2–8). Patients submitted to CC exhibited statistically higher mean pain scores (FPS: $P < 0.001$) when compared with the other surgical procedures. Postoperative complications were recorded in 27.7% of cases (edema in 18.7% of the cases, hematoma in 6.7% and both edema and hematoma in 2.3%). The majority (56.3%) were noted in children who underwent the CC procedure (**Table 3**). Regarding the intake of analgesics at home, a total of 111 doses of analgesics were given over the 1st, 2nd, and 3rd postoperative days: specifically, 70 (63%) doses were given on the 1st postoperative day, 27 (24.3%) on the 2nd and 14 (12.7%) on the 3rd postoperative day.

The majority (70.2%) were needed for those children who underwent the CC technique. The results of the multivariate regression analysis showed a strong correlation between the type of surgical technique (PP versus SP and CC versus SP) and the presence of postoperative pain as-

essed with the FPS [OR: 10.24 and 69.41 (95%CI: 2.568–40.87 and 13.051–369.14 respectively), $P < 0.001$, and $P < 0.001$ respectively] (**Table 4**).

On the 3rd postoperative day, 98 (32.6%) children presented NBMs. **Table 5** summarizes the results of the most significant factors that are influenced in the presence of NBMs based on the univariate logistic model. According to this, parental status, mother education level, birth order, no previous experience of surgery, the type of phimosis, the surgical technique, the nationality, the duration of surgery and the level of preoperative anxiety seems to influence the presence of NBMs. More specifically, NBMs were seen in cases of children whose parents were divorced and whose mothers had reached only a basic level of education. Significantly higher percentages of NBMs were associated with the first born child and those with no previous experience of surgery. Moreover, the proportion of children with NBMs on the 3rd postoperative day was significantly lower in SP cases as compared with both CC and PP ($P < 0.001$ for both comparisons), while PP cases exhibited fewer

Table 5. NBMs on 3rd Postoperative Day Associated with the Most Significant Demographic Factors and Clinical Characteristics

Variables	No NBMs (%)	NBMs (%)	P
Nationality			
Greek	159 (70.7)	63 (29.3)	0.015*
Other	44 (55.7)	35 (44.3)	
Familial status			
Married	180 (65.7)	94 (34.3)	0.039*
Divorced	23 (85.2)	4 (14.8)	
Mother's level of education			
Basic education	97 (52.4)	88 (47.6)	<0.001*
High education	106 (91.4)	10 (8.6)	
Birth order			
1st	71 (58.2)	51 (41.8)	0.005*
≥ 2nd	132 (73.7)	47 (26.3)	
Type of phimosis			
Congenital	187 (65.6)	98 (34.4)	0.004**
Acquired	16 (100.0)	0 (0.0)	
Surgical technique			
SP	91 (91.9)	8 (8.1)	<0.001*
PP	72 (77.4)	21 (22.6)	
CC	40 (36.7)	69 (63.3)	<0.001*
Low	92 (97.9)	2 (2.1)	
Moderate	109 (85.8)	18 (14.2)	
Increased	2 (2.5)	78 (97.5)	
Preoperative increased anxiety			
Yes	1 (0.6)	184 (99.4)	<0.001*
No	22 (18.9)	94 (81.1)	

*Pearson's chi-square test, **Fisher's exact test.

NBMs than those submitted to CC ($P < 0.001$). The proportion of children with NBMs on the 3rd postoperative day was significantly higher among children of non-Greek origin and in cases of congenital phimosis ($P; 0.015$). Additionally, the longer the duration of surgery, the higher the number of children with NBMs ($P < 0.001$). Finally, children with higher levels of preoperative anxiety were found to be at higher risk of developing NBMs postoperatively ($P < 0.001$).

However, when a multiple logistic regression model was applied, it was found that the surgical technique, mother's educational level, preoperative increased anxiety and history of previous surgery were all significantly associated with NBMs (**Table 6**).

More specifically, children who underwent PP or CC had an estimated greater risk (3.32- and 19.62-fold, respectively) of having NBMs as compared to those who underwent SP surgery. Children whose mothers had achieved

Table 6. Multiple Logistic Regression Results with NBMs Changes on the 3rd Postoperative Day as Dependent Variable

	OR (95% CI) †	P
Surgical technique		
SP	1.00*	
PP	3.32 (1.39–7.93)	0.007
Circumcision	19.62 (8.63–44.60)	<0.001
Mother's education		
Basic education	1.00	
High education	0.09 (0.03–0.24)	<0.001
Preoperative increased anxiety		
No	1.00	
Yes	4.93 (1.88–12.89)	0.001
History of previous surgery		
No	1.00	
Yes	7.96 (3.97–15.93)	<0.001

*Indicates reference category, †Odds Ratio (95% Confidence Interval).

a high level of education had 91% lower probability of having NBMs as compared with those whose mothers had reached a basic level of education [OR: 0.09 (95% CI: 0.03–0.24), $P < 0.001$] (Table 6). In addition, children with preoperative increased anxiety had 4.93 times greater probability of having NBMs, while those with experience of previous surgery had a likelihood of almost 8 times greater (Table 6).

DISCUSSION

A considerable number of previous studies have focused on the impact of various factors on postoperative NBMs. These factors included preoperative anxiety of parent and child, child's temperament, previous experience of hospitalization for surgery, length of hospitalization, preoperative preparation, preoperative premedication, the presence of parents at the induction of anesthesia and in the post-anesthetic care unit, or the different use of anesthetic drugs provided at induction in anesthesia, [5,16–24]. In the present study, along with some of the above mentioned factors, we studied new factors such as familial status, nationality, and the birth order of children in order to gain more specific information concerning the influence of day-case surgery on NBMs post-discharge.

Liu et al. [25] examined the behavioral problems of children of divorced parents and found that such children had negative consequences including aggressive behavior, withdrawal, and social problems as compared with children with both parents. Similarly, Weaver and Schofield [26] showed that children of divorced families exhibited more behavioral problems compared to those with intact families. However, while the results of our study showed a statistical significance of NBMs in children from divorced families when compared against children with both parents ($P: 0.039$), the regression analysis did not confirm this observation.

Various stress factors, such as language, discrimination, and parental acculturative stress may be implicated in the presence of psychological distress among immigrant children [27–29]. All these factors come under the term acculturation as they are related to the process of cultural psychological changes after the meeting of different cultures between individuals [30]. Acculturation has been studied at length in young adolescents and adults. However, it has not been studied extensively in children

[31]. In our study, a statistically significant increase of postoperative NBMs ($P < 0.001$) in children of non-Greek nationality was noticed which may reflect the effect of stress factors on the emotional reaction of these children.

Karling and Hägglöf [32] reported that the basic education level of parents acted as a protective factor against general anxiety. Furthermore, they identified that higher education acted as a risk factor for specific NBMs such as general anxiety, regression and apathy/withdrawal [32]. One possible explanation could be the capacity of a highly educated mother to have a better understanding of the harmful aspects of stress. Conversely, the results of both the univariate and multivariate analysis showed that the lower education level of parents may be a contributory factor for postoperative NBMs. Furthermore, in line with Karling and Hägglöf [32], the present study found no evidence that the father's educational level had an impact on postoperative NBMs.

In the present study, a history of previous surgery was a predictive factor for postoperative NBMs, as a statistical difference was observed among children with a surgery-free history and NBMs and those having had previous surgery ($P < 0.001$). Furthermore, the multivariate logistic model indicated that previous surgical history as an independent variable is a crucial factor for the occurrence of postoperative NBMs. Kain et al. [33], and Kotiniemi et al. [34] reported that children's surgical or medical histories may either worsen or minimize anxiety, depending on the quality of a dreadful event. All these conflicting results indicate that the influence of a previous experience of surgery has not yet been clarified for various reasons such as different methods for assessment of anxiety and different statistical analyses [5].

Notably, the statistical analysis showed that congenital phimosis was a contributory factor for postoperative NBMs, but this observation was not noticed in the multivariate analysis. Furthermore, children submitted to the SP surgical technique expressed less NBMs postoperatively, and this finding was confirmed in the logistic regression model. Although no accurate results could be extracted from this observation, one could hypothesize that this finding is related to less postoperative pain which was observed with the SP technique. Indeed, children undergoing SP suffered less pain when compared with the other two techniques ($P < 0.001$) and this finding was supported by the multivariate analysis. Another significant finding regarding the

surgical technique is that of the duration of surgery. Our result showed that this technique required less operating time, a factor that may influence the presence of postoperative anxiety, i.e. the longer the surgical intervention, the greater the chance of postoperative NBMs [17]. It is noteworthy that postoperative complications were not associated with postoperative NBMs.

Surprisingly, no impact of age on postoperative NBMs was noticed in this study, as opposed to several other studies [33,35,36] that showed age to be a potential factor for postoperative maladaptive behavioral changes. Most of the children affected were under four years of age [35,36]. However, our findings concur with those of Thompson and Vernon [36], in which no association between behavioral changes and age was found. Karling et al. [16] hypothesized that a lack of questions related to behavior in older children may be responsible for this discrepancy. Regarding birth order, our results showed that first born children suffered more from NBMs as compared with second born and third born children ($P: 0.005$). Although this finding was not verified as a predictor factor in the logistic analysis, it is a new finding which should be investigated in future studies.

Of particular interest was the observation that children with increased preoperative anxiety had a statistically greater risk of postoperative NBMs according to both categorical data and multivariate regression analysis. This observation is in line with the studies by Kain et al. [20] and Fortier et al. [37]. Kain et al. [38] speculated that a possible explanation could be the underlying temperament of the child and the feasibility of adaptation to stressful environmental stimuli. In contrast, preoperative preparation was found not to have any impact in the presence of NBMs postoperatively. Our results are consistent with the findings of Kain et al. [39] who reported that no benefits were gained from behavioral preparation programs in the postoperative period. On the contrary, Margolis et al. [40] noticed that a preparation program based on age adjustments may have a beneficial effect on children postoperatively. All in all, the results published in the international literature are conflicting and further investigation is needed to resolve this issue.

A substantial number of studies have shown that increased preoperative anxiety is associated with higher levels of postoperative pain in children undergoing various surgical procedures [20,41–43]. However, this correlation

was not found in the present study. Notably, the multivariate logistic analysis showed higher preoperative anxiety in children of divorced parents, and those of non-Greek nationality.

On the basis of our results, the mean FPS score was higher in younger children aged 5–7 years when compared to children aged 8–10 and 11–14 years ($P: 0.0018$ and $P: 0.003$ respectively). This evidence is supported by the study of Palermo and Drotar [44]. However, in the proposed multivariate analysis, no relationship between age and postoperative pain was found. This finding is in accordance with a study by Logan and Rose [45] who found no relationship between age and postoperative pain. Conversely, Crandall et al. [46] found that postoperative pain increases with age. In summary, results regarding the influence of age in the perception of pain are inconclusive.

In this study, previous surgical experience, as an independent factor, did not correlate with the intensity of postoperative pain; this finding was in accordance with the univariate statistical analysis. This topic has not yet been clarified. The study by Ericsson et al. [43] supports our results, while Crandall et al. [46] reported lower pain scores in children with previous surgical experience. All these contradictory results reflect the interaction of multiple factors such as the conditions of hospitalization, the reasons for surgery, the presence of postoperative complications, and outcome that may be involved with the presence of postoperative pain [43].

Although nationality, preoperative preparation and the duration of surgery seemed to have an influence on pain intensity postoperatively, these variables had no predictability as separate factors in our model of multivariate analysis. On the contrary, the type of surgical technique, absence of siblings, and presence of postoperative complications seem to affect the intensity of postoperative pain as independent factors. In a previous study [47], the authors noticed that postoperative pain in circumcision was less in cases where cyanoacrylate was used for approximation of wound closure rather than sutures. One possible explanation could be the traction effect of the trauma sutures once they come in contact with clothing [47]. This evidence was supported in our study in cases submitted to the SP technique in which no sutures were used. In addition, the presence of postoperative complications was a predictor of postoperative pain, and the SP technique was accompanied by a lower rate of complications.

Our study had a number of limitations. For example, the study only included children operated on for circumcision. Furthermore, preoperative parental anxiety, children's temperament, and preoperative anxiety of the children in the ward or at induction of anesthesia were not assessed. Furthermore, postoperative behavior alterations were measured once and may therefore not reflect a long-term impact on behavioral status of children. However, some new observations, such as the impact of nationality, the mothers' education level, and the order of birth, should be further investigated in future studies.

REFERENCES

- Beyaz SG. Comparison of postoperative analgesic efficacy of caudal block versus dorsal penile nerve block with levobupivacaine for circumcision in children. *Korean J Pain* 2011; 24: 31–5.
- Tree-Trakarn T, Pirayavaraporn S. Postoperative pain relief for circumcision in children: comparison among morphine, nerve block, and topical analgesia. *Anesthesiology* 1985; 62: 519–22.
- Wollin SR, Plummer JL, Owen H, Hawkins RM, Materazzo F, Morrison V. Anxiety in children having elective surgery. *J Pediatr Nurs* 2004; 19: 128–32.
- Ader R, Cohen N, Felten D. Psychoneuroimmunology: interactions between the nervous system and the immune system. *Lancet* 1995; 345: 99–103.
- Caumo W, Broenstrup JC, Fialho L, Petry SM, Brathwait O, Bandeira D, et al. Risk factors for postoperative anxiety in children. *Acta Anaesthesiol Scand* 2000; 44: 782–9.
- Bonica JJ. The need of a taxonomy. *Pain* 1979; 6: 247–8.
- Beesdo K, Knappe S, Pine DS. Anxiety and anxiety disorders in children and adolescents: developmental issues and implications for DSM–V. *Psychiatr Clin North Am* 2009; 32: 483–524.
- Kain ZN, Mayes LC, Cicchetti DV, Bagnall AL, Finley JD, Hofstadter MB. The Yale Preoperative Anxiety Scale: how does it compare with a "gold standard"? *Anesth Analg* 1997; 85: 783–8.
- Cumino Dde O, Cagno G, Gonçalves VF, Wajman DS, Mathias LA. Impact of preanesthetic information on anxiety of parents and children. *Braz J Anesthesiol* 2013; 63: 473–82.
- Christianakis E. Sutureless prepuceplasty with wound healing by second intention: an alternative surgical approach in children's phimosis treatment. *BMC Urol* 2008; 8: 6.
- Cuckow PM, Rix G, Mouriquand PD. Preputial plasty: a good alternative to circumcision. *J Pediatr Surg* 1994; 29: 561–3.
- Bieri D, Reeve RA, Champion GD, Addicoat L, Ziegler JB. The Faces Pain Scale for the self-assessment of the severity of pain experienced by children: development, initial validation, and preliminary investigation for ratio scale properties. *Pain* 1990; 41: 139–50.
- Gedaly-Duff V, Ziebarth D. Mothers' management of adenoid-tonsillectomy pain in 4- to 8-year-olds: a preliminary study. *Pain* 1994; 57: 293–9.
- Vernon DT, Schulman JL, Foley JM. Changes in children's behavior after hospitalization. Some dimensions of response and their correlates. *Am J Dis Child* 1966; 111: 581–93.
- Vernon DT, Thompson RH. Research on the effect of experimental interventions on children's behavior after hospitalization: a review and synthesis. *J Dev Behav Pediatr* 1993; 14: 36–44.
- Karling M, Stenlund H, Hägglöf B. Child behaviour after anaesthesia: associated risk factors. *Acta Paediatr* 2007; 96: 740–7.
- Kain ZN, Wang SM, Mayes LC, Caramico LA, Hofstadter MB. Distress during the induction of anesthesia and postoperative behavioral outcomes. *Anesth Analg* 1999; 88: 1042–7.
- Stargatt R, Davidson AJ, Huang GH, Czarnecki C, Gibson MA, Stewart SA, et al. A cohort study of the incidence and risk factors for negative behavior changes in children after general anesthesia. *Paediatr Anaesth* 2006; 16: 846–59.
- Yuki K, Daaboul DG. Postoperative maladaptive behavioral changes in children. *Middle East J Anaesthesiol* 2011; 21: 183–9.
- Kain ZN, Mayes LC, Caldwell-Andrews AA, Karas DE, McClain BC. Preoperative anxiety, postoperative pain, and behavioral recovery in young children undergoing surgery. *Pediatrics* 2006; 118: 651–8.
- Kain ZN, Mayes LC, Wang SM, Hofstadter MB. Postoperative behavioral outcomes in children: effects of sedative premedication. *Anesthesiology* 1999; 90: 758–65.
- Watson AT, Visram A. Children's preoperative anxiety and postoperative behaviour. *Paediatr Anaesth* 2003; 13: 188–204.
- Lardner DR, Dick BD, Crawford S. The effects of parental presence in the postanesthetic care unit on children's postoperative behavior: a prospective, randomized, controlled study. *Anesth Analg* 2010; 110: 1102–8.
- Aguilera IM, Patel D, Meakin GH, Masterson J. Perioperative anxiety and postoperative behavioural disturbances in children undergoing intravenous or inhalation induction of anaesthesia. *Paediatr Anaesth* 2003; 13: 501–7.
- Liu X, Guo C, Okawa M, Zhai J, Li Y, Uchiyama M, et al. Behavioral and emotional problems in Chinese children of divorced parents. *J Am Acad Child Adolesc Psychiatry* 2000; 39: 896–903.
- Weaver JM, Schofield TJ. Mediation and moderation of divorce effects on children's behavior problems. *J Fam Psychol* 2015; 29: 39–48.
- Duru E, Poyrazli S. Personality dimensions, psychosocial-

- demographic variables, and English language competency in predicting level of acculturative stress among Turkish international students. *Int J Stress Manag* 2007; 14: 99–110.
28. Suarez-Morales L, Lopez B. The impact of acculturative stress and daily hassles on pre-adolescent psychological adjustment: examining anxiety symptoms. *J Prim Prev* 2009; 30: 335–49.
 29. Behnke AO, MacDermid SM, Coltrane SL, Parke RD, Duffy S, Widaman KF. Family cohesion in the lives of Mexican American and European American parents. *J Marriage Fam* 2008; 70: 1045–59.
 30. Sam DL, Berry JW. Acculturation: when individuals and groups of different cultural backgrounds meet. *Perspect Psychol Sci* 2010; 5: 472–81.
 31. Leidy MS, Parke RD, Cladis M, Coltrane S, Duffy S. Positive marital quality, acculturative stress, and child outcomes among Mexican Americans. *J Marriage Fam* 2009; 71: 833–47.
 32. Karling M, Hägglöf B. Child behaviour after anaesthesia: association of socioeconomic factors and child behaviour checklist to the Post-Hospital Behaviour Questionnaire. *Acta Paediatr* 2007; 96: 418–23.
 33. Kain ZN, Mayes LC, O'Connor TZ, Cicchetti DV. Preoperative anxiety in children. Predictors and outcomes. *Arch Pediatr Adolesc Med* 1996; 150: 1238–45.
 34. Kotiniemi LH, Ryhänen PT, Moilanen IK. Behavioural changes in children following day-case surgery: a 4-week follow-up of 551 children. *Anaesthesia* 1997; 52: 970–6.
 35. Eckenhoff JE. Relationship of anesthesia to postoperative personality changes in children. *AMA Am J Dis Child* 1953; 86: 587–91.
 36. Thompson RH, Vernon DT. Research on children's behavior after hospitalization: a review and synthesis. *J Dev Behav Pediatr* 1993; 14: 28–35.
 37. Fortier MA, Del-Rosario AM, Martin SR, Kain ZN. Perioperative anxiety in children. *Paediatr Anaesth* 2010; 20: 318–22.
 38. Kain ZN, Caldwell-Andrews AA, Maranets I, McClain B, Gaal D, Mayes LC, et al. Preoperative anxiety and emergence delirium and postoperative maladaptive behaviors. *Anesth Analg* 2004; 99: 1648–54.
 39. Kain ZN, Caramico LA, Mayes LC, Genevro JL, Bornstein MH, Hofstadter MB. Preoperative preparation programs in children: a comparative examination. *Anesth Analg* 1998; 87: 1249–55.
 40. Margolis JO, Ginsberg B, Dear GL, Ross AK, Goral JE, Bailey AG. Paediatric preoperative teaching: effects at induction and postoperatively. *Paediatr Anaesth* 1998; 8: 17–23.
 41. Pederson C. Effect of imagery on children's pain and anxiety during cardiac catheterization. *J Pediatr Nurs* 1995; 10: 365–74.
 42. Bringuier S, Dadure C, Raux O, Dubois A, Picot MC, Capdevila X. The perioperative validity of the visual analog anxiety scale in children: a discriminant and useful instrument in routine clinical practice to optimize postoperative pain management. *Anesth Analg* 2009; 109: 737–44.
 43. Ericsson E, Wadsby M, Hultcrantz E. Pre-surgical child behavior ratings and pain management after two different techniques of tonsil surgery. *Int J Pediatr Otorhinolaryngol* 2006; 70: 1749–58.
 44. Palermo TM, Drotar D. Prediction of children's postoperative pain: the role of presurgical expectations and anticipatory emotions. *J Pediatr Psychol* 1996; 21: 683–98.
 45. Logan DE, Rose JB. Is postoperative pain a self-fulfilling prophecy? Expectancy effects on postoperative pain and patient-controlled analgesia use among adolescent surgical patients. *J Pediatr Psychol* 2005; 30: 187–96.
 46. Crandall M, Lammers C, Senders C, Braun JV. Children's tonsillectomy experiences: influencing factors. *J Child Health Care* 2009; 13: 308–21.
 47. Elemen L, Seyidov TH, Tugay M. The advantages of cyanoacrylate wound closure in circumcision. *Pediatr Surg Int* 2011; 27: 879–83.