

## STABILITY FOR A CUBIC FUNCTIONAL EQUATIONS IN NON-ARCHIMEDEAN NORMED SPACES

CHANG IL KIM\* AND CHANG HYEOb SHIN\*\*

ABSTRACT. In this paper, we investigate the functional equation  
 $f(3x+y) + f(3x-y) = f(x+2y) + 2f(x-y) + 6f(2x) + 3f(x) - 6f(y)$   
and prove the generalized Hyers-Ulam stability for it in non-Archimedean  
normed spaces.

### 1. Introduction and preliminaries

S. M. Ulam [15] raised a question concerning the stability of functional equations in 1940 : *Let  $G_1$  be a group and let  $G_2$  be a metric group with the metric  $d(\cdot, \cdot)$ . Given  $\epsilon > 0$ , does there exist a  $\delta > 0$  such that if a mapping  $h : G_1 \rightarrow G_2$  satisfies the inequality  $d(h(xy), h(x)h(y)) < \delta$  for all  $x, y \in G_1$ , then there exists a homomorphism  $H : G_1 \rightarrow G_2$  with  $d(h(x), H(x)) < \epsilon$  for all  $x \in G_1$ ?*

In 1941, Hyers [6] solved the Ulam problem for the case of approximately additive functions in Banach spaces. Since then, the stability of several functional equations have been extensively investigated by several mathematicians [2, 5, 7, 8]. Rassias [13], Jun and Kim [9] and Park and Jung [12] introduced the following functional equations

$$(1.1) \quad f(x + 2y) + 3f(x) = 3f(x + y) + f(x - y) + 6f(y)$$

and

$$(1.2) \quad f(2x + y) + f(2x - y) = 2f(x + y) + 2f(x - y) + 12f(x)$$

and

$$(1.3) \quad f(3x + y) + f(3x - y) = 3f(x + y) + 3f(x - y) + 48f(x)$$

---

Received November 13, 2014; Accepted July 22, 2015.

2010 Mathematics Subject Classification: Primary 39B82, 39B52.

Key words and phrases: generalized Hyers-Ulam stability, cubic functional equation, non-Archimedean space.

Correspondence should be addressed to Chang Hyeob Shin, [seashin@hanmail.net](mailto:seashin@hanmail.net).

and they established general solutions and the generalized Hyers-Ulam-Rassias stability problem for this functional equations, respectively. It is easy to see that the function  $f(x) = cx^3$  is a solution of the functional equations (1.1), (1.2) and (1.3). Thus, it is natural that (1.1), (1.2) and (1.3) are called a *cubic functional equations* and every solution of the cubic functional equation is said to be a *cubic mapping*.

In this paper, we consider the following functional equation

$$(1.4) \quad \begin{aligned} & f(3x + y) + f(3x - y) \\ &= f(x + 2y) + 2f(x - y) + 6f(2x) + 3f(x) - 6f(y). \end{aligned}$$

We prove the generalized Hyers-Ulam stability of (1.4) in complete non-Archimedean normed spaces.

A *valuation* is a function  $|\cdot|$  from a field  $\mathbb{K}$  into  $[0, \infty)$  such that, for any  $r, s \in \mathbb{K}$ , the following conditions hold: (i)  $|r| = 0$  if and only if  $r = 0$ , (ii)  $|rs| = |r||s|$ , (iii)  $|r + s| \leq \max\{|r|, |s|\}$ .

A field  $\mathbb{K}$  is called a *valued field* if  $\mathbb{K}$  carries a valuation. The usual absolute values of  $\mathbb{R}$  and  $\mathbb{C}$  are examples of valuations. If the triangle inequality is replaced by  $|r + s| \leq \max\{|r|, |s|\}$  for all  $r, s \in \mathbb{K}$ , then the valuation  $|\cdot|$  is called a *non-Archimedean valuation* and a field with a non-Archimedean valuation is called *non-Archimedean field*. If  $|\cdot|$  is a non-Archimedean valuation on  $\mathbb{K}$ , then clearly,  $|1| = |-1|$  and  $|n| \leq 1$  for all  $n \in \mathbb{N}$ .

**DEFINITION 1.1.** Let  $X$  be a vector space over a scalar field  $\mathbb{K}$  with a non-Archimedean nontrivial valuation  $|\cdot|$ . A function  $\|\cdot\| : X \rightarrow \mathbb{R}$  is called a *non-Archimedean norm (valuation)* if it satisfies the following conditions:

- (a)  $\|x\| = 0$  if and only if  $x = 0$ ,
- (b)  $\|rx\| = |r|\|x\|$ ,
- (c) the strong triangle inequality (ultrametric), that is,

$$\|x + y\| \leq \max\{\|x\|, \|y\|\}$$

for all  $x, y \in X$  and all  $r \in \mathbb{K}$ .

If  $\|\cdot\|$  is a non-Archimedean norm, then  $(X, \|\cdot\|)$  is called a *non-Archimedean normed space*.

Let  $(X, \|\cdot\|)$  be a non-Archimedean normed space. Let  $\{x_n\}$  be a sequence in  $X$ . Then  $\{x_n\}$  is said to be *convergent* if there exists an  $x \in X$  such that  $\lim_{n \rightarrow \infty} \|x_n - x\| = 0$ . In that case,  $x$  is called *the limit of the sequence*  $\{x_n\}$  and one denotes it by  $\lim_{n \rightarrow \infty} x_n = x$ . A sequence  $\{x_n\}$  is said to be *Cauchy* in  $(X, \|\cdot\|)$  if  $\lim_{n \rightarrow \infty} \|x_{n+p} - x_n\| = 0$  for all

$p \in \mathbb{N}$ . By (c) in Definition 1.1,

$$\|x_n - x_m\| \leq \max\{\|x_{j+1} - x_j\| \mid m \leq j \leq n-1\} \quad (n > m)$$

and hence a sequence  $\{x_n\}$  is Cauchy in  $(X, \|\cdot\|)$  if and only if sequence  $\{x_{n+1} - x_n\}$  converges to zero in  $(X, \|\cdot\|)$ . By a *complete non-Archimedean normed space* we mean one in which every Cauchy sequence is convergent.

Throughout this paper,  $X$  is a non-Archimedean normed space and  $Y$  a complete non-Archimedean normed space.

## 2. The Generalized Hyers-Ulam stability for (1.4)

In 2003, Jun and Kim [10] introduced the following cubic functional equation

$$(2.1) \quad f(x+2y) + f(x-2y) + 6f(x) = 4f(x+y) + 4f(x-y)$$

and proved the generalized Hyers-Ulam stability for it in Banach spaces. In this section, we prove the generalized Hyers-Ulam stability of functional equation (1.4) in complete non-Archimedean normed spaces. We start the following theorem.

**THEOREM 2.1.** *Let  $f : X \longrightarrow Y$  be a mapping. Then  $f$  satisfies (1.4) if and only if  $f$  is cubic.*

*Proof.* Suppose that  $f$  satisfies (1.4). Letting  $x = y = 0$  in (1.4), we have  $f(0) = 0$ . Letting  $y = 0$  in (1.4), we have

$$(2.2) \quad f(3x) - 3f(2x) - 3f(x) = 0$$

for all  $x \in X$  and letting  $x = 0$  in (1.4) and replacing  $y$  by  $x$ , we have

$$(2.3) \quad 7f(x) - f(-x) - f(2x) = 0$$

for all  $x \in X$ . Letting  $y = x$  in (1.4), we have

$$(2.4) \quad f(4x) - f(3x) - 5f(2x) + 3f(x) = 0$$

for all  $x \in X$ . By (2.2) and (2.4), we get

$$(2.5) \quad f(4x) = 2^3 f(2x)$$

for all  $x \in X$ . Replacing  $2x$  by  $x$  in (2.5), we get

$$(2.6) \quad f(2x) = 2^3 f(x)$$

for all  $x \in X$ . By (2.2) and (2.6), we get

$$(2.7) \quad f(3x) = 3^3 f(x)$$

for all  $x \in X$ . By (2.3) and (2.6), we get

$$(2.8) \quad f(-x) = -f(x)$$

for all  $x \in X$ . Replacing  $y$  by  $3y$  in (1.4), by (2.6) and (2.7), we have

$$(2.9) \quad 27f(x+y) + 27f(x-y) = f(x+6y) + 2f(x-3y) + 51f(x) - 6f(3y)$$

for all  $x, y \in X$ . Interchanging  $x$  and  $y$  in (2.9), by (2.8), we have

$$(2.10) \quad 27f(x+y) - 27f(x-y) = f(6x+y) - 2f(3x-y) + 51f(y) - 6f(3x)$$

for all  $x, y \in X$ . Replacing  $y$  by  $2y$  in (2.10), by (2.6), we have

$$(2.11) \quad \begin{aligned} & 27f(x+2y) - 27f(x-2y) \\ &= 8f(3x+y) - 2f(3x-2y) + 408f(y) - 6f(3x) \end{aligned}$$

for all  $x, y \in X$ . Letting  $y = -y$  in (2.11), by (2.8), we have

$$(2.12) \quad \begin{aligned} & 27f(x-2y) - 27f(x+2y) \\ &= 8f(3x-y) - 2f(3x+2y) - 408f(y) - 6f(3x) \end{aligned}$$

for all  $x, y \in X$ . By (2.11) and (2.12), we get

$$(2.13) \quad 8[f(3x+y) + f(3x-y)] - 2[f(3x+2y) + f(3x-2y)] - 12f(3x) = 0$$

for all  $x, y \in X$ . Letting  $x = \frac{x}{3}$  in (2.13), we have

$$f(x+2y) + f(x-2y) + 6f(x) = 4f(x+y) + 4f(x-y)$$

for all  $x, y \in X$  and so  $f$  is additive-quadratic-cubic [10]. By (2.6),  $f$  is cubic. The converse is trivial.  $\square$

For a given mapping  $f : X \rightarrow Y$ , we define the difference operator  $Df : X^2 \rightarrow Y$  by

$$\begin{aligned} Df(x, y) &= f(3x+y) + f(3x-y) \\ &\quad - f(x+2y) - 2f(x-y) - 6f(2x) - 3f(x) + 6f(y) \end{aligned}$$

for all  $x, y \in X$ .

**THEOREM 2.2.** *Let  $\phi : X^2 \rightarrow [0, \infty)$  be a mapping such that*

$$(2.14) \quad \lim_{n \rightarrow \infty} \frac{\phi(2^n x, 2^n y)}{|2|^{3n}} = 0$$

for all  $x, y \in X$  and let for each  $x \in X$ , the following limit

$$(2.15) \quad \lim_{n \rightarrow \infty} \max \left\{ \left\{ \frac{1}{|2|} \frac{\phi(2^{j-1}x, 0)}{|2|^{3(j-1)}} : 0 \leq j < n \right\} \cup \left\{ \frac{\phi(2^{j-1}x, 2^{j-1}x)}{|2|^{3(j-1)}} : 0 \leq j < n \right\} \right\}$$

denoted by  $\tilde{\phi}(x)$ , exist. Suppose that  $f : X \rightarrow Y$  is a mapping satisfying

$$(2.16) \quad \|Df(x, y)\| \leq \phi(x, y)$$

for all  $x, y \in X$ . Then there exists a cubic mapping  $C : X \rightarrow Y$  such that

$$(2.17) \quad \|C(x) - f(x)\| \leq \frac{1}{|2|^6} \tilde{\phi}(x)$$

for all  $x \in X$ . In addition, if the limit

$$(2.18) \quad \lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} \max \left\{ \left\{ \frac{1}{|2|} \frac{\phi(2^{j-1}x, 0)}{|2|^{3(j-1)}} : i \leq j < n + i \right\} \cup \left\{ \frac{\phi(2^{j-1}x, 2^{j-1}x)}{|2|^{3(j-1)}} : i \leq j < n + i \right\} \right\} = 0$$

exists for all  $x \in X$ , then  $C$  is the unique cubic mapping satisfying (2.17).

*Proof.* Putting  $x = y = 0$  in (2.16), we have

$$\|f(0)\| \leq \frac{1}{|2|^2} \phi(0, 0)$$

and since  $1 \leq \frac{1}{|2|}$ , we get

$$\|f(0)\| \leq \frac{1}{|2|^3} \phi(0, 0) \leq \frac{1}{|2|^{3n}} \phi(0, 0)$$

for all  $n \in \mathbb{N}$ . By (2.14),  $f(0) = 0$ .

Putting  $y = 0$  in (2.16), we have

$$(2.19) \quad \|f(3x) - 3f(2x) - 3f(x)\| \leq \frac{1}{|2|} \phi(x, 0)$$

for all  $x \in X$ . Putting  $y = x$  in (2.16), we have

$$(2.20) \quad \|f(4x) - f(3x) - 5f(2x) + 3f(x)\| \leq \phi(x, x)$$

for all  $x \in X$ . By (2.19) and (2.20), we get

$$(2.21) \quad \|f(4x) - 8f(2x)\| \leq \max \left\{ \frac{1}{|2|} \phi(x, 0), \phi(x, x) \right\}$$

for all  $x \in X$ . Replacing  $x$  by  $2^{n-1}x$  and dividing by  $|2|^{3(n+1)}$  in (2.21), we get

$$(2.22) \quad \begin{aligned} & \left\| \frac{f(2^{n+1}x)}{2^{3(n+1)}} - \frac{f(2^n x)}{2^{3n}} \right\| \\ & \leq \frac{1}{|2|^6} \max \left\{ \frac{1}{|2|} \frac{\phi(2^{n-1}x, 0)}{|2|^{3(n-1)}}, \frac{\phi(2^{n-1}x, 2^{n-1}x)}{|2|^{3(n-1)}} \right\} \end{aligned}$$

for all  $x \in X$ . By (2.14) and (2.22), we get  $\left\{ \frac{f(2^n x)}{2^{3n}} \right\}$  is Cauchy sequence. Since  $Y$  is complete, we conclude that  $\left\{ \frac{f(2^n x)}{2^{3n}} \right\}$  is convergent. Set

$$C(x) := \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^{3n}}.$$

Using induction one can show that

$$(2.23) \quad \begin{aligned} & \left\| \frac{f(2^n x)}{2^{3n}} - f(x) \right\| \leq \frac{1}{|2|^6} \max \left\{ \left\{ \frac{1}{|2|} \frac{\phi(2^{j-1}x, 0)}{|2|^{3(j-1)}} : 0 \leq j < n \right\} \cup \right. \\ & \left. \left\{ \frac{\phi(2^{j-1}x, 2^{j-1}x)}{|2|^{3(j-1)}} : 0 \leq j < n \right\} \right\} \end{aligned}$$

for all  $n \in \mathbb{N}$  and all  $x \in X$ . By taking  $n$  to infinity in (2.23) and by (2.15), we obtain (2.17). Replacing  $x$  and  $y$  by  $2^n x$  and  $2^n y$ , respectively, and dividing by  $|2|^{3n}$  in (2.16) and taking the limit as  $n \rightarrow \infty$ , by (2.14), we get

$$C(3x+y) + C(3x-y) = C(x+2y) + 2C(x-y) + 6C(2x) + 3C(x) - 6C(y)$$

for all  $x, y \in X$ . Therefore the mapping  $C : X \rightarrow Y$  satisfies (1.4) and so by Theorem 2.1,  $C$  is cubic.

Suppose that (2.18) holds. If  $C'$  is another cubic mapping satisfying (2.17), then by (2.18),

$$\begin{aligned} & \|C(x) - C'(x)\| = \lim_{i \rightarrow \infty} \frac{1}{|2|^{3i}} \|C(2^i x) - C'(2^i x)\| \\ & \leq \lim_{i \rightarrow \infty} \frac{1}{|2|^{3i}} \max \{ \|C(2^i x) - f(2^i x)\|, \|f(2^i x) - C'(2^i x)\| \} \\ & \leq \frac{1}{|2|^6} \lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} \max \left\{ \left\{ \frac{1}{|2|} \frac{\phi(2^{j-1}x, 0)}{|2|^{3(j-1)}} : i \leq j < n+i \right\} \cup \right. \\ & \quad \left. \left\{ \frac{\phi(2^{j-1}x, 2^{j-1}x)}{|2|^{3(j-1)}} : i \leq j < n+i \right\} \right\} = 0 \end{aligned}$$

for all  $x \in X$  and so  $C = C'$ . □

From Theorem 2.2, we obtain the following corollary concerning the stability of (1.4).

**COROLLARY 2.3.** *Let  $\alpha_i : [0, \infty) \rightarrow [0, \infty)$  ( $i = 1, 2, 3$ ) be mappings satisfying*

- (i)  $\alpha_i(|2|) \neq 0$ ,
- (ii)  $\alpha_i(|2|t) \leq \alpha_i(|2|)\alpha_i(t)$  for all  $t \geq 0$ , and
- (iii)  $\alpha_1(|2|) < |2|^{\frac{3}{2}}$ ,  $\alpha_2(|2|) < |2|^3$ , and  $\alpha_3(|2|) < |2|^3$ .

Let  $f : X \rightarrow Y$  be a mapping such that

$$\|Df(x, y)\| \leq \delta[\alpha_1(\|x\|)\alpha_1(\|y\|) + \alpha_2(\|x\|) + \alpha_3(\|y\|)]$$

for all  $x, y \in X$  and some  $\delta > 0$ . Suppose that  $|2| < 1$ . Then there exists a unique cubic mapping  $C : X \rightarrow Y$  such that

$$\|C(x) - f(x)\| \leq \frac{1}{|2|^6} \tilde{\phi}(x)$$

for all  $x \in X$ , where

$$\tilde{\phi}(x) = \delta|2|^2 \max \left\{ \frac{\alpha_2(\|x\|)}{\alpha_2(|2|)}, |2| \left[ \left( \frac{\alpha_1(\|x\|)}{\alpha_1(|2|)} \right)^2 + \frac{\alpha_2(\|x\|)}{\alpha_2(|2|)} + \frac{\alpha_3(\|x\|)}{\alpha_3(|2|)} \right] \right\}.$$

*Proof.* Let  $\phi(x, y) = \delta[\alpha_1(\|x\|)\alpha_1(\|y\|) + \alpha_2(\|x\|) + \alpha_3(\|y\|)]$ . Then for any  $n \in \mathbb{N}$

$$\begin{aligned} & \frac{\phi(2^n x, 2^n y)}{|2|^{3n}} \\ &= \frac{\delta}{|2|^{3n}} \left[ \alpha_1(|2|^n \|x\|)\alpha_1(|2|^n \|y\|) + \alpha_2(|2|^n \|x\|) + \alpha_3(|2|^n \|y\|) \right] \\ &\leq \delta \left[ \left( \frac{\alpha_1(|2|)}{|2|^3} \right)^n \alpha_1(\|x\|)\alpha_1(\|y\|) + \left( \frac{\alpha_2(|2|)}{|2|^3} \right)^n \alpha_2(\|x\|) \right. \\ &\quad \left. + \left( \frac{\alpha_3(|2|)}{|2|^3} \right)^n \alpha_3(\|y\|) \right] \end{aligned}$$

for all  $x, y \in X$ . By (iii), we have

$$\lim_{n \rightarrow \infty} \frac{\phi(2^n x, 2^n y)}{|2|^{3n}} = 0$$

for all  $x, y \in X$ . Hence  $\phi$  satisfies (2.14) in Theorem 2.2.

Let  $x \in X$  and  $j \in \mathbb{N} \cup \{0\}$ . Then

$$\frac{1}{|2|} \frac{\phi(2^{j-1}x, 0)}{|2|^{3(j-1)}} \leq \frac{\delta}{|2|} \left( \frac{\alpha_2(|2|)}{|2|^3} \right)^{j-1} \alpha_2(\|x\|)$$

and

$$\frac{\phi(2^{j-1}x, 2^{j-1}x)}{|2|^{3(j-1)}} \leq \delta \left[ \left( \frac{(\alpha_1(|2|))^2}{|2|^3} \right)^{j-1} (\alpha_1(\|x\|))^2 + \left( \frac{\alpha_2(|2|)}{|2|^3} \right)^{j-1} \alpha_2(\|x\|) + \left( \frac{\alpha_3(|2|)}{|2|^3} \right)^{j-1} \alpha_3(\|x\|) \right]$$

for all  $x \in X$ . By (iii), we obtain

$$\lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} \max \left\{ \left\{ \frac{1}{|2|} \frac{\phi(2^{j-1}x, 0)}{|2|^{3(j-1)}} : i \leq j < n + i \right\} \cup \left\{ \frac{\phi(2^{j-1}x, 2^{j-1}x)}{|2|^{3(j-1)}} : i \leq j < n + i \right\} \right\} = 0$$

for all  $x \in X$  and so  $\phi$  satisfies (2.18) in Theorem 2.2. Hence by Theorem 2.2, we have the result.  $\square$

EXAMPLE 2.4. Let  $\delta > 0$  and  $p$  be a real number with  $p > \frac{3}{2}$ . Suppose that  $|2| < 1$ . Let  $f : X \rightarrow Y$  is a mapping satisfying

$$\|Df(x, y)\| \leq \delta(\|x\|^p \|y\|^p + \|x\|^{2p} + \|y\|^{2p})$$

for all  $x, y \in X$ . Then there exists a unique cubic mapping  $C : X \rightarrow Y$  satisfying (1.4) such that

$$\|C(x) - f(x)\| \leq \delta |2|^{-2(p+2)} \max\{1, 3|2|\} \|x\|^{2p}$$

for all  $x \in X$ .

We have the following result which is analogous Theorem 2.2 for the functional equation (1.4).

THEOREM 2.5. Let  $\phi : X^2 \rightarrow [0, \infty)$  be a mapping such that

$$\lim_{n \rightarrow \infty} |2|^{3n} \phi\left(\frac{x}{2^n}, \frac{y}{2^n}\right) = 0$$

for all  $x, y \in X$  and for each  $x \in X$ , and let for each  $x \in X$ , the following limit

$$\lim_{n \rightarrow \infty} \max \left\{ \left\{ \frac{|2|^{3(j+2)}}{|2|} \phi\left(\frac{x}{2^{j+2}}, 0\right) : 0 \leq j < n \right\} \cup \left\{ |2|^{3(j+2)} \phi\left(\frac{x}{2^{j+2}}, \frac{x}{2^{j+2}}\right) : 0 \leq j < n \right\} \right\}$$

denoted by  $\phi_1(x)$ , exist. Suppose that  $f : X \rightarrow Y$  is a mapping satisfying  $f(0) = 0$  and

$$\|Df(x, y)\| \leq \phi(x, y)$$



for all  $x, y \in X$ . Then there exists a cubic mapping  $C : X \rightarrow Y$  satisfying (1.4) such that

$$(2.24) \quad \|C(x) - f(x)\| \leq \frac{1}{|2|^6} \phi_1(x)$$

for all  $x \in X$ . In addition, if the limit

$$\lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} \max \left\{ \left\{ \frac{|2|^{3(j+2)}}{|2|} \phi \left( \frac{x}{2^{j+2}}, 0 \right) : i \leq j < n + i \right\} \cup \left\{ |2|^{3(j+2)} \phi \left( \frac{x}{2^{j+2}}, \frac{x}{2^{j+2}} \right) : i \leq j < n + i \right\} \right\} = 0,$$

then  $C$  is the unique cubic mapping satisfying (2.24).

The following corollary is an immediate consequence of Theorem 2.5.

**COROLLARY 2.6.** Let  $\alpha_i : [0, \infty) \rightarrow [0, \infty)$  ( $i = 1, 2, 3$ ) be mappings satisfying

- (i)  $\alpha_i\left(\frac{1}{|2|}\right) \neq 0$ ,
- (ii)  $\alpha_i\left(\frac{t}{|2|}\right) \leq \alpha_i\left(\frac{1}{|2|}\right)\alpha_i(t)$  for all  $t \geq 0$ , and
- (iii)  $\alpha_1\left(\frac{1}{|2|}\right) < \frac{1}{|2|^{\frac{3}{2}}}$ ,  $\alpha_2\left(\frac{1}{|2|}\right) < \frac{1}{|2|^3}$ , and  $\alpha_3\left(\frac{1}{|2|}\right) < \frac{1}{|2|^3}$ .

Let  $f : X \rightarrow Y$  be a mapping such that  $f(0) = 0$  and

$$\|Df(x, y)\| \leq \delta[\alpha_1(\|x\|)\alpha_1(\|y\|) + \alpha_2(\|x\|) + \alpha_3(\|y\|)]$$

for all  $x, y \in X$  and some  $\delta > 0$ . Then there exists a unique cubic mapping  $C : X \rightarrow Y$  such that

$$\|C(x) - f(x)\| \leq \frac{1}{|2|^6} \phi_1(x)$$

for all  $x \in X$ , where

$$\phi_1(x) = \delta|2|^6 \max \left\{ \frac{1}{|2|} \left( \alpha_2\left(\frac{1}{|2|}\right) \right)^2 \alpha_2(\|x\|), \left( \alpha_1\left(\frac{1}{|2|}\right) \right)^4 (\alpha_1(\|x\|))^2 + \left( \alpha_2\left(\frac{1}{|2|}\right) \right)^2 \alpha_2(\|x\|) + \left( \alpha_3\left(\frac{1}{|2|}\right) \right)^2 \alpha_3(\|x\|) \right\}.$$

**EXAMPLE 2.7.** Let  $\delta > 0$  and  $p$  be a real number with  $p < \frac{3}{2}$ . Suppose that  $|2| < 1$ . Let  $f : X \rightarrow Y$  is a mapping satisfying  $f(0) = 0$  and

$$\|Df(x, y)\| \leq \delta(\|x\|^p \|y\|^p + \|x\|^{2p} + \|y\|^{2p})$$

for all  $x, y \in X$ . Then there exists a unique cubic mapping  $C : X \rightarrow Y$  satisfying (1.4) such that

$$\|C(x) - f(x)\| \leq \delta |2|^{-(4p+1)} \max\{1, 3|2|\} \|x\|^{2p}$$

for all  $x \in X$ .

### References

- [1] T. Aoki, *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan **2** (1950), 64-66.
- [2] S. Czerwik, *Functional equations and Inequalities in several variables*, World Scientific, New Jersey, London, 2002.
- [3] M. E. Gordji and M. B. Savadkouhi *Stability of cubic and quartic functional equations in non-Archimedean spaces*, Acta Appl. Math. (2010), 1321-1329.
- [4] M. E. Gordji and M. B. Savadkouhi *Stability of a mixed type cubic-quartic functional equation in non-Archimedean spaces*, Applied Mathematics Letter **23** (2010), 1198-1202.
- [5] G. L. Forti, *Hyers-Ulam stability of functional equations in several variables*, Aequationes Math. **50** (1995), 143-190.
- [6] D. H. Hyers, *On the stability of linear functional equation*, Proc. Natl. Acad. Sci. USA **27** (1941), 222-224.
- [7] D. H. Hyers, G. Isac, and T. M. Rassias, *Stability of functional equations in several variables*, Birkhäuser, Boston, 1998.
- [8] D. H. Hyers and T. M. Rassias, *Approximate homomorphisms*, Aequationes Math. **44** (1992), 125-153.
- [9] K.-W. Jun and H.-M. Kim *The generalized Hyers-Ulam-Rassias stability of a cubic functional equation*, J. Math. Anal. Appl. **274** (2002), 867-878.
- [10] K.-W. Jun and H.-M. Kim *On the Hyers-Ulam-Rassias stability of a general cubic functional equation*, Math. Inequal. Appl. **6** (2003), 289-302.
- [11] M. Sal Moslehian and G. Sadeghi, *Stability of two types of cubic functional equations in non-Archimedean spaces*, Real Analysis Exchange **33** (2007), no. 2, 375-384.
- [12] K. H. Park and Y. S. Jung, *Stability of a cubic functional equation on groups*, Bull. Korean Math. Soc. **41** (2004), 347-357.
- [13] J. M. Rassias, *Solution of the Ulam stability problem for cubic mappings*, Glasnik Matematički **36** (2001), 63-72.
- [14] F. Skof, *Approssimazione di funzioni  $\delta$ -quadratic su dominio ristretto*, Atti. Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. **118** (1984), 58-70.
- [15] S. M. Ulam, *A collection of mathematical problems*, Interscience Publ., New York, 1960.

Stability for a cubic functional equations in non-Archimedean normed spaces 363

\*

Department of Mathematics Education  
Dankook University  
Yongin 448-701, Republic of Korea  
*E-mail:* kci206@hanmail.net

\*\*

Department of Mathematics  
Soongsil University  
Seoul 156-743, Republic of Korea  
*E-mail:* seashin@hanmail.net