

## Research Report

# Development of a Single-nucleotide Polymorphism Marker for the *Sw-5b* Gene Conferring Disease Resistance to *Tomato spotted wilt virus* in Tomato

Hyung Jin Lee<sup>1</sup>, Boyoung Kim<sup>1</sup>, Chungyun Bae<sup>1</sup>, Won-Hee Kang<sup>2</sup>, Byoung-Cheorl Kang<sup>2</sup>, Inhwa Yeam<sup>3</sup>, and Chang-Sik Oh<sup>1\*</sup>

<sup>1</sup>Department of Horticultural Biotechnology and Institute of Life Science & Resources, College of Life Science, Kyung Hee University, Yongin 446-701, Korea

<sup>2</sup>Department of Plant Science, Seoul National University, Seoul 151-742, Korea

<sup>3</sup>Horticulture and Breeding Major, School of Bioresource Sciences, Andong National University, Andong 760-749, Korea

**Abstract:** *Tomato spotted wilt virus* (TSWV) causes one of the most destructive viral diseases that threatens global tomato production. *Sw-5b* was reported as the resistance gene effective against TSWV. The objective of this research was to develop a single-nucleotide polymorphism (SNP) marker to distinguish tomato cultivars resistant to TSWV from susceptible cultivars for marker-assisted breeding. First, we determined genotypes for TSWV resistance in 32 commercial tomato cultivars using the previously reported *Sw-5b* gene-based marker. Then, DNA sequences of *Sw-5b* alleles in tomato cultivars showing resistant or susceptible genotypes were analyzed; a single SNP was found to distinguish tomato cultivars resistant to TSWV from susceptible cultivars. Based on the confirmed SNP, a SNP primer pair was designed. Using this new SNP sequence and high-resolution melting analysis, the same 32 tomato cultivars were screened. The results were perfectly correlated with those from screening with the *Sw-5b* gene-based marker. These results indicate that the SNP marker developed in this study will be useful for better tracking of resistance to TSWV in tomato breeding.

**Additional key words:** Genotypes, HRM, marker-assisted selection, SNP marker, TSWV

## Introduction

*Tomato spotted wilt virus* (TSWV) causes a very destructive viral disease that threatens worldwide tomato production (*Solanum lycopersicum* L. syn *Lycopersicon esculentum* Mill.) (Saidi and Warade, 2008). TSWV is a virus belonging to the genus of negative RNA virus *Tospovirus* within the family *Bunyaviridae*. Disease symptoms caused by TSWV include bronzing, curling, necrotic spots and streaks, and stunting of stems and leaves (German et al., 1992). This virus is effectively transmitted by thrips; adult thrips can transmit viruses to a new, healthy plant only 30 minutes after feeding on viruses from infected plants. Prevention of TSWV in crop plants is difficult due to the wide range of plant

hosts. *Tomato chlorotic spot virus* (TCSV) and *Groundnut ringspot virus* (GRSV), which both belong to *Tospovirus* and are closely related to TSWV, also cause a decrease in the total production of tomato crops (Brommonschenkel et al., 2000).

The use of resistance genes is a very efficient way to reduce disease. Resistance sources to *Tospoviruses* have been found in diverse cultivars of *Solanum lycopersicum* (Saidi, 2008). Some resistance genes for TSWV have also been identified in accessions of *S. habrochaites* (previously *L. hirsutum* PI127826 and LA1353) and in *S. habrochaites* var. *glabratum* (PI134417 and LA1223). To date, eight resistance genes for TSWV, *Sw1a*, *Sw1b*, *sw2*, *sw3*, *sw4*, *Sw-5b*, *Sw-6*, and *Sw-7*, have been reported (Finlay, 1953;

\*Corresponding author: co35@khu.ac.kr

※ Received 24 February 2015; Revised 1 June 2015; Accepted 12 June 2015. This work was supported by the Golden Seed Project (Center for Horticultural Seed Development, No. 213003-04-3-SBG20), Ministry of Agriculture, Food and Rural Affairs (MAFRA), Ministry of Oceans and Fisheries (MOF), Rural Development Administration (RDA), and Korea Forest Service (KFS) and also by Technology Commercialization Support Program (No. 113044-3) of Korea Institute of Planning & Evaluation for Technology in Food, Agriculture Forestry & Fisheries.

© 2015 Korean Society for Horticultural Science

Price et al., 2007; Saidi, 2008). Among them, *Sw-5b* has been widely deployed in tomato breeding because of its durable and reliable resistance to multiple *Tospoviruses* (Stevens et al., 1991). The genetic location of *Sw-5b* gene has been mapped between restriction fragment length polymorphism (RFLP) markers CT71 and CT220 near a telomeric region of the long arm of chromosome 9 in tomato (Stevens et al., 1995). This gene was cloned, and the protein encoded by this gene consists of the coiled-coil (CC) domain, the nucleotide binding-adaptor shared by APAF-1, R proteins and a CED-4 (NB-ARC) and a leucine-rich repeat (LRR) domain (Spasova et al., 2001). This gene is similar to two known resistance genes in tomato: the nematode resistance gene *Mi* (51% identity at the amino acid level) and the *Prf* gene resistant to *Pseudomonas syringae* (43% identity at the amino acid level) (Spasova et al., 2001).

Maker-assisted selection (MAS) has been widely applied for the breeding of disease-resistant cultivars in many crops, including tomato, by utilizing genetic markers linked to corresponding resistance genes (Foolad, 2007; Foolad and Sharma, 2004). Among diverse cultivars, single nucleotide polymorphisms (SNPs) in certain genes responsible for important phenotypes, such as disease resistance, have been widely used in genome mapping, association studies, diversity analysis, and tagging of economic important genes in plant genomics (Giancola et al., 2006; Rafalski, 2002). Most SNPs have been discovered and verified in tomato (Labate and Baldo, 2005; Yang et al., 2004) and successfully used in selection of various bacterial disease resistances (Yang et al., 2005). Molecular markers, such as RFLP and cleaved amplified polymorphic sequence (CAPS) markers, are based on the presence of SNPs or insertion/deletion (INDEL) mutations (Nasu et al., 2002). SNPs can also be used directly for developing SNP markers. High-resolution melting (HRM) analysis or SNP chips like Fluidigm have been recently developed and are currently being widely used (Wu et al., 2008).

Molecular markers linked to TSWV resistance have also been developed and reported in tomato (Foolad and Panthee, 2012). In particular, for TSWV resistance conferred by the *Sw-5b* gene, diverse molecular markers such as RAPD (random amplified polymorphic DNA), RFLP, SCAR (sequence characterized amplified regions), CAPS, and INDEL markers have been reported (Dianese et al., 2010; Garland et al., 2005; Langella et al., 2004; Roselló et al., 2001; Śmiech et al., 2000; Stevens et al., 1995). Since the successful cloning of the *Sw-5b* gene, a gene-based marker and two SNPs specific to the *Sw-5b* allele for TSWV resistance have been

reported (Shi et al., 2011). However, SNP analysis between resistant and susceptible alleles of the *Sw-5b* gene has not yet been intensively examined.

In this study, we first detected SNPs between resistant and susceptible alleles of *Sw-5b* genes in 32 commercial tomato cultivars. Then, SNP markers for HRM using this SNP information were developed.

## Materials and Methods

### Plant materials

Thirty-two commercial tomato cultivars were used in this study (Table 1): thirteen cultivars from Nongwoo Bio (Suwon, Korea), eight cultivars from Syngenta (Basel, Switzerland), five cultivars from Monsanto Korea (St. Louis, Missouri, United States), four cultivars from Asia Seed (Seoul, Korea), and one cultivar each from Takii (Kyoto, Japan) and Sakata (Sakata, Japan). Moneymaker and M82 were included as TSWV-susceptible cultivars. Only two cultivars, 'TY sense Q' and 'Titichal' among the 32 commercial cultivars used in this study were described by the company as being resistant to TSWV. These tomato cultivars were grown in a growth chamber (14 h light/ 10 h dark, 26-28°C, 63.03  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  light intensity) for 4-5 weeks for genomic DNA isolation.

### Genomic DNA extraction

Genomic DNA (gDNA) was extracted from young leaves using the cetyl-trimethyl-ammonium bromide (CTAB) method with minor modifications (Murray and Thompson, 1980). Briefly, fresh, young leaves (0.2 g) were ground in a 2 mL microfuge tube with 300  $\mu\text{L}$  extraction buffer [(2% w/v) CTAB, 1.4 M NaCl, 20 mM EDTA, 100 mM Tris-HCl pH 8.0, 1% PVP, 0.1% sodium bisulfite, and 0.2% (v/v) 2-mercaptoethanol] for gDNA extraction. Extracted gDNA was quantified using a Nanodrop 2000/UV-Vis spectrophotometer (Thermo Scientific, Wilmington, DE, USA) and diluted to 10  $\text{ng}\cdot\mu\text{L}^{-1}$  for polymerase chain reaction (PCR).

### PCR amplification and DNA sequencing

For the primer pairs Sw5b-f1/r1 and Sw5b-f2/r2 (Table 2), PCR amplification was conducted using procedures described by Shi et al. (2011). PCR fragments were separated by gel electrophoresis with 1% agarose gel in 0.5 X TAE buffer and visualized under UV light using Dual UV transilluminator after staining with EcoDye™ DNA Staining Solution (Biofact Co., Daejeon, Korea). The PCR fragments amplified from the primer pairs Sw5-f1/r1 were sequenced in Macrogen

**Table 1.** Genotype evaluation for TSWV resistance in 32 commercial tomato cultivars with the designated genetic markers.

Number	Cultivar name	Source	Genotype per markers			
			Gene-based marker <sup>x</sup>	<i>Sw-5b</i> -SNP <sup>y</sup>	SNP1 <sup>x</sup>	SNP2 <sup>x</sup>
1	TY sense Q*		R	R	R	S
2	Savera		R	R	V <sup>z</sup>	S
3	TY tiny		R	R	R	S
4	Titichal*		R	R	R	S
5	Tiara		S	S	S	S
6	Redpang		S	S	S	S
7	Prime alexander.	Nongwoo Bio	S	S	S	S
8	Minimaru		S	S	S	S
9	Gold minichal		S	S	S	S
10	TY altorang		S	S	S	S
11	Alexander delux		S	S	S	S
12	Minichal		S	S	S	S
13	Pinktop		S	S	S	S
14	Daylose		R	R	V	R
15	Dafnis		S	S	V	S
16	Medison		S	S	S	S
17	TP-7 plus		S	S	S	S
18	Mamirio	Syngenta	S	S	S	S
19	Dunne		S	S	S	S
20	Gayachal plus		S	S	S	S
21	Rapsody		S	S	S	S
22	Unicon		S	S	S	S
23	Olle TY		S	S	S	S
24	Stix TY	MonsantoKorea	S	S	S	S
25	Bacchus		S	S	S	S
26	Rapito		S	S	S	S
27	Shinheuksu		R	R	R	S
28	Cadillak		S	S	S	S
29	Miniheuksu	Asia Seed	S	S	S	S
30	TY250		S	S	S	S
31	Dotaerang TY winner	Takii	S	S	S	S
32	Oyama	Sakata	S	S	S	S
33	Moneymaker		S	S	S	S
34	M82		S	S	S	S

\*These cultivars were described by the company as being resistant to TSWV.

<sup>z</sup>Variable HRM result.

<sup>y</sup>Developed in this study.

<sup>x</sup>Obtained from Shi et al. (2011).

R, resistant genotype; S, susceptible genotype.

**Table 2.** PCR primers used in this study.

Primer name	Primer (5'→3')	T <sub>m</sub> (°C)	Product size (bp)	Reference or source	
Sw5b-f1	AACCACTAGGGGCAGTCCTT	63.7	662	Shi et al., 2011	
Sw5b-r1	CTCACTATGTGGCTGCTCCA	64.1			
Sw5b-f2	CGGAACCTGTAACCTTGACTG	59.8	541		
Sw5b-r2	GAGCTCTCATCCATTTTCCG	59.8			
SNP1 F	GAGAGAGTTATTACAAGAT	48.7	93		
SNP1 R	TTTCTCCTCAACATGTCGGC	58.4			
SNP2 F	GGTATGGGATGACTTAAGGC	58.4	85		
SNP2 R	ACCACTTCTTCAAGTCGAG	58.4			
Sw5b-SNP-F	TTATTGTTTCTCGCTTTGATGTTTCG	60.9	100		This study
Sw5b-SNP-R	GAACCTGTAACCTTGACTGAAAATATC	61.6			

Inc. (Seoul, Korea). DNA sequences of the PCR products amplified with the primer pair Sw5-f1/r1 were aligned with ClustalW2 database from EMBL-EBI (<https://www.ebi.ac.uk/Tools/msa/clustalw2/>).


#### High-resolution melting (HRM) analysis

For HRM analysis, the Sw5b-SNP primer pair and the SNP1 and SNP2 primer pairs reported in Shi et al. (2011) (Table 2) were designed to amplify the DNA region carrying the target SNP. PCR was carried out in 20 µL reaction mixtures containing 10X *h-Taq* Reaction Buffer (25 mM MgCl<sub>2</sub> mix), 10 mM of each dNTP Mix, 10 pmol of each primer, 2.5 U/µL DNA polymerase, 20X EvaGreen™ (Solgent, Daejeon, Korea), 5X Band Doctor™ and 10ng of gDNA, using a CFX Connect™ Real-Time PCR Detection System (BIO-RAD, Hercules, USA). Cycling conditions were 95°C for 15 min, followed by 40 cycles of 95°C for 20s, 60°C for 15 s, and 72°C for 30 s, and HRM was analyzed for every increment of 0.2°C between 65 and 95°C.

## Results and Discussion

#### Evaluation of 32 commercial tomato cultivars for TSWV resistance using the *Sw-5b* gene-based marker

To examine whether commercial tomato cultivars currently available in Korea carry resistance to TSWV, 32 cultivars were collected from the seed companies listed in Table 1. The dominant PCR marker specific to *Sw-5b* with the primer pair Sw5b-f2/Sw5b-r2 (Table 2) was previously reported (Shi et al., 2011). Collected tomato cultivars were screened by PCR with this primer pair, including the TSWV-susceptible cultivars, Moneymaker and M82. PCR amplified 541-bp DNA products from only six cultivars ('TY sense




**Fig. 1.** Determination of genotypes for TSWV resistance in 32 commercial tomato cultivars using PCR with the primer pair Sw5b-f2/ Sw5b-r2. Cultivars showing the resistant genotype (1 to 6) has an obvious PCR band at 541-bp. M, 1 kb DNA ladder; 1, TY sense Q; 2, Titchal; 3, Savera; 4, Daylose; 5, TY tiny; 6, Shinheuksu; 7, Prime alexander; 8, Minimaruru; 9, Gold minichal; 10, TY altorang; 11, Alexander deluxe; 12, Tiara; 13, Minichal; 14, Pinktop; 15, Rapsody; 16, Mamirio; 17, Medison; 18, TP-7plus; 19, Dafnis; 20, Dunne; 21, Gayachal plus; 22, Dotaerang TY winner; 23, Oyama; 24, Unicon; 25, Miniheuksu; 26, Cadillac; 27, Redpang; 28, TY 250; 29, Olle TY; 30, Stix TY; 31, Bacchus; 32, Rapito; 33, Moneymaker; 34, M82.

Q', 'Titchal', 'Savera', 'TY tiny', 'Daylose' and 'Shinheuksu') of 32 cultivars (Fig. 1), indicating that only these six cultivars carry the *Sw-5b* allele for TSWV-resistance. Except for 'TY sense Q' and 'Titchal' that were already described by the provider to carry resistance to TSWV, 'Savera', 'TY tiny', 'Daylose' and 'Shinheuksu' were newly identified as

cultivars carrying resistance to TSWV in this study.

Previously, Shi et al. (2011) also screened 14 tomato accessions and 10 commercial tomato cultivars for TSWV-resistance and found that only three commercial cultivars ('Talladega', 'Sophya', and 'BHN-444') and one accession (LA3667) carried the *Sw-5b* allele for resistance to TSWV as shown in the Stevens cultivar, the original source of

the *Sw-5b* allele for resistance to TSWV. In this study, completely different commercial cultivars from what Shi et al. (2011) used were screened, and only six of them were determined to be TSWV-resistant. Both studies indicate that not many commercial cultivars in the market carry resistance to TSWV, a situation that needs to be remedied to reduce the large amount of agricultural loss.



**Fig. 2.** Identification of the single SNP in *Sw-5b* resistance gene alleles from tomato cultivars showing resistant and susceptible genotypes. (A) Location of primers and the functional SNP in *Sw-5b* gene used in this study. Numbers indicate the position of bases. (B) The part of the DNA sequences of PCR products amplified with Sw5b-SNP-F/R primers in 32 commercial tomato cultivars. The number indicates the position of the SNP from the translation start site of Sw-5b allele that was underlined and bold. \* indicates bases identical in all 32 cultivars.

### Detection of a functional SNP to distinguish resistant and susceptible *Sw-5b* alleles

To find SNPs for distinguishing between resistant and susceptible *Sw-5b* alleles, another primer pair Sw5b-f1/Sw5b-r1 (Table 2, Fig. 2A) was designed based on a previous report (Shi et al., 2011). PCR with this primer pair amplified 662-bp DNA products from both resistant and susceptible *Sw-5b* alleles in all 32 cultivars (data not shown). All PCR products were sequenced, and their DNA sequences were aligned using CluswalW2. In addition, the alleles from the resistant cultivar 'Stevens' (GenBank Accession No. AY007366) and the susceptible cultivar 'NY07-464' (GenBank accession No. FJ686042.1) were retrieved from the GenBank database and included in this alignment. Sequence alignment revealed several SNPs in this region, including G/A at 1,796-bp from the translational start site of *Sw-5b* allele (Fig. 2C). Among these SNPs, the G/A SNP at 1,796-bp was 100% correlated with genotypes for TSWV resistance. All six resistant cultivars ('TY sence Q', 'Titichal', 'Savera', 'TY tiny', 'Daylose' and 'Shinheuksu') have a G base like the AY007366 allele, while all 22 susceptible cultivars have an A base like the NY07-464 allele. This base change was a non-synonymous substitution, i.e., G to A base substitution results in change of the 599<sup>th</sup> amino acid, glutamine (Gln<sup>599</sup>) to arginine (Arg<sup>599</sup>). Among the three domains of the *Sw-5b* protein, the 599<sup>th</sup> amino acid is located in the NB-ARC domain (Fig. 2A). It is likely that, due to different features between these two amino acids, the tertiary structure of the *Sw-5b* protein changes rendering the protein non-functional.


To date, only one paper has reported SNPs specific to TSWV resistance in tomato (Shi et al., 2011). In that report, six SNPs were found in the *Sw-5b* allele, and all of them are located in the DNA region, encoding the NB-ARC domain of the *Sw-5b* protein. Of these six, only two SNPs [A/T at 31,279 position and A/G at 31,392 position in *Sw-5b* (AY007366) of 'Stevens' cultivar] were confirmed to distinguish the TSWV-resistant cultivars from the susceptible cultivars. It has not been determined if the other four SNPs (C/T at 31,415 position, G/A at 31,317 position, G/A at 31,434 position, and C/T at 31,459 position in AY007366) distinguish the TSWV-resistant cultivars from the susceptible cultivars in SNP genotyping. Interestingly, the C/T SNP at position 31,459 is exactly the same SNP confirmed in this study, both of which are the reverse-complement bases of the G/A SNP shown in Fig. 2. In our study, no other SNPs that can distinguish the TSWV-resistant cultivars from the susceptible cultivars were found. This implies that only the G/A SNP can be used for determining TSWV resistance

in commercial tomato cultivars available in Korea.

### Development of the SNP marker for TSWV-resistance

To examine whether the G/A SNP can be used as a SNP marker for selecting TSWV-resistance, the primer pair Sw5b-SNP-F/R was designed. The expected size of the PCR product was 100 bp (Table 2, Fig. 2A) and HRM analysis was carried out. In general, HRM polymorphism is determined by changes in the melting curve shape and location. HRM analysis with the primer pair Sw5b-SNP-F/R clearly distinguished two genotypes at temperatures between 75°C and 80°C (Fig. 3). As summarized in Table 1, only six cultivars ('TY sence Q', 'Titichal', 'Savera', 'TY tiny', 'Daylose' and 'Shiheuksu') were categorized as the resistant group, while all others were the susceptible group. This grouping exactly matched the results from the gene-specific PCR marker shown in Fig. 1. These results indicate that the SNP primer set can be used as a SNP marker for TSWV resistance in commercial tomato cultivars available in Korea.

To compare the accuracy between the SNP marker developed in this study and two SNP markers reported in Shi et al. (2011), HRM analysis was conducted with two SNP markers, SNP1 and SNP2, in the 32 commercial tomato cultivars listed in Table 1. As summarized in Table 1, both SNP1 and SNP2 markers were less accurate than the *Sw-5b* SNP marker developed in this study. Based on these results, G/A SNP at the 1,796-bp site from the translational start site of *Sw-5b* allele (corresponding to 31,459 position in 'Stevens' cultivar allele AY007366) is more accurate for determining TSWV resistance in commercial tomato cultivars.



**Fig. 3.** HRM analysis of PCR products amplified with Sw5b-SNP-F/R primers in 32 commercial tomato cultivars. The green and red melting curves indicate groups of tomato showing resistant (R) or susceptible (S) genotypes, respectively. Results are summarized in Table 1.

## Literature Cited

- Brommonschenkel, S. H., A. Frary, A. Frary, A., and S. D. Tanksley. 2000. The broad-spectrum *tospovirus* resistance gene *Sw-5* of tomato is a homolog of the root-knot nematode resistance gene *Mi*. *Mol. Plant-Microbe Interact.* 13:1130-1138.
- Dianese, É. C., M. E. N, de Fonseca, R. Goldbach, R. Kormelink, A. K. Inoue-Nagata, R. O. Resende, and L. S. Boiteux. 2010. Development of a locus-specific, co-dominant SCAR marker for assisted-selection of the *Sw-5* (*Tospovirus* resistance) gene cluster in a wide range of tomato accessions. *Mol. Breed.* 25:133-142.
- Finlay, K. 1953. Inheritance of spotted wilt resistance in the tomato II. Five genes controlling spotted wilt resistance in four tomato types. *Aust. J. Biol. Sci.* 6:153-163.
- Foolad, M. R. 2007. Genome mapping and molecular breeding of tomato. *Int. J. Plant Genomics* 2007:64358.
- Foolad, M. R. and D. R. Panthee. 2012. Marker-assisted selection in tomato breeding. *Crit. Rev. Plant Sci.* 31:93-123.
- Foolad, M. R. and A. Sharma. 2004. Molecular markers as selection tools in tomato breeding. *Acta Hort.* 695:225-240.
- Garland, S., M. Sharman, D. Persley, and D. McGrath. 2005. The development of an improved PCR-based marker system for *Sw-5*, an important TSWV resistance gene of tomato. *Crop Pasture Sci.* 56:285-289.
- German, T. L., D. E. Ullman, and J. W. Moyer. 1992. *Tospoviruses*: diagnosis, molecular biology, phylogeny, and vector relationships. *Annu. Rev. Phytopathol.* 30:315-348.
- Giancola, S., H. I. McKhann, A. Berard, C. Camilleri, S. Durand, S. P. Libeau, F. Roux, X. Reboud, I. G. Gut, and D. Brunel. 2006. Utilization of the three high-throughput SNP genotyping methods, the GOOD assay, Amplifluor and TaqMan, in diploid and polyploid plants. *Theor. Appl. Genet.* 112:1115-1124.
- Labate, J. A. and A. M. Baldo. 2005. Tomato SNP discovery by EST mining and resequencing. *Mol. Breed.* 16:343-349.
- Langella, R., M. Ercolano, L. Monti, L. Frusciante, and A. Barone. 2004. Molecular marker assisted transfer of resistance to TSWV in tomato elite lines. *J. Hort. Sci. Biotechnol.* 79:806-810.
- Murray, M. and W. F. Thompson. 1980. Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Res.* 8:4321-4326.
- Nasu, S., J. Suzuki, R. Ohta, K. Hasegawa, R. Yui, N. Kitazawa, L. Monna, and Y. Minobe. 2002. Search for and analysis of single nucleotide polymorphisms (SNPs) in rice (*Oryza sativa*, *Oryza rufipogon*) and establishment of SNP markers. *DNA Res.* 9:163-171.
- Price, D. L., F. D. Memmott, J. W. Scott, S. M. Olson, and M. R. Stevens. 2007. Identification of molecular markers linked to a new Tomato spotted wilt virus resistance source in tomato. *Tomato Genet. Coop.* 57:35-36.
- Rafalski, A. 2002. Applications of single nucleotide polymorphisms in crop genetics. *Curr. Opin. Plant Biol.* 5:94-100.
- Roselló, S., B. Ricarte, M. J. Diez, and F. Nuez. 2001. Resistance to Tomato spotted wilt virus introgressed from *Lycopersicon peruvianum* in line UPV 1 may be allelic to *Sw-5* and can be used to enhance the resistance of hybrids cultivars. *Euphytica* 119:357-367.
- Saidi, M. and S. D. Warade. 2008. Tomato breeding for resistance to Tomato spotted wilt virus (TSWV): an overview of conventional and molecular approaches. *Czech J. Genet. Plant Breed.* 44:83-92.
- Shi, A., R. Vierling, R. Grazzini, P. Chen, H. Caton, and D. Panthee. 2011. Identification of molecular markers for *Sw-5* gene of tomato spotted wilt virus resistance. *Am. J. Biotechnol. Mol. Sci.* 1:8-16.
- Śmiech, M., Z. Rusinowski, S. Malepszy, and K. Niemirowicz-Szczytt. 2000. New RAPD markers of Tomato spotted wilt virus (TSWV) resistance in *Lycopersicon esculentum* Mill. *Acta Physiol. Plantarum* 22:299-303.
- Spasova, M. I., T. W. Prins, R. T. Folkertsma, R. M. Klein-Lankhorst, J. Hille, R. W. Goldbach, and M. Prins. 2001. The tomato gene *Sw5* is a member of the coiled coil, nucleotide binding, leucine-rich repeat class of plant resistance genes and confers resistance to TSWV in tobacco. *Mol. Breed.* 7:151-161.
- Stevens, M., E. Lamb, and D. Rhoads. 1995. Mapping the *Sw-5* locus for tomato spotted wilt virus resistance in tomatoes using RAPD and RFLP analyses. *Theor. Appl. Genet.* 90:451-456.
- Stevens, M., S. Scott, and R. Gergerich. 1991. Inheritance of a gene for resistance to Tomato spotted wilt virus (TSWV) from *Lycopersicon peruvianum* Mill. *Euphytica* 59:9-17.
- Wu, S.-B., M. G. Wirthensohn, P. Hunt, J. P. Gibson, J., and M. Sedgley. 2008. High resolution melting analysis of almond SNPs derived from ESTs. *Theor. Appl. Genet.* 118, 1-14.
- Yang, W., X. Bai, E. Kabelka, C. Eaton, S. Kamoun, E. van der Knaap, and D. Francis. 2004. Discovery of single nucleotide polymorphisms in *Lycopersicon esculentum* by computer aided analysis of expressed sequence tags. *Mol. Breed.* 14:21-34.
- Yang, W., S. Miller, J. Scott, J. Jones, and D. Francis. 2005. Mining tomato genome sequence databases for molecular markers: application to bacterial resistance and marker assisted selection. *Acta Hort.* 695:241-250.