http://e-nrp.org

# Dietary intake of fats and fatty acids in the Korean population: Korea National Health and Nutrition Examination Survey, 2013

Yeji Baek<sup>1\*</sup>, Ji-Yun Hwang<sup>2\*</sup>, Kirang Kim<sup>3</sup>, Hyun-Kyung Moon<sup>4</sup>, Sanghui Kweon<sup>1</sup>, Jieun Yang<sup>1</sup>, Kyungwon Oh<sup>1§</sup> and Jae Eun Shim<sup>5§</sup>

<sup>1</sup>Division of Health and Nutrition Survey, Korea Centers for Disease Control and Prevention, Osong Health Technology Administration Complex, 187,

Osongsaengmyeong2-ro, Cheongju-si, Chungbuk 28159, Korea

<sup>2</sup>Nutrition Education Major, Graduate School of Education, Sangmyung University, Seoul 03016, Korea

<sup>3</sup>Department of Food Science and Nutrition, Dankook University, Chungnam 31116, Korea

<sup>4</sup>Department of Food Science and Nutrition, Dankook University, Gyeonggi 16890, Korea

<sup>5</sup>Department of Food and nutrition, Daejeon University, 62, Daehak-ro, Dong-gu, Daejeon 34520, Korea

**BACKGROUND/OBJECTIVES:** The aim of this study was to estimate average total fat and fatty acid intakes as well as identify major food sources using data from the Korea National Health and Nutrition Examination Survey (KNHANES) VI-1 (2013). **SUBJECTS/METHODS:** Total fat and fatty acid intakes were estimated using 24-hour dietary recall data on 7,048 participants aged  $\geq$  3 years from the KNHANES VI-1 (2013). Data included total fat, saturated fatty acid (SFA), monounsaturated fatty acid (MUFA), polyunsaturated fatty acid (PUFA), n-3 fatty acid (n-3 FA), and n-6 fatty acid (n-6 FA) levels. Population means and standard errors of the mean were weighted in order to produce national estimates and separated based on sex, age, income, as well as residential region. Major food sources of fat, SFA, MUFA, PUFA, n-3 FA, and n-6 FA were identified based on mean consumption amounts of fat and fatty acids in each food.

**RESULTS:** The mean intake of total fat was 48.0 g while mean intakes of SFA, MUFA, PUFA, n-3 FA, and n-6 FA were 14.4 g, 15.3 g, 11.6 g, 1.6 g, and 10.1 g, respectively. Intakes of MUFA and SFA were each higher than that of PUFA in all age groups. Pork was the major source of total fat, SFA, and MUFA, and soybean oil was the major source of PUFA. Milk and pork were major sources of SFA in subjects aged 3-11 years and  $\geq$  12 years, respectively. Perilla seed oil and soybean oil were main sources of n-3 FA in subjects aged  $\geq$  50 years and aged < 50 years, respectively.

**CONCLUSIONS:** Estimation of mean fatty acid intakes of this study using nationally represented samples of the Korean population could be useful for developing and evaluating national nutritional policies.

Nutrition Research and Practice 2015;9(6):650-657; doi:10.4162/nrp.2015.9.6.650; pISSN 1976-1457 eISSN 2005-6168

Keywords: Fatty acid, fat, food sources, KNHANES

## INTRODUCTION

The current dietary environment has changed due to industrialization and economic development, which have had significant impacts on nutrition and health. The prevalence of chronic diseases is rapidly growing worldwide, and the major proportion of deaths due to chronic diseases is attributable to cardiovascular disease [1].

Dietary factors associated with reduced risk of CVD include fruits, vegetables, fish, and foods with high contents of linoleic acid, EPA, DHA, and potassium [2]. Dietary fats have been investigated as risk factors associated with coronary heart disease. Generally, intake of saturated fatty acid (SFA) raises blood cholesterol, especially low-density lipoprotein (LDL) cholesterol, whereas individual fatty acids have different effects [3-6]. An effective way to reduce risk of coronary heart disease is substitution of unsaturated fatty acids for SFA. There is an optimum range of individual fatty acid intake according to the type of fatty acid (e.g. polyunsaturated fatty acid (PUFA), monounsaturated fatty acid (MUFA), n-3 fatty acid (n-3 FA), n-6 fatty acid (n-6 FA)) [7-10].

The World Health Organization (WHO) has set nutrient intake guidelines for populations in order to establish national and regional dietary recommendations for prevention of diet-related chronic diseases such as cardiovascular diseases [2]. Korea has also set adequate intake ranges of fat and fatty acids based on Korean Dietary Reference Intakes (i.e. acceptable macronutrient distribution ranges, AMDR), but a fatty acid database for estimation of fatty acid consumption in the Korean population is not yet available [11].

Fat consumption has gradually increased with each nationally representative nutrition survey [12]. Even though current

Received: March 25, 2015, Revised: August 12, 2015, Accepted: August 17, 2015

\* These two authors contributed to this work equally.

<sup>&</sup>lt;sup>§</sup> Corresponding Author: Jae Eun Shim, Tel. 82-42-280-2469, Fax. 82-42-280-2468, Email. jshim@dju.kr

Kyungwon Oh, Tel. 82-43-719-7460, Fax. 82-43-719-7527, Email. kwoh27@korea.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-nc/3.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Recently, a fatty acid composition table was constructed based on food lists in the Korea National Health and Nutrition Examination Survey (KNHANES) [13,14]. Population-level intakes of fatty acids can be evaluated using recently released KNHANES data. Thus, this study was conducted to produce estimates of population intake status and major sources of fatty acids, which could provide data for policy development and health promotion strategies related to dietary fats.

## SUBJECTS AND METHODS

#### Study population

Subjects were participants of KNHANES VI-1 (2013), which was a nationwide cross-sectional survey assessing the health and nutritional status of Koreans [15]. The KNHANES has been conducted every year by the Korea Centers for Disease Control and Prevention on a target population of non-institutionalized civilians in Korea with a sampling plan of multi-stage clustered probability design.

In this analysis, we included 7,048 participants aged 3 years or older who completed a 24-h dietary recall. Children aged 1 and 2 years (n = 194) were excluded due to systematic underestimation of fat and fatty acid intakes since the KNHANES did not include breast milk. Men were 43.9% of subjects and 80.1% of those lived in urban areas.

## Assessment of total fat and fatty acid intakes

Dietary intake data were obtained using the 24-h dietary recall method. Total fat intakes were calculated using a food composition table published by the Rural Development Administration of Korea [16]. Intakes of various fatty acids were calculated using a fatty acid composition table containing fatty acid content data on 2,143 foods published by national agencies from Korea, United States, and Japan. Fatty acid composition was matched according to food name with same species and same form (dried, boiled, etc.), and domestic sources were selected preferentially compared to foreign sources. If the same species or form of food was not available in the literature, imputation was performed using the calculation from the same species with different forms (dried, boiled, etc.) or by substitution with familiar species in a biosystematic grouping. The process of developing the fatty acid composition table has been reported in detail [13,14]. The major food sources of total fat and fatty acids were evaluated based on the mean amounts of fat and fatty acids of each food.

#### Statistical analysis

Intakes of total fat and fatty acids were expressed as means and standard errors by groups according to gender, age, region, and income. Ratios of n-6 FA to n-3 FA as well as PUFA to SFA were highly skewed to the right, and geometric means were calculated. Food sources of total fat and fatty acids were calculated and expressed as intake amounts and as percentage contribution to total intake (%). Taylor series method was performed to account for the complex survey design of KNHANES. All statistical analyses were performed using statistical software package SAS version 9.4.

## RESULTS

## Total fat and fatty acid intakes

Total fat and fatty acid intakes according to sociodemographic characteristics are shown in Table 1. Mean intake of total fat was 48.0 g while mean intakes of SFA, MUFA, PUFA, n-3 FA, and n-6 FA were 14.4 g, 15.3 g, 11.6 g, 1.6 g, and 10.1 g, respectively. Intakes of MUFA and SFA were each higher than that of PUFA in all age groups. Total fat and fatty acid intakes of men were higher than those of women, and total fat and individual fatty acid intakes were highest in subjects aged 19-29 years. Subjects who lived in urban areas consumed a relatively higher amount of PUFA and lower amounts of SFA and MUFA compared to residents in rural areas. Total fat and fatty acid intakes of high-income subjects were higher than those of low-income subjects. Geometric means and standard errors of PUFA/SFA and n-6 FA/n-3 FA were 0.821  $\pm$  0.002 and 6.914  $\pm$ 0.023, respectively (data not shown).

#### Food sources of total fat and fatty acids

The major source of total fat was milk (15%) in children 3-5 years of age, followed by pork (10%) and cookies and biscuits (9%) (Table 2). Pork was the highest ranked source of total fat in subjects 6 years of age and over, providing 12-18% of total fat. Pork, soybean oil, and beef were the top three sources of total fat in subjects 30 years of age and over.

The main contributor to SFA intake was milk, followed by pork in children 3-11 years of age (Table 3). In children 3-5 years of age, milk, pork, and cheese contributed 48% to SFA intake. Cookies, biscuits, snack, ice cream, instant noodles, and chocolate contributed 15-18% to SFA intake among subjects aged 3-18 years. Pork was the highest ranked source of SFA among subjects aged 12 years or older, providing 16-23% of SFA intake. The four major sources of SFA were pork, instant noodles, milk, and beef in subjects 19-49 years of age as well as pork, milk, beef, and coffee in subjects 50 years of age and over. These four sources provided almost 40% of SFA intake.

The highest ranked source of MUFA was milk in children 3-5 years of age and pork in those 6 years of age and over, which were very close to the SFA results (Table 4). The main contributors to MUFA intake were meat (pork, beef, or chicken) and vegetable oils (soybean oil, sesame oil, or rape seed oil). Meat contributed 18-35% to MUFA intake, and vegetable oils contributed 5-14% to MUFA intake. Egg and milk were also major food sources, and other sources were different depending on age; snack and bread in subjects 3-18 years of age; instant noodles, mayonnaise, and bread in subjects 19-49 years of age; instant noodles, peanuts, and almonds in subjects 50-64 years of age; and over.

Vegetable oils (soybean oil, sesame oil, or rape seed oil) were the main source of PUFA in all age groups, providing 22-32% of PUFA intake, whereas soybean oil was the highest ranked food source (Table 5). Pork and mayonnaise were also major contributors, and intakes of PUFA from pulses (soybean milk,

## Fatty acid intake in Korean

**Table 1.** Total fat and fatty acid intake<sup>1)</sup>

Table 1. Total fat ar	nd fatty acid inta	ike <sup>1)</sup>				(g/day/capita	a)
		Total fat	SFA	MUFA	PUFA	n-3 FA	n-6 FA
		$\text{Mean} \pm \text{SE}$	Mean ± SE	Mean ± SE	Mean ± SE	Mean ± SE	Mean ± SE
Total	7,048	$48.0\pm0.8$	$14.4\pm0.3$	$15.3\pm0.3$	$11.6 \pm 0.2$	$1.59\pm0.03$	$10.1 \pm 0.2$
Gender							
Men	3,097	55.7 ± 1.1	$16.7\pm0.3$	$17.9\pm0.4$	$13.3\pm0.3$	$1.80\pm0.05$	$11.6 \pm 0.2$
Women	3,951	$40.2\pm0.8$	$12.0\pm0.3$	$12.7\pm0.3$	$9.8\pm0.2$	$1.39\pm0.04$	$8.5 \pm 0.2$
Age (yrs)							
3-5	313	$41.2 \pm 3.2$	$15.0 \pm 1.2$	$13.2 \pm 1.1$	$7.9\pm0.9$	$1.00\pm0.09$	$7.0 \pm 0.8$
6-11	640	50.9 ± 1.6	$17.4 \pm 0.6$	$16.5\pm0.5$	$10.4\pm0.4$	$1.26\pm0.06$	$9.2\pm0.4$
12-18	654	$60.2 \pm 1.6$	$18.8\pm0.5$	$19.7\pm0.6$	$13.5\pm0.5$	$1.55 \pm 0.07$	$12.0\pm0.4$
19-29	670	$64.8\pm2.2$	$19.7\pm0.7$	$21.0\pm0.8$	$15.2\pm0.5$	$1.79\pm0.08$	$13.5\pm0.5$
30-49	2,023	52.6 ± 1.1	$15.4 \pm 0.4$	$16.9\pm0.4$	$12.9\pm0.3$	$1.77 \pm 0.05$	$11.3 \pm 0.2$
50-64	1,450	$37.7 \pm 0.9$	$10.5\pm0.3$	$11.8\pm0.3$	$10.0\pm0.3$	$1.67\pm0.08$	$8.4\pm0.2$
65+	1,298	$23.7\pm0.7$	$6.8\pm0.2$	$7.1 \pm 0.3$	$6.2\pm0.2$	$1.08\pm0.06$	$5.2 \pm 0.2$
Region <sup>2)</sup>							
Urban	5,642	$50.0\pm0.8$	$15.0\pm0.3$	$16.1 \pm 0.3$	$12.0\pm0.2$	$1.62\pm0.04$	$10.5 \pm 0.2$
Rural	1,406	$50.7\pm2.2$	$15.9\pm0.9$	$16.2 \pm 0.8$	$11.4 \pm 0.4$	$1.52\pm0.05$	$10.0\pm0.3$
Income <sup>2,3)</sup>							
Low	1,712	46.8 ± 1.3	$14.2 \pm 0.4$	$15.0\pm0.5$	$11.1 \pm 0.3$	$1.46\pm0.06$	$9.8\pm0.3$
Middle-low	1,791	48.0 ± 1.1	$14.6\pm0.3$	$15.4\pm0.4$	$11.5 \pm 0.3$	$1.52\pm0.05$	$10.1 \pm 0.3$
Middle-high	1,751	51.2 ± 1.4	$15.5\pm0.5$	$16.4\pm0.5$	$12.2\pm0.4$	$1.60\pm0.05$	$10.7\pm0.3$
High	1,742	54.4 ± 1.9	$16.3\pm0.7$	$17.5 \pm 0.7$	$12.9\pm0.4$	$1.84\pm0.07$	$11.1 \pm 0.4$

SFA, saturated fatty acid; MUFA, monounsaturated fatty acid; PUFA, polyunsaturated fatty acid; n-3 FA, n-3 fatty acid; n-6 FA, n-6 fatty acid, <sup>1)</sup>All estimates were weighted and calculated by considering the complex survey design. <sup>2)</sup>Age-standardized using the 2005 Korea population estimates, <sup>3)</sup>Divided into quartiles based on monthly household income (household income/  $\sqrt{}$  number of household members) by gender and 5-year age group.

Table 2. Food sources contributing to total fat intake<sup>1)</sup>

					3-5 ye	ars			6-11 years					12-18 years						
	Food name	Intake (g)	SE	% <sup>2)</sup>	Cum % <sup>3)</sup>	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	47.96	0.77	-	-	Total	41.16	3.20	-	-	Total	50.91	1.56	-	-	Total	60.17	1.60	-	-
1	Pork	7.52	0.34	15.7	15.7	Milk	6.23	0.36	15.1	15.1	Pork	6.31	0.67	12.4	12.4	Pork	9.08	0.85	15.1	15.1
2	Soybean oil	4.40	0.16	9.2	24.9	Pork	4.02	0.55	9.8	24.9	Milk	5.21	0.34	10.2	22.6	Soybean oil	6.42	0.62	10.7	25.8
3	Beef	2.82	0.15	5.9	30.7	Cookies, biscuits	3.68	2.49	9.0	33.9	Soybean oil	4.22	0.43	8.3	30.9	Instant noodle	3.84	0.40	6.4	32.1
4	Egg	2.58	0.07	5.4	36.1	Soybean oil	2.63	0.28	6.4	40.3	Egg	2.72	0.19	5.3	36.2	Beef	3.62	0.38	6.0	38.2
5	Instant noodle	2.33	0.16	4.9	41.0	Cheese	2.46	1.79	6.0	46.2	Beef	2.38	0.27	4.7	40.9	Egg	3.25	0.21	5.4	43.6
6	Milk	2.16	0.07	4.5	45.5	Egg	2.44	0.22	5.9	52.1	Bread	1.95	0.24	3.8	44.7	Milk	3.02	0.25	5.0	48.6
7	Mayonnaise	1.68	0.09	3.5	49.0	Cake	1.89	0.84	4.6	56.7	Instant noodle	1.67	0.23	3.3	48.0	Bread	2.86	0.33	4.8	53.3
8	Sesame oil	1.60	0.05	3.3	52.3	Bread	1.69	0.24	4.1	60.8	Snack	1.44	0.20	2.8	50.9	Chicken	1.93	0.33	3.2	56.5
9	Bread	1.59	0.10	3.3	55.6	Beef	1.40	0.19	3.4	64.2	Cookies, biscuits	1.40	0.31	2.8	53.6	Mayonnaise	1.72	0.24	2.9	59.4
10	Tofu	1.19	0.05	2.5	58.1	Ice cream	1.25	0.32	3.0	67.3	Chicken	1.37	0.27	2.7	56.3	Tofu	1.49	0.29	2.5	61.9
	19-29 years					30-49 years					50-64 years					65+ years				
	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	64.78	2.21	-	-	Total	52.62	1.15	-	-	Total	37.70	0.93	-	-	Total	23.74	0.73	-	-
1	Pork	11.75	1.29	18.1	18.1	Pork	8.80	0.60	16.7	16.7	Pork	5.42	0.40	14.4	14.4	Pork	2.81	0.36	11.8	11.8
2	Soybean oil	6.96	0.54	10.8	28.9	Soybean oil	5.07	0.27	9.6	26.4	Soybean oil	2.78	0.21	7.4	21.8	Beef	1.84	0.23	7.7	19.6
3	Instant noodle	3.98	0.39	6.2	35.0	Beef	3.16	0.25	6.0	32.4	Beef	2.15	0.21	5.7	27.5	Soybean oil	1.30	0.13	5.5	25.0
4	Beef	3.76	0.53	5.8	40.8	Egg	2.92	0.11	5.6	37.9	Egg	2.12	0.14	5.6	33.1	Milk	1.11	0.15	4.7	29.7
5	Egg	3.26	0.23	5.0	45.9	Instant noodle	2.67	0.35	5.1	43.0	Sesame oil	1.66	0.08	4.4	37.5	Egg	1.09	0.09	4.6	34.3
6	Mayonnaise	3.20	0.34	4.9	50.8	Mayonnaise	2.05	0.18	3.9	46.9	Milk	1.32	0.10	3.5	41.0	Soybean	1.00	0.09	4.2	38.6
7	Milk	2.36	0.20	3.7	54.4	Sesame oil	1.93	0.08	3.7	50.6	Instant noodle	1.23	0.14	3.3	44.3	Sesame oil	0.94	0.07	3.9	42.5
8	Bread	2.17	0.40	3.3	57.8	Milk	1.84	0.10	3.5	54.0	Tofu	1.22	0.09	3.2	47.5	Tofu	0.85	0.07	3.6	46.1
9	Chicken	2.14	0.42	3.3	61.1	Bread	1.62	0.15	3.1	57.1	Mayonnaise	1.14	0.14	3.0	50.5	Well polished rice	0.84	0.02	3.5	49.6
10	Sesame oil	1.76	0.15	2.7	63.8	Tofu	1.36	0.08	2.6	59.7	Coffee	1.04	0.06	2.8	53.3	Instant noodle	0.81	0.11	3.4	53.1

<sup>1)</sup> All estimates were weighted and calculated by considering the complex survey design. <sup>2)</sup> % of total intake, <sup>3)</sup> Cumulative % of total intake,

Table 3. Food sources contributing to SFA intake  $^{1\!\mathrm{)}}$ 

		3+ ye	ars				3-5 ye	ars			6-11 years					12-18 years				
	Food name	Intake (g)	SE	% <sup>2)</sup>	Cum % <sup>3)</sup>	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	14.36	0.26	-	-	Total	14.97	1.21	-	-	Total	17.43	0.61	-	-	Total	18.84	0.50	-	-
1	Pork	2.82	0.13	19.7	19.7	Milk	4.21	0.24	28.1	28.1	Milk	3.52	0.23	20.2	20.2	Pork	3.35	0.33	17.8	17.8
2	Milk	1.45	0.05	10.1	29.8	Pork	1.50	0.22	10.0	38.2	Pork	2.35	0.26	13.5	33.7	Milk	2.03	0.17	10.8	28.5
3	Instant noodle	1.03	0.07	7.2	37.0	Cheese	1.44	1.05	9.7	47.8	Beef	0.78	0.09	4.5	38.1	Instant noodle	1.70	0.18	9.0	37.6
4	Beef	0.88	0.05	6.2	43.1	Cookies, biscuits	1.05	0.50	7.0	54.8	Instant noodle	0.74	0.10	4.2	42.3	Beef	1.15	0.13	6.1	43.7
5	Egg	0.68	0.02	4.7	47.8	Ice cream	0.70	0.18	4.6	59.5	Egg	0.70	0.05	4.0	46.4	Soybean oil	0.90	0.09	4.8	48.5
6	Soybean oil	0.62	0.02	4.3	52.1	Egg	0.64	0.06	4.3	63.8	Ice cream	0.70	0.11	4.0	50.4	Egg	0.86	0.05	4.5	53.0
7	Bread	0.43	0.03	3.0	55.1	Beef	0.46	0.06	3.1	66.8	Chocolate	0.63	0.11	3.6	54.0	Ice cream	0.80	0.14	4.3	57.3
8	Coffee	0.43	0.02	3.0	58.1	Snack	0.44	0.08	3.0	69.8	Soybean oil	0.59	0.06	3.4	57.4	Bread	0.75	0.09	4.0	61.2
9	Ice cream	0.37	0.03	2.6	60.7	Bread	0.44	0.07	3.0	72.8	Snack	0.59	0.10	3.4	60.8	Cookies, biscuits	0.49	0.09	2.6	63.8
10	Cookies, biscuits	0.30	0.03	2.1	62.8	Instant noodle	0.40	0.23	2.7	75.5	Cheese	0.56	0.35	3.2	64.0	Chocolate	0.46	0.10	2.4	66.3
		19-29 y	ears				30-49 y	ears				50-64 y	ears				65+ ye	ars		
	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	19.67	0.75	-	-	Total	15.40	0.40	-	-	Total	10.52	0.26	-	-	Total	6.79	0.22	-	-
1	Pork	4.44	0.51	22.6	22.6	Pork	3.31	0.23	21.5	21.5	Pork	2.04	0.16	19.4	19.4	Pork	1.05	0.14	15.5	15.5
2	Instant noodle	1.77	0.17	9.0	31.6	Milk	1.24	0.07	8.0	29.5	Milk	0.90	0.07	8.5	27.9	Milk	0.75	0.10	11.0	26.6
3	Milk	1.58	0.13	8.0	39.6	Instant noodle	1.17	0.15	7.6	37.2	Beef	0.70	0.07	6.7	34.6	Beef	0.61	0.07	8.9	35.5
4	Beef	1.10	0.13	5.6	45.2	Beef	0.99	0.08	6.5	43.6	Coffee	0.56	0.03	5.3	39.9	Coffee	0.43	0.02	6.3	41.8
5	Soybean oil	0.97	0.08	5.0	50.1	Egg	0.77	0.03	5.0	48.6	Egg	0.56	0.04	5.3	45.2	Instant noodle	0.36	0.05	5.3	47.1
6	Egg	0.85	0.06	4.3	54.5	Soybean oil	0.71	0.04	4.6	53.2	Instant noodle	0.55	0.06	5.2	50.4	Well polished rice	0.34	0.01	4.9	52.1
7	Ice cream	0.59	0.11	3.0	57.5	Coffee	0.68	0.04	4.4	57.6	Soybean oil	0.39	0.03	3.7	54.1	Egg	0.28	0.02	4.1	56.2
8	Bread	0.55	0.09	2.8	60.3	Bread	0.45	0.05	2.9	60.5	Well polished rice	0.30	0.01	2.9	57.0	Bread	0.21	0.03	3.1	59.3
9	Chicken	0.50	0.11	2.5	62.8	Ice cream	0.35	0.06	2.3	62.8	Bread	0.29	0.04	2.8	59.7	Soybean oil	0.18	0.02	2.7	62.0
10	Mayonnaise	0.49	0.05	2.5	65.3	Mayonnaise	0.31	0.03	2.0	64.8	Sesame oil	0.24	0.01	2.2	62.0	Soybean	0.16	0.01	2.4	64.4

SFA, saturated fatty acid, <sup>1)</sup> All estimates were weighted and calculated by considering the complex survey design, <sup>2)</sup> % of total intake, <sup>3)</sup> Cumulative % of total intake,

**Table 4.** Food sources contributing to MUFA intake $^{1)}$ 

		3+ ye	ars				3-5 ye	ars					12-18 years							
	Food name	Intake (g)	SE	% <sup>2)</sup>	Cum % <sup>3)</sup>	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	15.32	0.28	-	-	Total	13.21	1.12	-	-	Total	16.54	0.54	-	-	Total	19.66	0.57	-	-
1	Pork	3.31	0.15	21.6	21.6	Milk	1.78	0.10	13.5	13.5	Pork	2.78	0.30	16.8	16.8	Pork	4.01	0.38	20.4	20.4
2	Beef	1.09	0.06	7.1	28.8	Pork	1.77	0.24	13.4	26.9	Milk	1.49	0.10	9.0	25.8	Soybean oil	1.49	0.14	7.6	28.0
3	Soybean oil	1.02	0.04	6.7	35.4	Cookies, biscuits	1.39	0.99	10.5	37.4	Egg	1.00	0.07	6.0	31.8	Beef	1.38	0.15	7.0	35.0
4	Egg	0.95	0.03	6.2	41.6	Egg	0.91	0.08	6.9	44.2	Soybean oil	0.98	0.10	5.9	37.7	Instant noodle	1.28	0.13	6.5	41.5
5	Instant noodle	0.77	0.05	5.1	46.7	Cheese	0.66	0.48	5.0	49.2	Beef	0.94	0.11	5.7	43.4	Egg	1.21	0.08	6.1	47.6
6	Milk	0.62	0.02	4.0	50.7	Soybean oil	0.61	0.06	4.6	53.8	Bread	0.60	0.08	3.6	47.0	Bread	0.88	0.10	4.5	52.1
7	Sesame oil	0.59	0.02	3.9	54.6	Bread	0.56	0.09	4.3	58.1	Instant noodle	0.56	0.08	3.4	50.4	Milk	0.87	0.07	4.4	56.5
8	Bread	0.48	0.03	3.2	57.7	Beef	0.56	0.07	4.2	62.3	Cookies, biscuits	0.50	0.12	3.0	53.4	Chicken	0.67	0.14	3.4	59.9
9	Chicken	0.37	0.04	2.4	60.2	Cake	0.47	0.20	3.6	65.9	Rape seed oil	0.50	0.11	3.0	56.5	Sesame oil	0.54	0.04	2.8	62.6
10	Mayonnaise	0.37	0.02	2.4	62.6	Snack	0.38	0.08	2.9	68.8	Snack	0.48	0.07	2.9	59.4	Snack	0.46	0.07	2.4	65.0
		19-29 y	ears			1	30-49 y	ears				50-64 y	ears				65+ ye	ars		
	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	21.00	0.85	-	-	Total	16.89	0.42	-	-	Total	11.76	0.34	-	-	Total	7.05	0.25	-	-
1	Pork	5.18	0.57	24.7	24.7	Pork	3.88	0.26	23.0	23.0	Pork	2.38	0.17	20.3	20.3	Pork	1.24	0.16	17.6	17.6
2	Soybean oil	1.62	0.13	7.7	32.4	Beef	1.24	0.10	7.3	30.3	Beef	0.89	0.09	7.5	27.8	Beef	0.76	0.09	10.7	28.3
3	Beef	1.33	0.16	6.4	38.7	Soybean oil	1.18	0.06	7.0	37.2	Egg	0.78	0.05	6.7	34.5	Egg	0.40	0.03	5.6	33.9
4	Instant noodle	1.32	0.13	6.3	45.0	Egg	1.08	0.04	6.4	43.7	Soybean oil	0.65	0.05	5.5	40.0	Sesame oil	0.35	0.03	4.9	38.9
5	Egg	1.20	0.09	5.7	50.7	Instant noodle	0.89	0.12	5.3	48.9	Sesame oil	0.62	0.03	5.2	45.2	Milk	0.32	0.04	4.5	43.4
6	Chicken	0.76	0.20	3.6	54.3	Sesame oil	0.71	0.03	4.2	53.2	Instant noodle	0.41	0.05	3.5	48.7	Soybean oil	0.30	0.03	4.3	47.6
7	Mayonnaise	0.70	0.07	3.3	57.7	Milk	0.53	0.03	3.1	56.3	Milk	0.38	0.03	3.2	51.9	Instant noodle	0.27	0.04	3.8	51.5
8	Milk	0.68	0.06	3.2	60.9	Bread	0.50	0.05	2.9	59.2	Rape seed oil	0.34	0.09	2.9	54.8	Bread	0.22	0.03	3.2	54.6
9	Sesame oil	0.65	0.06	3.1	64.0	Mayonnaise	0.45	0.04	2.7	61.8	Peanut	0.33	0.05	2.9	57.6	Rape seed oil	0.21	0.04	3.1	57.7
10	Bread	0.65	0.11	3.1	67.1	Rape seed oil	0.34	0.04	2.0	63.8	Almond	0.29	0.06	2.4	60.1	Tofu	0.17	0.02	2.4	60.1

MUFA, monounsaturated fatty acid. <sup>1)</sup> All estimates were weighted and calculated by considering the complex survey design. <sup>2)</sup> % of total intake, <sup>3)</sup> Cumulative % of total intake,

Table 5. Food sources contributing to PUFA intake  $^{\rm 1\! l}$ 

		3+ ye	ars				3-5 ye	ars			6-11 years					12-18 years				
	Food name	Intake (g)	SE	% <sup>2)</sup>	Cum % <sup>3)</sup>	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	11.56	0.19	-	-	Total	7.95	0.86	-	-	Total	10.37	0.40	-	-	Total	13.47	0.48	-	-
1	Soybean oil	2.53	0.09	21.9	21.9	Soybean oil	1.51	0.16	19.0	19.0	Soybean oil	2.42	0.25	23.4	23.4	Soybean oil	3.69	0.35	27.4	27.4
2	Pork	0.95	0.04	8.2	30.0	Cookies, biscuits	0.91	0.81	11.5	30.4	Pork	0.81	0.08	7.8	31.2	Pork	1.20	0.10	8.9	36.3
3	Mayonnaise	0.95	0.05	8.2	38.2	Pork	0.51	0.06	6.5	36.9	Egg	0.57	0.05	5.5	36.6	Mayonnaise	0.97	0.14	7.2	43.5
4	Sesame oil	0.68	0.02	5.9	44.1	Egg	0.48	0.05	6.0	42.9	Mayonnaise	0.52	0.10	5.0	41.6	Tofu	0.75	0.15	5.6	49.0
5	Tofu	0.59	0.03	5.1	49.2	Cake	0.46	0.23	5.7	48.6	Tofu	0.51	0.07	5.0	46.6	Egg	0.64	0.05	4.7	53.7
6	Egg	0.51	0.02	4.5	53.7	Bread	0.38	0.06	4.8	53.4	Sesame oil	0.51	0.04	4.9	51.5	Bread	0.63	0.10	4.7	58.5
7	Soybean	0.35	0.02	3.1	56.7	Tofu	0.30	0.04	3.8	57.1	Bread	0.43	0.07	4.2	55.7	Sesame oil	0.62	0.05	4.6	63.1
8	Instant noodle	0.35	0.03	3.1	59.8	Sesame oil	0.30	0.03	3.8	60.9	Instant noodle	0.26	0.04	2.5	58.2	Instant noodle	0.57	0.06	4.2	67.3
9	Bread	0.33	0.03	2.9	62.6	Mayonnaise	0.24	0.07	3.0	63.9	Rape seed oil	0.22	0.05	2.1	60.3	Cake	0.28	0.08	2.1	69.3
10	Perilla seed oil	0.20	0.02	1.7	64.3	Milk	0.22	0.01	2.7	66.6	Snack	0.22	0.04	2.1	62.4	Chicken	0.23	0.04	1.7	71.1
		19-29 y	ears			:	30-49 y	ears			50-64 years						65+ ye	ars		
	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	15.19	0.54	-	-	Total	12.93	0.28	-	-	Total	9.98	0.30	-	-	Total	6.23	0.22	-	-
1	Soybean oil	4.00	0.31	26.3	26.3	Soybean oil	2.91	0.16	22.5	22.5	Soybean oil	1.60	0.12	16.0	16.0	Soybean oil	0.75	0.08	12.0	12.0
2	Mayonnaise	1.80	0.19	11.8	38.1	Mayonnaise	1.15	0.10	8.9	31.4	Sesame oil	0.71	0.04	7.1	23.1	Soybean	0.61	0.05	9.8	21.8
3	Pork	1.46	0.14	9.6	47.8	Pork	1.10	0.07	8.5	39.9	Pork	0.67	0.05	6.7	29.8	Tofu	0.42	0.04	6.8	28.5
4	Sesame oil	0.75	0.07	4.9	52.7	Sesame oil	0.82	0.04	6.4	46.3	Mayonnaise	0.64	0.08	6.5	36.3	Sesame oil	0.40	0.03	6.4	34.9
5	Egg	0.65	0.06	4.3	57.0	Tofu	0.67	0.04	5.2	51.5	Tofu	0.60	0.05	6.0	42.3	Pork	0.36	0.04	5.7	40.6
6	Instant noodle	0.58	0.06	3.8	60.8	Egg	0.58	0.03	4.5	55.9	Soybean	0.58	0.04	5.8	48.1	Soybean milk	0.32	0.04	5.1	45.8
7	Tofu	0.53	0.08	3.5	64.2	Instant noodle	0.42	0.07	3.3	59.2	Egg	0.42	0.03	4.3	52.4	Perilla seed oil	0.24	0.05	3.9	49.7
8	Bread	0.46	0.13	3.0	67.3	Bread	0.34	0.04	2.7	61.8	Perilla seed oil	0.31	0.05	3.1	55.5	Egg	0.24	0.02	3.8	53.4
9	Chicken	0.25	0.04	1.7	68.9	Soybean	0.27	0.02	2.1	64.0	Peanut	0.26	0.04	2.7	58.1	Mayonnaise	0.23	0.05	3.7	57.1
10	Soybean	0.25	0.04	1.6	70.6	Walnut	0.20	0.05	1.6	65.5	Soybean milk	0.23	0.03	2.3	60.5	Well polished rice	0.19	0.00	3.1	60.2

PUFA, polyunsaturated fatty acid, <sup>1)</sup> All estimates were weighted and calculated by considering the complex survey design, <sup>2)</sup> % of total intake, <sup>3)</sup> Cumulative % of total intake,

Table 6. Food sources contributing to n-3 FA intake  $^{1\!\mathrm{)}}$ 

					3-5 ye	ars				6-11 y	ears			12-18 years						
	Food name	Intake (g)	SE	% <sup>2)</sup>	Cum % <sup>3)</sup>	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	1.593	0.034	-	-	Total	1.003	0.092	-	-	Total	1.263	0.061	-	-	Total	1.545	0.066	-	-
1	Soybean oil	0.337	0.012	21.2	21.2	Soybean oil	0.201	0.021	20.1	20.1	Soybean oil	0.323	0.033	25.6	25.6	Soybean oil	0.492	0.047	31.9	31.9
2	Perilla seed oil	0.158	0.017	10.0	31.1	Cake	0.098	0.056	9.8	29.9	Perilla seed oil	0.109	0.031	8.6	34.2	Mayonnaise	0.116	0.016	7.5	39.4
3	Mayonnaise	0.113	0.006	7.1	38.2	Perilla seed oil	0.075	0.025	7.5	37.3	Rape seed oil	0.063	0.014	5.0	39.2	Perilla seed oil	0.089	0.020	5.8	45.1
4	Mackerel	0.070	0.006	4.4	42.6	Cookies, biscuits	0.063	0.054	6.3	43.7	Mayonnaise	0.062	0.012	5.0	44.1	Tofu	0.078	0.015	5.0	50.2
5	Tofu	0.062	0.003	3.9	46.5	Rape seed oil	0.045	0.012	4.5	48.1	Tofu	0.054	0.008	4.3	48.4	Pork	0.061	0.005	4.0	54.1
6	Perilla seed	0.057	0.007	3.6	50.1	Bread	0.035	0.007	3.5	51.6	Mackerel	0.045	0.012	3.5	51.9	Bread	0.055	0.010	3.6	57.7
7	Pork	0.049	0.002	3.1	53.1	Perilla seed	0.032	0.016	3.2	54.8	Pork	0.042	0.004	3.3	55.3	Cake	0.049	0.019	3.2	60.9
8	Soybean	0.046	0.002	2.9	56.0	Mackerel	0.032	0.009	3.2	57.9	Egg	0.041	0.005	3.3	58.5	Egg	0.042	0.005	2.7	63.7
9	Rape seed oil	0.037	0.004	2.3	58.3	Egg	0.031	0.005	3.1	61.1	Bread	0.036	0.006	2.9	61.4	Rape seed oil	0.037	0.007	2.4	66.0
10	Eel	0.037	0.011	2.3	60.6	Tofu	0.031	0.004	3.1	64.2	Cake	0.030	0.010	2.4	63.7	Hamburger	0.027	0.007	1.7	67.8
		19-29 y	/ears			:	30-49 y	ears				50-64 y	/ears				65+ ye	ears		
	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)		%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	1.787	0.079	-	-	Total	1.773	0.054	-	-	Total	1.673	0.078	-	-	Total	1.082	0.058	-	-
1	Soybean oil	0.534	0.041	29.9	29.9	Soybean oil	0.388	0.021	21.9	21.9	Perilla seed oil	0.246	0.044	14.7	14.7	Perilla seed oil	0.194	0.038	17.9	17.9
2	Mayonnaise	0.215	0.023	12.1	41.9	Perilla seed oil	0.149	0.026	8.4	30.3	Soybean oil	0.213	0.016	12.7	27.5	Soybean oil	0.100	0.010	9.2	27.1
3	Perilla seed oil	0.106	0.037	6.0	47.9	Mayonnaise	0.138	0.012	7.8	38.1	Perilla seed	0.102	0.022	6.1	33.6	Perilla seed	0.082	0.017	7.6	34.7
4	Pork	0.076	0.008	4.2	52.1	Mackerel	0.085	0.013	4.8	42.9	Mackerel	0.101	0.015	6.0	39.6	Soybean	0.079	0.007	7.3	42.0
5	Tofu	0.055	0.008	3.1	55.2	Tofu	0.070	0.004	4.0	46.9	Mayonnaise	0.077	0.009	4.6	44.2	Mackerel	0.069	0.009	6.4	48.4
6	Egg	0.044	0.005	2.5	57.6	Pork	0.057	0.004	3.2	50.1	Soybean	0.075	0.005	4.5	48.7	Tofu	0.044	0.004	4.1	52.5
7	Mackerel	0.044	0.014	2.5	60.1	Eel	0.051	0.024	2.9	52.9	Eel	0.070	0.031	4.2	52.9	Soybean milk	0.038	0.005	3.5	56.0
8	Perilla seed	0.040	0.015	2.2	62.3	Perilla seed	0.046	0.010	2.6	55.5	Tofu	0.063	0.005	3.8	56.6	Anchovy	0.037	0.003	3.5	59.5
9	Bread	0.032	0.010	1.8	64.1	Anchovy	0.043	0.003	2.4	58.0	Anchovy	0.045	0.003	2.7	59.4	Doenjang	0.032	0.002	3.0	62.5
10	Soybean	0.032	0.005	1.8	65.9	Rape seed oil	0.042	0.005	2.4	60.3	Rape seed oil	0.043	0.011	2.6	61.9	Mayonnaise	0.027	0.006	2.5	65.0

n-3 FA, n-3 fatty acid. <sup>1)</sup> All estimates were weighted and calculated by considering the complex survey design. <sup>2)</sup> % of total intake, <sup>3)</sup> Cumulative % of total intake,

Table 7. Food sources contributing to n-6 FA intake<sup>1)</sup>

						6-11 ye	ears			12-18 years										
	Food name	Intake (g)	SE	% <sup>2)</sup>	Cum % <sup>3)</sup>	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	10.06	0.16	-	-	Total	7.04	0.80	-	-	Total	9.22	0.36	-	-	Total	12.01	0.43	-	-
1	Soybean oil	2.26	0.08	22.5	22.5	Soybean oil	1.35	0.14	19.1	19.1	Soybean oil	2.16	0.22	23.5	23.5	Soybean oil	3.29	0.32	27.4	27.4
2	Pork	0.97	0.04	9.7	32.1	Cookies, biscuits	0.85	0.76	12.0	31.2	Pork	0.82	0.09	8.9	32.4	Pork	1.20	0.11	10.0	37.4
3	Mayonnaise	0.83	0.05	8.3	40.4	Pork	0.53	0.07	7.5	38.7	Sesame oil	0.49	0.04	5.3	37.7	Mayonnaise	0.85	0.12	7.1	44.5
4	Sesame oil	0.66	0.02	6.5	46.9	Egg	0.40	0.04	5.7	44.4	Egg	0.48	0.04	5.2	43.0	Tofu	0.69	0.14	5.8	50.3
5	Tofu	0.55	0.02	5.4	52.4	Cake	0.36	0.18	5.1	49.5	Tofu	0.48	0.07	5.2	48.1	Sesame oil	0.60	0.04	5.0	55.3
6	Egg	0.44	0.01	4.4	56.7	Bread	0.32	0.05	4.5	53.9	Mayonnaise	0.46	0.09	5.0	53.1	Instant noodle	0.55	0.06	4.6	59.8
7	Instant noodle	0.34	0.03	3.4	60.1	Sesame oil	0.29	0.03	4.1	58.0	Bread	0.36	0.06	3.9	57.0	Egg	0.54	0.04	4.5	64.3
8	Soybean	0.31	0.02	3.1	63.2	Tofu	0.28	0.03	3.9	61.9	Instant noodle	0.25	0.04	2.7	59.8	Bread	0.52	0.09	4.4	68.7
9	Bread	0.27	0.03	2.7	65.9	Milk	0.26	0.02	3.7	65.7	Milk	0.22	0.01	2.4	62.1	Cake	0.23	0.06	1.9	70.6
10	Soybean milk	0.17	0.02	1.7	67.6	Mayonnaise	0.21	0.06	3.0	68.6	Snack	0.20	0.04	2.2	64.4	Chicken	0.21	0.03	1.8	72.4
		19-29 y	ears				30-49 y	ears				50-64 y	ears				65+ ye	ars		
	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %	Food name	Intake (g)	SE	%	Cum %
Rank	Total	13.55	0.49	-	-	Total	11.26	0.25	-	-	Total	8.37	0.24	-	-	Total	5.18	0.19	-	-
1	Soybean oil	3.57	0.28	26.4	26.4	Soybean oil	2.60	0.14	23.1	23.1	Soybean oil	1.43	0.11	17.1	17.1	Soybean oil	0.67	0.07	12.9	12.9
2	Mayonnaise	1.58	0.17	11.7	38.0	Pork	1.14	0.08	10.1	33.2	Pork	0.70	0.05	8.3	25.4	Soybean	0.53	0.05	10.3	23.1
3	Pork	1.52	0.16	11.2	49.2	Mayonnaise	1.01	0.09	9.0	42.2	Sesame oil	0.68	0.03	8.2	33.5	Tofu	0.39	0.03	7.6	30.7
4	Sesame oil	0.72	0.06	5.3	54.6	Sesame oil	0.79	0.03	7.0	49.2	Mayonnaise	0.57	0.07	6.8	40.3	Sesame oil	0.38	0.03	7.4	38.1
5	Instant noodle	0.56	0.06	4.1	58.7	Tofu	0.62	0.04	5.5	54.8	Tofu	0.56	0.04	6.6	47.0	Pork	0.36	0.05	7.0	45.1
6	Egg	0.55	0.05	4.1	62.8	Egg	0.49	0.02	4.4	59.1	Soybean	0.51	0.04	6.1	53.0	Soybean milk	0.28	0.03	5.4	50.5
7	Tofu	0.49	0.07	3.6	66.4	Instant noodle	0.40	0.07	3.6	62.7	Egg	0.36	0.03	4.3	57.3	Egg	0.20	0.02	3.9	54.4
8	Bread	0.38	0.12	2.8	69.2	Bread	0.29	0.04	2.6	65.3	Peanut	0.29	0.05	3.4	60.8	Mayonnaise	0.20	0.04	3.9	58.3
9	Chicken	0.23	0.04	1.7	70.9	Soybean	0.24	0.02	2.1	67.4	Soybean milk	0.21	0.03	2.5	63.2	Well polished rice	0.19	0.00	3.6	61.9
10	Soybean	0.22	0.03	1.6	72.5	Walnut	0.18	0.04	1.6	68.9	Sesame seed	0.19	0.02	2.3	65.6	Peanut	0.12	0.03	2.4	64.2

n-6 FA, n-6 fatty acid

<sup>1)</sup> All estimates were weighted and calculated by considering the complex survey design, <sup>2)</sup> % of total intake, <sup>3)</sup> Cumulative % of total intake,

tofu, soybean, or walnut) were relatively high in subjects 50 years of age and over.

The main sources of n-3 FA were soybean oil and perilla seed oil (Table 6). Cake, bread, or hamburger contributed 5-13% to n-3 FA intake in subjects 3-18 years of age. Mayonnaise contributed 8-12% in subjects 12-49 years of age, and perilla seed contributed 6-8% to n-3 FA intake in subjects 50 years of age and over. Fish (mackerel, eel, and anchovy) contributed 10% to n-3 FA intake in subjects 30 years of age and over.

The main source of n-6 FA in all age groups was soybean oil, providing 13-27% of n-6 FA intake (Table 7). Pork, mayonnaise, sesame oil, tofu, and egg also considerably affected intakes of n-6 FA.

The top ten sources of fats and fatty acids in men and women were similar. Highest ranked foods were the same (i.e. pork for total fat, SFA, and MUFA and soybean oil for PUFA, n-3 FA, and n-6 FA), and listed foods were similar. Ranked orders except for the first were slightly different (data not shown).

## DISCUSSION

This study produced population-level estimates of fatty acid intake and food sources according to socio-demographic status as a first attempt to estimate intake status of the Korean population using a comprehensive and reliable fatty acid composition table containing dietary intake data from KNHANES. Population mean of total fat intake was 48.0 g. PUFA was consumed less than SFA or MUFA, and mean intake levels of SFA, MUFA, and PUFA were 14.4 g, 15.3 g, and 11.6 g, respectively. Mean intakes of n-3 FA and n-6 FA were 1.6 g and 10.1 g, respectively, and the mean ratio of n-6 FA to n-3 FA was 6.9. Pork was the major source of total fat, SFA, and MUFA while soybean oil was the major source of PUFA. Milk and pork were major sources of SFA in subjects aged 3-11 years and  $\geq$  12 years, respectively. Perilla seed oil and soybean oil were the main sources of n-3 FA in subjects aged  $\geq$  50 years and < 50 years, respectively.

To reduce risk of coronary heart disease, substitution of unsaturated fatty acids for SFA, which raise low-density lipoprotein (LDL) cholesterol, is generally recommended. The ratio of unsaturated fatty acids to SFA was approximately 1.9 in the current study, which analyzed data from the KNHANES VI-1 (2013) containing a newly constructed food composition table, and this value was lower than 2.2 from the KNHANES III (2005) [11]. Direct comparison is not possible due to differences in the study conditions, including the food composition table. The ratio of n-6 FA to n-3 FA was within the current recommendation of 4-10 for Koreans [11].

The total fat consumption level in this study was lower than that of the US (48 g versus 79 g) estimated in 2004 using the 1999 to 2000 NHANES [17]. Average levels of fatty acid consumption were lower than those of the US [17] and Japan [18], although direct comparison is not feasible due to differences in the study conditions, including the study population (all age groups in US and middle-aged men with abdominal obesity or metabolic syndrome in Japan) and food composition table. The mean SFA intake level (14.4 g) was lower than those of the US (27 g) [17] and Japan (16.1 g) [18]. The average MUFA (15.3 g) and PUFA (11.6 g) intakes were lower than those of the US (MUFA 30 g and PUFA 16 g) [17] and Japan (MUFA 19.6 g and PUFA 15.8 g) [18].

Even though population averages for fat consumption were within adequate ranges, intake trends suggest that it will increase further and generate health concerns [12]. Examination of foods contributing to total fat and individual fatty acid intakes in this study provided information about food sources of fat and fatty acids in the Korean population. Better food choices can improve intakes of total fat and SFA and simultaneously increase proportions of PUFA and MUFA in the diet. The major sources of SFA and MUFA were somewhat similar, and the major source was pork in subjects aged 12 or more years. The major food sources of n-6 FA and n-3 FA were also similar, and the major source was soybean oil. Pork was the major source of total fat and fatty acids due to its high consumption in the Korean population. As fat intake increases at the population level along with increases in meat intake, food-based recommendations need to be developed to promote increased fat intake from plant-based foods such as oils, nuts, and seeds [rich sources of MUFA and alpha-linolenic acid (ALA)] as well as fish [primary food source of eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA)]. Due to the complexity of various fatty acids existing in the same foods, particularly coexistence of n-3 and n-6 FA in many food sources, nutritional education needs more complex responses than merely suggesting substitution of one food for another. Advances in processing that can produce foods with improved fatty acid compositions will also likely bring about population changes.

The Eskimo [19] and Japanese [20] are examples of populations that traditionally have high n-3 FA intakes, in particular higher intakes of long chain n-3 FA, and corresponding lower n-6 FA intakes. These intakes are mainly based on considerably high fish consumption as well as very lean red meat. Interestingly, soybean oil and perilla seed oil were the main food sources contributing to n-3 FA intake in our study. Perilla seed oil and raw perilla seed have often been used in traditional Korean cuisine. Similar to flaxseed oil, perilla seed oil contains 40-64% ALA, along with small amounts of other types of PUFA such as LA, which competes with ALA in the synthesis of EPA from ALA [21]. Large amounts of ALA in perilla seed oil have been reported to increase plasma levels of EPA and DHA in mice [22] and anti-atherosclerosis properties in Japanese quail [23]. Therefore, dietary intake of perilla seed oil may have cardioprotective effects.

There is inconsistent evidence linking SFA intake and cardiovascular disease. SFA intake is not associated with increased risk of CHD, stroke, or CVD according to the metaanalysis of 347,747 subjects during a follow-up of 5-23 y [24]. However, a JPHC study reported a positive association between SFA intake and cardiovascular disease, as the first report on the Asian population [25]. These inconsistent results may be attributable to dietary differences between the study populations. In this regard, population-specific differences, i.e., East and West, in the distribution of fat should be considered.

Korean dietary recommendations and population-based dietary

guidelines aim to provide adequate nutrient intakes and prevent chronic diseases. The consistent increase in meat and fat consumption as well as the dramatic increase in prevalence of overweight, metabolic disorders, chronic diseases in Korea highlights the need for better understanding of associations between food sources of fat and various fatty acids as well as disease risk. Therefore, further studies and health objectives should continue to focus on relations between fatty acid intake and related chronic diseases using this newly constructed data.

In conclusion, this study estimated fatty acid intakes in the Korean population based on a newly constructed fatty acid composition table using data from the KNHANES VI-1 (2013). This population-based descriptive data on fatty acid intake will serve as a basis for investigation of associations between dietary fatty acid intake and related chronic diseases as well as the development and evaluation of public health recommendations and national nutrition policies in Korea.

## REFERENCES

- World Health Organization (CH). The World Health Report 2002: Reducing Risks, Promoting Healthy Life. Geneva: World Health Organization; 2002.
- World Health Organization (CH). Diet, Nutrition and the Prevention of Chronic Diseases: Report of a Joint WHO/FAO Expert Consultation. WHO Technical Report Series 916. Geneva: World Health Organization; 2003.
- 3. Kris-Etherton P, Daniels SR, Eckel RH, Engler M, Howard BV, Krauss RM, Lichtenstein AH, Sacks F, St Jeor S, Stampfer M, Eckel RH, Grundy SM, Appel LJ, Byers T, Campos H, Cooney G, Denke MA, Howard BV, Kennedy E, Krauss RM, Kris-Etherton P, Lichtenstein AH, Marckmann P, Pearson TA, Riccardi G, Rudel LL, Rudrum M, Sacks F, Stein DT, Tracy RP, Ursin V, Vogel RA, Zock PL, Bazzarre TL, Clark J. Summary of the scientific conference on dietary fatty acids and cardiovascular health: conference summary from the nutrition committee of the American Heart Association. Circulation 2001;103: 1034-9.
- Grundy SM, Vega GL. Plasma cholesterol responsiveness to saturated fatty acids. Am J Clin Nutr 1988;47:822-4.
- Katan MB, Zock PL, Mensink RP. Dietary oils, serum lipoproteins, and coronary heart disease. Am J Clin Nutr 1995;61:13685-13735.
- Mensink RP, Katan MB. Effect of dietary fatty acids on serum lipids and lipoproteins. A meta-analysis of 27 trials. Arterioscler Thromb 1992;12:911-9.
- Hu FB, Stampfer MJ, Manson JE, Rimm E, Colditz GA, Rosner BA, Hennekens CH, Willett WC. Dietary fat intake and the risk of coronary heart disease in women. N Engl J Med 1997;337:1491-9.
- Kris-Etherton PM; American Heart Association. Nutrition Committee. AHA Science Advisory. Monounsaturated fatty acids and risk of cardiovascular disease. Circulation 1999;100:1253-8.
- Kris-Etherton PM, Harris WS, Appel LJ; American Heart Association. Nutrition Committee. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. Circulation 2002;106:2747-57.
- Harris WS, Mozaffarian D, Rimm E, Kris-Etherton P, Rudel LL, Appel LJ, Engler MM, Engler MB, Sacks F. Omega-6 fatty acids and risk for cardiovascular disease: a science advisory from the American Heart Association Nutrition Subcommittee of the Council on Nutrition, Physical Activity, and Metabolism; Council on Cardio-

vascular Nursing; and Council on Epidemiology and Prevention. Circulation 2009;119:902-7.

- 11. The Korean Nutrition Society. Dietary Reference Intakes for Koreans. 1st rev. ed. Seoul: The Korean Nutrition Society; 2010.
- 12. Ministry of Health and Welfare, Korea Centers for Disease Control and Prevention. Korea Health Statistics 2013: Korea National Health and Nutrition Examination Survey (KNHANES VI-1). Cheongju: Ministry of Health and Welfare, Korea Centers for Disease Control and Prevention; 2014.
- Yoon MO, Kim K, Hwang JY, Lee HS, Son TY, Moon HK, Shim JE. Development of fatty acids database using the Korea National Health and Nutrition Examination Survey data. J Nutr Health 2014;47:435-42.
- Baek Y, Kweon S, Oh K. Development of fatty acid composition table and intakes of fatty acids in Korea National Health and Nutrition Examination Survey (KNHANES). Public Health Wkly Rep 2015;8:75-81.
- Kweon S, Kim Y, Jang MJ, Kim Y, Kim K, Choi S, Chun C, Khang YH, Oh K. Data resource profile: the Korea National Health and Nutrition Examination Survey (KNHANES). Int J Epidemiol 2014;43: 69-77.
- Rural Development Administration, National Academy of Agricultural Science. Standard Food Composition Table. 8th rev. ed. Seoul: Kyomoonsa; 2011.
- Ervin RB, Wright JD, Wang CY, Kennedy-Stephenson J. Dietary intake of fats and fatty acids for the United States population: 1999-2000. Adv Data 2004:1-6

- Kawashima A, Sugawara S, Okita M, Akahane T, Fukui K, Hashiuchi M, Kataoka C, Tsukamoto I. Plasma fatty acid composition, estimated desaturase activities, and intakes of energy and nutrient in Japanese men with abdominal obesity or metabolic syndrome. J Nutr Sci Vitaminol (Tokyo) 2009;55:400-6.
- Bang HO, Dyerberg J, Sinclair HM. The composition of the Eskimo food in north western Greenland. Am J Clin Nutr 1980;33:2657-61.
- 20. Sugano M. Characteristics of fats in Japanese diets and current recommendations. Lipids 1996;31 Suppl:S283-6.
- Rao S, Abdel-Reheem M, Bhella R, McCracken C, Hildebrand D. Characteristics of high alpha-linolenic acid accumulation in seed oils. Lipids 2008;43:749-55.
- Chung KH, Hwang HJ, Shin KO, Jeon WM, Choi KS. Effects of perilla oil on plasma concentrations of cardioprotective (n-3) fatty acids and lipid profiles in mice. Nutr Res Pract 2013;7:256-61.
- Sadi AM, Toda T, Oku H, Hokama S. Dietary effects of corn oil, oleic acid, perilla oil, and evening [corrected] primrose oil on plasma and hepatic lipid level and atherosclerosis in Japanese quail. Exp Anim 1996;45:55-62.
- Siri-Tarino PW, Sun Q, Hu FB, Krauss RM. Meta-analysis of prospective cohort studies evaluating the association of saturated fat with cardiovascular disease. Am J Clin Nutr 2010;91:535-46.
- Yamagishi K, Iso H, Kokubo Y, Saito I, Yatsuya H, Ishihara J, Inoue M, Tsugane S; JPHC Study Group. Dietary intake of saturated fatty acids and incident stroke and coronary heart disease in Japanese communities: the JPHC Study. Eur Heart J 2013;34:1225-32.