



Effects of *Candida norvegensis* Live Cells on *In vitro* Oat Straw Rumen Fermentation

Oscar Ruiz, Yamicela Castillo^{1*}, Claudio Arzola, Eduviges Burrola, Jaime Salinas²,
Agustín Corral, Michael E. Hume³, Manuel Murillo⁴, and Mateo Itza¹

College of Animal Science and Ecology, Autonomous University of Chihuahua, Chihuahua, Chih. 31000, Mexico

ABSTRACT: This study evaluated the effect of *Candida norvegensis* (*C. norvegensis*) viable yeast culture on *in vitro* ruminal fermentation of oat straw. Ruminal fluid was mixed with buffer solution (1:2) and anaerobically incubated with or without yeast at 39°C for 0, 4, 8, 16, and 24 h. A fully randomized design was used. There was a decrease in lactic acid (quadratic, $p = 0.01$), pH, (quadratic, $p = 0.02$), and yeasts counts (linear, $p < 0.01$) across fermentation times. However, *in vitro* dry matter disappearance (IVDMD) and ammonia-N increased across fermentation times (quadratic; $p < 0.01$ and $p < 0.02$, respectively). Addition of yeast cells caused a decrease in pH values compared over all fermentation times ($p < 0.01$), and lactic acid decreased at 12 h ($p = 0.05$). Meanwhile, yeast counts increased ($p = 0.01$) at 12 h. *C. norvegensis* increased ammonia-N at 4, 8, 12, and 24 h ($p < 0.01$), and IVDMD of oat straw increased at 8, 12, and 24 h ($p < 0.01$) of fermentation. Yeast cells increased acetate ($p < 0.01$), propionate ($p < 0.03$), and butyrate ($p < 0.03$) at 8 h, while valerate and isovalerate increased at 8, 12, and 24 h ($p < 0.01$). The yeast did not affect cellulolytic bacteria ($p = 0.05$), but cellulolytic fungi increased at 4 and 8 h ($p < 0.01$), whereas production of methane decreased ($p < 0.01$) at 8 h. It is concluded that addition of *C. norvegensis* to *in vitro* oat straw fermentation increased ruminal fermentation parameters as well as microbial growth with reduction of methane production. Additionally, yeast inoculum also improved IVDMD. (**Key Words:** Rumen, Fermentation, Yeast, Oat Straw, Methane)

INTRODUCTION

Agricultural by-products, such as cereal straw from oats, wheat, and corn, constitute a great potential source of ruminant feed energy. Straws have low nutritional value, because of their low nitrogen and high indigestible fiber

content. In recent years, yeast-based additives, primarily *Saccharomyces cerevisiae* (*S. cerevisiae*), have been used to increase rumen feed utilization efficiency (Williams et al., 1991; Miller-Webster et al., 2002; Lila et al., 2004; Doležal et al., 2011; Chaucheyras-Durand et al., 2012). The beneficial effects associated with *S. cerevisiae* in animal studies include a greater dry matter (DM) and neutral detergent fiber digestibility. (Plata et al., 1994), as well as a higher feed utilization and milk production (Moallem et al., 2009). *In vitro* studies have also shown that yeast cultures favourably alter microbial fermentation (Marrero et al., 2013; Ye et al., 2014) and stimulate DM and cellulose digestion (Miller-Webster et al., 2002; Lila et al., 2004; Tang et al., 2008). In the same way, Marrero et al. (2015) showed that inclusion of two strains of yeast (Levazot 15 and Levica 25) in the *in vitro* fermentation of oat straw the accumulated gas production had a twofold increase as a result of yeast effect compared to control. Similar findings were reported by Marrero et al. (2014) when a yeast culture

* Corresponding Author: Yamicela Castillo. Tel: +52-636-110-9227, E-mail: ycastillo75@yahoo.com

¹ Department of Veterinary Medicine, Multidisciplinary Division, Autonomous University of Juarez City, Nuevo Casas Grandes, Chih. 31803, México.

² College of Veterinary Medicine and Animal Science, Autonomous University of Tamaulipas, Cd. Victoria, Tamps. 87000, México.

³ Agricultural Research Service, Southern Plains Research Center, Food and Feed Safety Research Unit, United States Department of Agriculture, College Station, TX 77843, USA.

⁴ College of Veterinary Medicine and Animal Science, Juarez University of Durango State, Durango, Dgo. 34000, Mexico.

Submitted Feb. 27, 2015; Revised May 18, 2015; Accepted May 31, 2015

(Levica 27) was included in the fermentation of corn stover. Aldo et al. (2006) also reported significant increases in the *in vitro* forage degradability when rice bran was treated with *Candida utilis* attributing this to the stimulation of rumen microbes by yeast. However, studies examining yeast cultures other than *S. cerevisiae* as rumen feed additives are scarce (Shin et al., 2002; Oeztuerk et al., 2005; Ando et al., 2006). In a previous study, Castillo (2009) isolated and identified the yeast *Candida norvegensis* and demonstrated the favourable effect of a non-*Saccharomyces* yeast addition on some fermentative parameters, such as gas production. The aim of the current experiment was to investigate the inoculation of a yeast (*Candida norvegensis*) and the effects on the *in vitro* ruminal fermentation of oat straw.

MATERIALS AND METHODS

Additives and Substrates

The yeast strain Levazoot 15 (*Candida norvegensis*) from the UACH yeast collection was used, with record number of the Gen Bank: JQ519367.1 GI: 386785959, was obtained in a previous study (Castillo, 2009) when the yeast was selected, isolated and identified from the rumen environment of dairy cows. Yeast strain samples from a culture of the *Candida norvegensis* strain were plated on malt extract broth and incubated for 20 h at 30°C±2°C. The probiotic was prepared by placing 30 mL of this inoculum into 1,200 mL of malt extract broth at identical conditions and incubation time, and containing 2.5×10⁸ live organisms/mL. Oat straw was used as substrate on a DM basis for *in vitro* incubation. The oat straw had been ground in a Wiley mill (Thomas-Wiley Model 4 Thomas Scientific, Swedesboro, NJ, USA) to pass through a 1 mm screen. The chemical composition of the cow's diet and the oat straw are shown in Table 1.

In vitro batch fermentation

Two cannulated lactating Holstein dairy cows (550±25.5 kg BW) were fed twice daily with 4.0 kg of a mix grain and 4 kg of corn silage (DM basis) (Table 1), and used as donor animals for ruminal liquor. The ruminal liquor was drawn before feeding (0600) from each cow with a vacuum pump and deposited in a 2-L hermetically sealed insulated flask that had been brought to the proper temperature and flushed with CO₂, and immediately taken to the laboratory. The mixed sample was strained through several layers of surgical gauze into a 1-L Erlenmeyer flask. The buffer solution had the following composition in 1.0 L: 9.8 g of NaHCO₃, 7 g of Na₂HPO₄·7H₂O, 0.57 g of KCl, 0.47 g of CaCl₂, 0.12 g of MgSO₄·7H₂O, 0.917 g of urea, and 0.917 g of glucose. The entire procedure was conducted under a CO₂ atmosphere to ensure anaerobic conditions and efforts were made to keep the temperature at 39°C. For each

Table 1. Chemical composition of feedstuffs

Item	Corn silage	Concentrate ¹	Oat straw ²
Percentage of daily ration	69.5	30.5	-
Chemical composition ³			
OM	94.7	96.4	88.7
CP	8.3	19.1	5.3
EE	2.9	2.6	2.3
ADF	31.0	9.5	45.8
NDF	75.4	20.2	70.3

OM, organic matter; CP, crude protein; EE, ether extract; ADF, acid detergent fiber; NDF, neutral detergent fiber.

¹ Contained (DM basis) 51.0% corn, 23.5% wheat bran, 10% cottonseed meal, 8.49% corn gluten meal, 2.0% sugarcane molasses, 1.5% soybean meal, 1.0% bypass fat, 0.8% CaCO₃, 0.5% urea, 0.5% animal fat, 0.2% NaCl, 0.5% trace mineral and vitamin premix. Trace mineral and vitamin premix contained: Mg, 0.003 Co, 0.001% Se, 0.140% Zn, 0.092% Mn, 0.052% Cu, 0.140% Fe, 2,500 IU of vitamin A/g, and 50 IU vitamin D/g.

² Substrate for *in vitro* fermentation.

³ Percentage dry mass basis.

sample, 50 mL of filtered rumen fluid and 100 mL of buffer solution were mixed in a 1:2 ratio. After mixing, 120 mL of diluted ruminal fluid was transferred under anaerobic conditions to 250 mL serum bottles containing 1.5 g of the substrate (oat straw) on a DM basis. Forty bottles were anaerobically sealed with butyl rubber stoppers and capped with aluminium. One set of 20 bottles, the control, was incubated at 39°C for 24 h with shaking at 30 rpm (New Brunswick Model Innova 4000, Nijmegen, Netherlands) and each bottle contained only the filtered rumen liquor and buffer solution. The remaining bottles (20) contained the same medium and was inoculated with 30 mL of the yeast probiotic and were incubated under identical conditions as the control group. Five sample times were set, 0, 4, 8, 12, and 24 h, (except for methane at 36 h) from the start of fermentation, with four repetitions for each time point. Consequently, the fermentation was conducted in a total of 40 bottles.

Microbiological determination

At every incubation time, 1 mL samples were collected to determine colony forming units (Log₁₀ colony-forming unit [CFU]/mL) of cellulolytic bacteria, cellulolytic fungi, and yeast using the Hungate (1969) culture roll-tube technique under strict anaerobic conditions. The cellulolytic bacteria were grown in culture media as described by Caldwell and Bryant (1966) with the modification described by Elías (1971). Dilutions of 10³, 10⁴, and 10⁵ were used to cultivate the bacteria and the fungi. The same dilutions were used for the fungi, and the culture media used was as specified by Joblin (1981). For the yeasts, malt extract agar was utilized with 0.01 g/L of chloramphenicol and three dilutions were used, 10⁴, 10⁵, and 10⁶. The number of CFU was determined by visual inspection of colonies on the tube

rolls under a magnifying glass.

Analyses

At the end of incubation methane gas was measured according to Theodorou et al. (1994), and total gas was measured by inserting a needle connected to a pressure transducer (Festo Co. Chihuahua, Chih. Mexico) where amount of gas was recorded and collected in a previously sealed tube using a 50-mL syringe. Tubes were retained and methane was measured by gas chromatography (Pye Unicam LTD, Cambridge, UK) using a flame ionization detector. The carrier gas was nitrogen, temperature of furnace was 52°C and detector temperature was 230°C. Silica gel columns were utilized. Methane concentration produced in each fermentation was calculated according to the general equation of gases ($N = PV/RT$). Atmospheric pressure was 10.1×10^4 Pa and room temperature was 299.5°K. The pH was immediately determined after bottles were uncapped with a portable pH meter (Hannah Instruments, Model HI 9017, Arvore-Vila do Conde, Portugal). Lactic acid (LA) concentrations ($\mu\text{g/mL}$) were determined by a colorimetric method as described by Taylor (1996). For analysis of ammonia nitrogen (mMol) and volatile fatty acids (VFA) (mMol), 1 mL of 25% metaphosphoric acid (wt/vol) was added to 5 mL of fermentation fluid and was stored at -20°C . The $\text{NH}_3\text{-N}$ was determined by the colorimetric method of Broderick and Kang (1980) using a spectrophotometer (Hach DR500, Loveland, CO, USA). After thawing, 1 mL of fermentation fluid was centrifuged ($10,000 \times g$ for 10 min) and VFA levels (mMol) were determined by gas chromatography (Perkin Elmer Model Clarus 800, Perkin Elmer Inc., Waltham, MA, USA) using a column (30 m \times 0.32 mm i.d.) and a flame ionization detector (column temperature 180°C, detector temperature 320°C, and injector temperature 240°C). The carrier gas was hydrogen with a flow rate of 1.40 mL/min. For *in vitro* DM disappearance (IVDMD) was determined according to Capetillo et al. (2002). the bottles' contents were transferred into test tubes (20 cc) previously dried at 105°C and centrifuged at $1,000 \times g$ for 5 min and the pellets were dried at 55°C for 48 h. The IVDMD was calculated by subtracting the dry residue weight (pellet) from the original weight of oat straw divided by the original sample weight, and the values multiplied by 100 to derive the percentage of IVDMD.

Statistical analysis

An analysis of variance was performed with the PROC MIXED procedure of SAS. (2002) (SAS Inst., Inc., Cary, NC, USA). The adjusted model included the fixed effects of yeast strain treatment and the control group, fermentation time, and the interaction between treatment and the time effect.

The model as fitted was as follows:

$$Y_{ijk} = \mu + T_i + M_j + L \times M_{(ij)} + \varepsilon_{ijk}$$

where Y_{ijk} = the dependent variable; μ = overall mean; T_i = treatment effect ($i = 1, 2$); M_j = Time effect ($j = 1, 2, 3, 4, 5$); $L \times M_{(ij)}$ = Interaction effect, and ε_{ijk} = random residual error.

The identity of the nested balloon flask (experimental unit) was used in the treatment as a random effect. A trend analysis was performed on the variables across fermentation times in each treatment group, using fermentation time as an indicator variable. Mean values are reported with standard errors, and p-values are declared statistically different when $p < 0.05$ or as indicated. Comparison of means were made using the predicted difference procedure of SAS (2002).

RESULTS AND DISCUSSION

pH value and lactic acid content

As shown in Table 2, the pH and LA concentrations in the medium decreased quadratically over time ($p < 0.01$ and $p < 0.02$, respectively). Treatment with yeast reduced pH relative to controls at all fermentation times ($p < 0.01$). Williams et al. (1991) reported similar results; they observed decreased pH when *S. cerevisiae* (10 g/d) was added to oat hay diet of young bulls. and by Lynch and Martin (2002), who studied the *in vitro* effect of *S. cerevisiae* on fermentation of Bermuda hay and alfalfa hay. Lila et al. (2004), Oeztuerk et al. (2005), Lattimer et al. (2007), Longuski et al. (2009), and Inal et al. (2010), reported *in vitro* studies where pH remained unchanged. LA concentration in the yeast treatment was lower than the control group at 12 h of fermentation ($p < 0.05$). At 24 h, the reduction was not statistically significant, although there was an appreciable numerical difference. These results confirmed findings by Williams et al. (1991), Erasmus et al. (1992), and Lila et al. (2004) who reported that the presence of yeast significantly reduced rumen LA concentrations and this reduction could be attributed to yeast cells stimulating the activity of *Selenomona ruminantum*, which consumes LA and thus contributes to pH stabilization.

Ammonia nitrogen concentration

The concentrations of ammonia-N (Table 3) increased quadratically with time of fermentation ($p < 0.01$). Yeast treatment reduced ammonia-N concentrations in the medium at 4, 8, 12, and 24 h. These results agreed with those of Erasmus et al. (1992) and Moallem et al. (2009) who found significant ammonia-N concentration decreases when yeast was added to dairy cow diets. Chaucheyras-Durand and Fonty (2001) also reported that ammonia-N

Table 2. Least squares means (\pm SE) of pH and lactic acid during *in vitro* oat straw rumen fermentation

Variable	Treatment			SEM	Trend		Treatment effect
	Time (h)	Control	Yeast		L	Q	
pH	0	7.56 ^a	7.36 ^b	0.045	***	**	***
	4	7.55 ^a	7.16 ^b	0.045			***
	8	7.37 ^a	7.02 ^b	0.045			***
	12	7.28 ^a	6.91 ^b	0.045			***
	24	7.08 ^a	6.68 ^b	0.045			***
Lactic acid (μ g/mL)	0	20.36 ^a	19.08 ^a	1.11	***	**	NS
	4	18.19 ^a	19.65 ^a	1.11			NS
	8	17.42 ^a	16.05 ^a	1.11			NS
	12	17.18 ^a	14.01 ^b	1.11			*
	24	16.58 ^a	14.68 ^a	1.11			NS

SEM, standard error of the mean; L, linear trend across fermentation times; Q, quadratic trend across fermentation times; NS, non significant.

Different letters in the same row indicate significant differences (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$) between treatments.

Means ($n = 40$) in rows with the same superscript are not significantly different (ns) due to treatment effect.

concentrations decreased when yeast was added to alfalfa hay and feeds for growing sheep. This ammonia-N reduction may be due to higher ammonia intake by microbial cells, perhaps as a direct result of stimulation of rumen microbial activity (Williams and Newbold, 1990).

Percentage of *in vitro* dry matter digestibility

As shown in Table 3, oat straw IVDMD increased quadratically with greater fermentation time ($p < 0.02$). Yeast treatment only increased IVDMD relative to controls at 24 h ($p < 0.01$), in agreement with *in vitro* results reported by Miller-Webster et al. (2002) and Lila et al. (2004). Williams et al. (1991) reported that stimulation of cellulose degradation by yeast cultures is associated with a decrease in the lag phase, which also causes an initial increase in digestion rate. These authors also attributed the improvement in DM digestibility to pH stabilization. Ando et al. (2004) as well, reported an increment in the total

degradability of forage when they added dried beer yeast attributing it to the activation of rumen microbes and partially due to the addition of nitrogen sources. Newbold et al. (1995) reported that some yeast cultures increased total and cellulolytic bacterial counts in the rumen, thus increasing fibre digestion. Another explanation may be that yeast cells in the rumen produce ethanol (Kung et al., 1997), which is converted by rumen microorganisms into valeric and isovaleric acids. The increases in these acids that were observed with the yeast treatment in this study are consistent with this latter possibility (Table 4). Valeric, capric, isobutyric, and isovaleric acids were found to stimulate rumen cellulolysis in studies involving fractional distillation of the rumen fluid (Elias, 1983). Likewise, in our study, higher concentrations of valeric and isovaleric acids were associated with increased rumen cellulolysis and *in vitro* disappearance of DM. The molar concentrations of acetic ($p < 0.01$), butyric ($p < 0.01$), propionic ($p < 0.01$),

Table 3. Least squares means (\pm SE) of ammonium nitrogen and *in vitro* dry matter disappearance during *in vitro* oat straw rumen fermentation

Variable	Treatment			SEM	Trend		Treatment effect
	Time (h)	Control	Yeast		L	Q	
NH ₃ -N	0	3.27 ^a	2.76 ^a	0.25	***	**	NS
	4	7.45 ^a	5.33 ^b	0.25			***
	8	7.04 ^a	5.51 ^b	0.25			***
	12	7.72 ^a	5.46 ^b	0.25			***
	24	9.44 ^a	6.46 ^b	0.25			***
IVDMD	0	9.76 ^a	12.63 ^a	2.74	***	**	NS
	4	10.96 ^a	14.59 ^a	2.74			NS
	8	17.64 ^a	21.27 ^a	2.74			NS
	12	25.00 ^a	30.33 ^a	2.74			NS
	24	46.64 ^a	56.71 ^b	2.74			**

SEM, standard error of the mean; L, linear trend across fermentation times; Q, quadratic trend across fermentation times;

NH₃-N, ammonium nitrogen (mMol); NS, non significant; IVDMD, *in vitro* dry matter disappearance (%).

Different letters in the same row indicate significant differences (** $p < 0.01$; *** $p < 0.001$) between treatments.

Means ($n = 40$) in rows with the same superscript are not significantly different (ns) due to treatment effect.

Table 4. Least squares mean (\pm SE) of concentrations of acetic, propionic, butyric, valeric and isovaleric acids during *in vitro* oat straw rumen fermentation

Volatile fatty acid (mMol)	Treatment			SEM	Trend		Treatment effect
	Time (h)	Control	Yeast		L	Q	
Acetic acid	0	24.40 ^a	20.00 ^a	1.97	***	*	NS
	4	24.56 ^a	26.82 ^a	1.97			NS
	8	29.50 ^a	37.42 ^b	1.97			***
	12	32.36 ^a	40.89 ^b	1.97			***
	24	45.04 ^a	48.84 ^a	1.97			NS
Propionic acid	0	8.55 ^a	6.72 ^b	0.58	***	NS	*
	4	8.22 ^a	9.09 ^a	0.58			NS
	8	9.99 ^a	11.90 ^b	0.58			*
	12	11.50 ^a	13.46 ^b	0.58			*
	24	17.26 ^a	17.79 ^a	0.58			NS
Butyric acid	0	5.81 ^a	4.48 ^b	0.42	***	**	*
	4	5.27 ^a	6.23 ^a	0.42			NS
	8	6.73 ^a	8.12 ^b	0.42			*
	12	7.61 ^a	8.50 ^a	0.42			NS
	24	9.73 ^a	9.68 ^a	0.42			NS
Valeric acid	0	0.50 ^a	0.45 ^a	0.06	***	NS	NS
	4	0.46 ^a	0.63 ^a	0.06			NS
	8	0.60 ^a	0.99 ^b	0.06			***
	12	0.67 ^a	1.31 ^b	0.06			***
	24	0.96 ^a	1.77 ^b	0.06			***
Isovaleric acid	0	0.44 ^a	0.47 ^a	0.08	*	**	NS
	4	0.42 ^a	0.46 ^a	0.08			NS
	8	0.48 ^a	0.78 ^b	0.08			**
	12	0.58 ^a	1.02 ^b	0.08			***
	24	0.99 ^a	1.69 ^b	0.08			***

SEM, standard error of the mean; L, linear trend across fermentation times; Q, quadratic trend across fermentation times; NS, non significant.

Different letters in the same row indicate significant differences (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$) between treatments.

Means ($n = 40$) in rows with the same superscript are not significantly different (ns) due to treatment effect.

valeric ($p < 0.01$), and isovaleric acids ($p < 0.02$) increased linearly over fermentation times (Table 4).

Acetic acid content

In the present study (Table 4), acetic acid concentrations were higher in the yeast treatment groups than in the controls at 8 h ($p < 0.01$) and 12 h ($p < 0.01$). Propionic acid was lower when yeast was added at 0 h ($p < 0.03$), but higher at 8 h ($p < 0.03$) and 12 h ($p < 0.02$). Meanwhile, butyric acid concentration was also lower at 0 h and higher at 8 h with yeast treatment. Valeric and isovaleric acid concentrations with yeast treatment were higher than that in control solutions at 8, 12, and 24 h ($p < 0.01$). Oeztuerk et al. (2005) studied the effects of live yeast on *in vitro* rumen fermentation of a hay and grain diet and found that acetic, butyric, valeric, and isovaleric acids increased, while only propionic acid did not increase significantly. Similar results were found by Erasmus et al. (2005); Miller-Webster et al. (2002). Diaz et al. (2011); Křižova et al. (2011); Kowalik et al. (2012). The stimulating effect of the strain under study on production of VFAs may be related to the chemical

composition of the yeast cell wall and other cell components. Yeast cell walls, which account for ~20% of the yeast cell weight, are primarily made up of β -1.3 and β -1.6 glucans, and chitin (Moukadiri et al., 1997). These structures, which form appropriate substrates for microbial rumen fermentation, independent of the state of the yeast (Oeztuerk et al., 2005).

Microorganisms counts

Table 5 shows the resulting CFU counts in viable yeast, cellulolytic bacteria, and cellulolytic fungi in the rumen. The quantity of viable yeast inoculated into the rumen decreased linearly with fermentation time ($p < 0.01$, Table 5). Similar results were reported by Arambel and Rung-Syin (1987) in studies examining the growth of *S. cerevisiae* in a ruminal environment. These authors indicate that yeasts are incapable of sustaining a productive population within a ruminal environment, because of inhibiting factors for yeast growth, such as non-optimal temperature. The optimal temperature range for yeast growth is 28°C to 30°C, with survival remaining possible up to 37°C through formation

Table 5. Least squares mean (\pm SE) of yeast and microorganism counts during *in vitro* oat straw rumen fermentation

Variable	Treatment			SEM	Trend		Treatment effect
	Time (h)	Control	Yeast		L	Q	
Yeast	0	5.42 ^a	7.30 ^b	0.10	***	***	***
	4	5.73 ^a	6.65 ^b	0.10			***
	8	5.69 ^a	6.25 ^b	0.10			***
	12	5.50 ^a	5.91 ^b	0.10			**
	24	5.45 ^a	5.67 ^a	0.10			NS
Cellulolytic fungi	0	5.88 ^a	5.78 ^a	0.19	NS	NS	NS
	4	5.26 ^a	6.09 ^b	0.19			**
	8	5.20 ^a	6.15 ^b	0.19			***
Cellulolytic bacteria	0	6.26 ^a	6.29 ^a	0.11	NS	NS	NS
	4	6.36 ^a	6.58 ^a	0.11			NS
	8	6.23 ^a	6.50 ^a	0.11			NS

SEM, standard error of the mean; L, linear trend across fermentation times; q, quadratic trend across fermentation times; NS, non significant.

Different letters in the same row indicate significant differences (** $p < 0.01$; *** $p < 0.001$) between treatments.

Means (Log_{10} CFU/mL, $n = 40$) in rows with the same superscript are not significantly different (ns) due to treatment effect.

of ascospores (Dengis et al., 1995). At 39°C, the typical temperature in the ruminal environment, growth and viability are reduced. (Mendoza, 1993). Williams et al. (1990) postulated that yeast do not establish themselves permanently in the rumen. In the current study, the inoculated rumen materials maintained a higher yeast population than the control for up to 12 h ($p < 0.01$), and was not significantly different until 24 h post-inoculation. After 12 h, the yeast cells entered a lethal no-growth phase, which corroborates the hypothesis of Williams et al. (1990). Also it was observed that, after 12 h, the differences in yeast populations between the inoculated and control samples were smaller, reaching a nadir at 24 h. In the present study, rumen material was tested for resident yeast soon after collection, and it was demonstrated that a pool of yeast populations existed in the rumen. Fermentation and treatment time did not affect cellulolytic bacterial counts ($p < 0.05$) (Table 5). The effects of live yeast on cellulolytic bacterial counts are very diverse. Similar to the present findings, Erasmus et al. (1992) found no effect of *S. cerevisiae* on the cellulolytic bacterial populations of dairy cows fed an energy-rich diet. Conversely, Newbold et al. (1996) and Lila et al. (2004) reported a positive effect of adding yeast cultures to *in vitro* fermentation of hay and

concentrate mixes. Counts of cellulolytic fungi (Table 5) did not exhibit any trends across the fermentation times. However, treatment with yeast cultures positively affected cellulolytic fungal populations in the rumen after 4 h ($p < 0.01$) and 8 h ($p < 0.01$) of fermentation. Chaucheyras et al. (1995) also report that yeast stimulate the growth of the ruminal fungus *Neocallimastix frontalis*. This stimulation effect on ruminal microorganisms, particularly on those that break down cellulose, such as cellulolytic fungi, may be explained by specific mechanisms of action in the rumen. When yeast are plasmolized, over time they supply growth factors such as peptides, amino acids, B-complex vitamins, and other components that favour bacterial and fungal growth (Elías, 1971). Also, live yeast help to eliminate a small amount of oxygen (~1%), that enters the rumen when the animal ingests feed, through aerobic respiration, and this process facilitates growth of more stringent anaerobic microorganisms, such as cellulolytic bacteria and fungi (Newbold et al., 1996).

Methane production

As shown in Table 6, methane production did not exhibit any trends across fermentation times. A reduction in methane production was observed only at 8 h by yeast

Table 6. Least squares mean (\pm SE) of methane production during *in vitro* oat straw rumen fermentation

Item	Treatment			SEM	Trend ²		Treatment effect
	Time (h)	Control	Yeast		L	Q	
Methane (mL)	4	45.00 ^a	28.78 ^a	11.53	NS	NS	NS
	8	66.00 ^a	15.06 ^b	11.53			**
	12	33.05 ^a	15.91 ^a	11.53			NS
	24	29.93 ^a	13.12 ^a	11.53			NS
	36	34.50 ^a	24.94 ^a	11.53			NS

SEM, standard error of the mean; L, linear trend across fermentation times; q, quadratic trend across fermentation times; NS, non significant.

Different letters in the same row indicate significant differences (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$) between treatments.

Means ($n = 40$) in rows with the same superscript are not significantly different (ns) due to treatment effect.

treatment ($p < 0.01$). At other time points, there were noticeable numerical decreases, although not statistically significant. Numerous studies have demonstrated that methane production is affected by addition of yeast. Mutsvangwa et al. (1992) observed a marked decrease in methane production with the addition of *S. cerevisiae* in intensive fattening of bulls. Lynch and Martin (2002) and Lila et al. (2004) also found reduced methane production when they studied the behaviour of yeasts added to *in vitro* fermentation of alfalfa hay and hay-concentrate mix. This drop in methane production may be due to the yeast stimulating utilization of metabolic hydrogen by acetogenic bacteria in the generation of acetic acid. (Chaucheyras et al., 1995). Based on the above results, addition of the yeast *Candida norvegensis* to *in vitro* oat straw fermentation positively influenced ruminal fermentation parameters as well as microbial growth with reduction of methane production. Additionally, yeast inoculum also improved IVDM.

CONFLICT OF INTEREST

We certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

REFERENCES

- Ando, S., Y. Nishiguchi, K. Hayasaka, H. Iefuji, and J. Takahashi. 2006. Effects of *Candida utilis* treatment on the nutrient value of rice bran and the effect of *Candida utilis* on the degradation of forages *in vitro*. Asian Australas. J. Anim. Sci. 19:806-810.
- Ando, S., R. I. Khan, J. Takahasi, Y. Gamo, R. Morikawa, Y. Nishiguchi, and K. Hayasaka. 2004. Manipulation of rumen fermentation by yeast: The effects of dried beer yeast on the *in vitro* degradability of forages and methane production. Asian Australas. J. Anim. Sci. 17:68-72.
- Arambel, M. J. and T. Rung-Syin. 1987. Evaluation of *Saccharomyces cerevisiae* growth in the rumen ecosystem. Memories 19th Biennial Conference on Rumen Function, Chicago, IL, USA. pp. 17-19.
- Broderick, G. A. and J. H. Kang. 1980. Automated simultaneous determination of ammonia and total amino acids in ruminal fluid and *in vitro* media. J. Dairy. Sci. 63:64-75.
- Caldwell, D. R. and M. P. Bryant. 1966. Medium without fluid for non-selective enumeration and isolation of rumen bacteria. Appl. Microbiol. 14:794-801.
- Capetillo, L. C. M., P. E. Herrera, and C. C. C. Sandoval. 2002. Chemical composition of boherhavia erecta L, digestibility and gas production *in vitro*. Arch. Zootec. 51:461-464.
- Castillo, Y. 2009. *In vitro* Fermentation to Obtain the Yeast *Candida norvegensis* in Mixes of Alfalfa with Fermented Apple Waste and Effects on the Microbial Activity. PhD Thesis. Facultad de Zootecnia y Ecología. Universidad Autónoma de Chihuahua. Chihuahua, México.
- Chaucheyras-Durand, F., G. Fonty, G. Bertin, and P. Gouet. 1995. Effects of live *Saccharomyces cerevisiae* cells on zoospore germination, growth, and cellulolytic activity of the rumen anaerobic fungus, *Neocallimastix frontalis* MCH3. Curr. Microbiol. 31:201-205.
- Chaucheyras-Durand, F. and G. Fonty. 2001. Establishment of cellulolytic bacteria and development of fermentative activities in the rumen of gnotobiotically-reared lambs receiving the microbial additive *Saccharomyces cerevisiae* CNCM I-1077. Reprod. Nutr. Dev. 41:57-68.
- Chaucheyras-Durand, F., E. Chevaux, C. Martin, E. Forano. 2012. Use of yeast probiotics in ruminants: Effects and mechanisms of action on rumen pH, fibre degradation, and microbiota according to the diet. Chapter 7. In: Probiotic in Animals. Edited by Everlon Cid Rigobelo. INTECH, <http://dx.doi.org/10.5772/50192>. pp.119-152.
- Dengis, P. D., L. R. Nelissen, and P. G. Rouxhet. 1995. Mechanisms of yeast flocculation comparison of top- and bottom-fermenting strains. Appl. Environ. Microbiol. 61:718-728.
- Diaz, A., C. Saro, M. L. Tejido, A. Sosa, M. E. Martinez, J. Galindo, M. D. Carro, and M. J. Ranilla. 2011. Effects of a yeast enzymatic hydrolyzate on *in vitro* ruminal fermentation. In: Challenging strategies to promote the sheep and goat sector in the current global context (Eds. M. J. Ranilla, M. D. Carro, H. Ben Salem, and P. Morand-Fehr). Zaragoza: CIHEAM, CSIC, Universidad de León, FAO. pp. 181-186. <http://om.ciheam.org/article.php?IDPDF=801554> Accessed June 2, 2014.
- Doležal, P., J. Dvořáček, J. Doležal, J. Čermáková, L. Zeman, and K. Szwedziak. 2011. Effect of feeding yeast culture on ruminal fermentation and blood indicators of Holstein dairy cows. Acta Vet. Brno. 80:139-145.
- Elias, A. 1971. The Rumen Bacteria of Animals Fed on a High Molasses-urea Diet. Ph. D Thesis. University of Aberdeen, Aberdeen, UK.
- Elias, A. 1983. Digestion of grasslands and tropical forages. In: The grasslands in Cuba, vol. 2. Ed. EDICA. La Habana, Cuba. pp.187-246.
- Erasmus, L. J., P. M. Botha, and A. Kistner. 1992. Effect of yeast culture supplement on production, rumen fermentation, and duodenal nitrogen flow in dairy cows. J. Dairy Sci. 75:3056-3065.
- Erasmus, L. J., P. H. Robinson, A. Ahmadi, R. Hinders, and J. E. Garrett. 2005. Influence of prepartum and postpartum supplementation of a yeast culture and monensin, or both, on ruminal fermentation and performance of multiparous dairy cows. Anim. Feed. Sci. Technol. 122:219-239.
- Hungate, R. E. 1969. A roll tube method for cultivation in microbiology (Eds. J. B. Morris and D. B. Ribbons). Academic Press Inc., New York, NY, USA. 117 p.
- Inal, F., E. Gürbüz, B. Coşkun, M. S. Malataş, O. B. Cital, E. S. Polat, E. Seker, and C. Ozcan. 2010. The Effects of live yeast culture (*Saccharomyces cerevisiae*) on rumen fermentation and nutrient degradability in yearling lambs. Kafkas Univ. Vet. Fak. 16:799-804.
- Joblin, K. N. 1981. Isolation, enumeration, and maintenance of rumen anaerobic fungi in roll tubes. Appl. Environ. Microbiol. 42:1119-1122.
- Kowalik, B., J. Skomial, J. J. Pająk, M. Taciak, M. Majewska, and G. Belżeczki. 2012. Population of ciliates, rumen fermentation

- indicators and biochemical parameters of blood serum in heifers fed diets supplemented with yeast (*Saccharomyces cerevisiae*) preparation. Anim. Sci. Pap. Rep. 30: 329-338.
- Křižová, L., M. Richter, J. Trínacty, J. Řiha, and D. Kumprechtová. 2011. The effect of feeding live yeast cultures on ruminal pH and redox potential in dry cows as continuously measured by a new wireless device. Czech J. Anim. Sci. 56:37-45.
- Kung, L. Jr., E. M. Kreck, R. S. Tung, A. O. Hession, A. C. Sheperd, M. A. Cohen, H. E. Swain, and J. A. Leedle. 1997. Effects of a live yeast culture and enzymes on *in vitro* ruminal fermentation and milk production of dairy cows. J. Dairy Sci. 80:2045-2051.
- Lattimer, J. M., S. R. Cooper, D.W. Freeman, and D. L. Lalman. 2007. Effect of yeast culture on *in vitro* fermentation of a high-concentrate or high-fiber diet using equine fecal inoculums in a Daisy II incubator. J. Anim. Sci. 85:2484-2491.
- Lila, Z. A., N. Mohammed, T. Yasui, Y. Kurokawa, S. Kanda, and H. Itabashi. 2004. Effects of a twin strain of *Saccharomyces cerevisiae* live cells on mixed ruminal microorganism fermentation *in vitro*. J. Anim. Sci. 82:1847-1854.
- Longuski, R. A., Y. Ying, and M. S. Allen. 2009. Yeast culture supplementation prevented milk fat depression by a short-term dietary challenge with fermentable starch. J. Dairy Sci. 92: 160-167.
- Lynch, H. A. and S. A. Martin. 2002. Effects of *Saccharomyces cerevisiae* culture and *Saccharomyces cerevisiae* live cells on *in vitro* mixed ruminal microorganism fermentation. J. Dairy Sci. 85:2603-2608.
- Marrero Y., M. E. Burrola-Barraza, Y. Castillo, L. C. Basso, C. A. Rosa, O. Ruiz, and E. González-Rodríguez. 2013. Identification of *Levica* yeasts as a potential ruminal microbial additive. Czech J. Anim. Sci. 58:460-469.
- Marrero, Y., O. Ruiz, A. Corrales, O. Jay, J. Galindo, Y. Castillo, and N. Madera. 2014. *In vitro* gas production of fibrous substrates with the inclusion of yeast. Cuban J. Agric. Sci. 48: 119-123.
- Marrero, Y., Y. Castillo, O. Ruiz, E. Burrola, and C. Angulo. 2015. Feeding of yeast (*Candida* spp.) improves *in vitro* ruminal fermentation of fibrous substrates. J. Integr. Agric. 14:514-519.
- Mendoza, M. G. D. and R. Ricalde-Velasco. 1993. Alimentación de ganado bovino con dietas altas en grano. Universidad Autónoma Metropolitana. Cap. 9. Uso de aditivos alimenticios. p. 97.
- Miller-Webster, T., W. H. Hoover, M. Holt, and J. E. Nocek. 2002. Influence of yeast culture on ruminal microbial metabolism in continuous culture. J. Dairy Sci. 85:2009-2014.
- Moallem, U., H. Lehrer, L. Livshitz, M. Zachut, and S. Yakoby. 2009. The effects of live yeast supplementation to dairy cows during the hot season on production feed efficiency, and digestibility. J. Dairy Sci. 92:343-351.
- Moukadiri, I., J. Armero, A. Abad, R. Sentandreu, and J. Zueco. 1997. Identification of a mannoprotein present in the inner layer of the cell wall of *Saccharomyces cerevisiae*. J. Bacteriol. 179:2154-2162.
- Mutsvangwa, T., I. E. Edwards, J. H. Topps, and G. F. M. Paterson. 1992. The effect of dietary inclusion of yeast culture (Yea-Sacc) on patterns of rumen fermentation, food intake and growth of intensively fed bulls. Anim. Prod. 55:35-40.
- Newbold, C. J., R. J. Wallace, X. B. Chen, and F. M. McIntosh. 1995. Different strains of *Saccharomyces cerevisiae* differ in their effects on ruminal bacterial numbers *in vitro* and in sheep. J. Anim. Sci. 73:1811-1818.
- Newbold, C. J., R. J. Wallace, and F. M. McIntosh. 1996. Mode of action of the yeast *Saccharomyces cerevisiae* as a feed additive for ruminants. Br. J. Nutr. 76:249-261.
- Oeztuerk, H., B. Schroeder, M. Beyerbach, and G. Breves. 2005. Influence of living and autoclaved yeasts of *Saccharomyces boulardii* on *in vitro* ruminal microbial metabolism. J. Dairy Sci. 88:2594-2600.
- Plata, P. F., M. G. D. Mendoza, J. R. Barcena-Gama, and M. S. Gonzalez. 1994. Effect of a yeast culture (*Saccharomyces cerevisiae*) on neutral detergent fiber digestion in steers fed oat straw based diets. Anim. Feed. Sci. Technol. 49:203-210.
- SAS. Institute. 2002. SAS. User's Guide. SAS Institute Inc. Cary, NC, USA.
- Shin, H. T., Y. Beom Lim, J. Ho Koh, J. Yun Kim, S. Young Baig, and J. Heung Lee. 2002. Growth of *Issatchenkia orientalis* in aerobic batch and fed-batch cultures. J. Microbiol. 40:82-85.
- Tang, S. X., G. O. Tayo, Z. L. Tan, Z. H. Sun, L. X. Shen, C. S. Zhou, W. J. Xiao, G. P. Ren, X. F. Han, and S. B. Shen. 2008. Effects of yeast culture and fibrolytic enzyme supplementation on *in vitro* fermentation characteristics of low-quality cereal straws. J. Anim. Sci. 86:1164-1172.
- Taylor, K. A. C. C. 1996. A simple colorimetric assay for muramic acid and lactic acid. Appl. Biochem. Biotechnol. 56:49-58.
- Theodorou, M. K., B. A. Williams, M. S. Dhanoa, A. B. McAllan, and J. France. 1994. A simple gas production method using a pressure transducer to determine the fermentation kinetics of ruminant feeds. Anim. Feed. Sci. Technol. 48:185-197.
- Williams, P. E., A. Tait, G. M. Innes, and C. J. Newbold. 1991. Effects of the inclusion of yeast culture (*Saccharomyces cerevisiae* plus growth medium) in the diet of dairy cows on milk yield and forage degradation and fermentation patterns in the rumen of steers. J. Anim. Sci. 69:3016-3026.
- Williams, J. G. K., A. R. Kubelik, K. J. Livak, J. A. Rafalski, and S. V. Tingey. 1990. DNA-polymorphism amplified by arbitrary primers is useful as genetic markers. Nucl. Acids Res. 18:6531-6535.
- Williams, P. E. V. and C. J. Newbold. 1990. Rumen probiosis: the effects of novel microorganisms on rumen fermentation and rumen productivity. In: (Eds. W. Haresing, and D. J. A. Cole), Recent Advances in Animal Nutrition. Butterworths, London, England. p. 211.
- Ye, G., Y. Zhu, J. Liu, X. Chen, and K. Huang. 2014. Preparation of glycerol-enriched yeast culture and its effect on blood metabolites and ruminal fermentation in goats. PLOS ONE 9(4):e94410.