

## ON SEMIDERIVATIONS OF NEAR-RINGS

KYUNG HO KIM\*

ABSTRACT. In this paper, we introduce the notion of a semiderivation on near-rings, and we try to generalize some properties of prime rings with derivations to prime near-rings with semiderivations.

### 1. Introduction

J. C. Chang [7] studied on semiderivations of prime rings. He obtained some results of derivations of prime rings into semiderivations. H. E. Bell and W. S. Martindale III [1] investigated the commutativity property of a prime ring by means of semiderivations. C. L. Chuang [8] studied on the structure of semiderivations in prime rings. He obtained some remarkable results in connection with semiderivations. J. Bergen and P. Grzeszczuk [3] obtained the commutativity properties of semiprime rings with the help of skew(semi)-derivations. A. Firat [9] generalized some results of prime rings with derivations to the prime rings with semiderivations. In this paper, we introduce the notion of a semiderivation on near-rings, and we try to generalize some properties of prime rings with derivations to prime near-rings with semiderivations.

### 2. Preliminaries

In this section, we include some elementary aspects that are necessary for this paper.

By a *near-ring* we mean a non-empty set  $N$  with two binary operations “+” and “.” satisfying the following axioms:

- (a)  $(N, +)$  is a group,

---

Received September 01, 2016; Accepted October 20, 2016.

2010 Mathematics Subject Classification: Primary 03G25, 06B10, 06D99, 06B35, 06B99.

Key words and phrases: near-rings, semiderivation, prime, 2-torsion free, commutative.

This work is supported by Korea National University of Transportation in 2016.

- (b)  $(N, \cdot)$  is a semigroup,  
 (c)  $x \cdot (y + z) = x \cdot y + x \cdot z$  for all  $x, y, z \in N$ .

Precisely speaking, it is a left near-ring because it satisfies the left distributive law. We will use the word “near-ring” in stead of “left near-ring”. We denote  $xy$  instead of  $x \cdot y$ . Note that  $x0 = 0$  and  $x(-y) = -xy$  but in general  $0x \neq 0$  for some  $x \in N$ . Let  $N$  be a near-ring. Then

$N$  is said to be *prime* if  $aNb = 0$  implies  $a = 0$ , or  $b = 0$  for all  $a, b \in N$ .  $N$  is said to be *2-torsion free* if  $2a = 0$  implies  $a = 0$  for all  $a \in N$ .

For any  $x, y \in N$ ,  $[x, y], x \circ y$  represent  $xy - yx, xy + yx$  respectively. The symbol  $Z(N)$  represent the multiplicative center of  $N$ , that is,  $Z(N) = \{x \in N | xy = yx \text{ for all } y \in N\}$ . A mapping  $f : N \rightarrow N$  is said to be *commuting* on  $N$  if  $[f(x), x] = 0$  for all  $x \in N$  and is said to be *centralizing* on  $N$  if  $[f(x), x] \in Z(N)$  for all  $x \in N$ . Let  $N$  be a near-ring. An additive mapping  $f : N \rightarrow N$  is called a *derivation* if

$$f(xy) = f(x)y + xf(y)$$

for all  $x, y \in N$ .

### 3. Semiderivations of prime near-rings

In what follows, let  $N$  denote a near-ring with center  $Z(N)$  unless otherwise specified.

**DEFINITION 3.1.** Let  $N$  be a near-ring. An additive mapping  $f : N \rightarrow N$  is called a *semiderivation* associated with a surjective function  $g : N \rightarrow N$  if

- (a)  $f(xy) = f(x)g(y) + xf(y) = f(x)y + g(x)f(y)$ ,  
 (b)  $f(g(x)) = g(f(x))$ , for all  $x, y \in N$ .

If  $g = I$ , i.e., an identity mapping of  $N$ , then all semiderivations associated with  $g$  are merely ordinary derivations. If  $g$  is any endomorphism, then semiderivations are of the form  $f(x) = x - g(x)$ .

**EXAMPLE 3.2.** Let  $N_1$  and  $N_2$  be two semiprime near-rings and  $N = N_1 \oplus N_2$ . If  $\alpha_1 : N_1 \rightarrow N_1$  be an additive map,  $\alpha_2 : N_2 \rightarrow N_2$  be a left and right  $N_2$  module map, which is not a derivation. Define a mapping  $f : N \rightarrow N$  such that  $f((r_1, r_2)) = (0, \alpha_2(r_2))$  and  $g : N \rightarrow N$  such that  $g((r_1, r_2)) = (\alpha_1(r_1), 0)$ , where  $r_1 \in N_1, r_2 \in N_2$ . Define addition and multiplication on  $N$  by  $(a_1, b_1) + (a_2, b_2) = (a_1 + a_2, b_1 + b_2)$  and  $(a_1, b_1)(a_2, b_2) = (a_1a_2, b_1b_2)$  for all  $a_1, a_2 \in N_1, b_1, b_2 \in N_2$ . Then it can

be easily verified that  $f$  is a semiderivations on  $N$ , with associated map  $g$  which is not a derivation.

We begin with the following lemma, which is essential in developing the proof of our main result.

LEMMA 3.3. *Let  $N$  be a prime near-ring and  $a \in N$ . If  $f$  is a nonzero semiderivation associated with a surjective function  $g : N \rightarrow N$ . If  $af(N) = 0$ , then  $a = 0$ .*

*Proof.* Suppose that  $af(N) = 0$  and  $x \in N$ . Since  $f \neq 0$ , there is  $y \in N$  such that  $f(y) \neq 0$ , and

$$af(xy) = 0.$$

Hence we obtain

$$(3.1) \quad a(f(x)g(y) + xf(y)) = af(x)g(y) + axf(y) = 0.$$

Since  $af(x) = 0$ , we have  $axf(y) = 0$  for every  $x \in N$ . Since  $N$  is prime and  $f(y) \neq 0$ , we get  $a = 0$ . □

LEMMA 3.4. *Let  $N$  be a prime near-ring and  $f$  be a semiderivation associated with a surjective function  $g : N \rightarrow N$ . Then*

$$(f(x)g(y) + xf(y))z = f(x)g(y)z + xf(y)z$$

for all  $x, y, z \in N$ .

*Proof.* For any  $x, y \in N$ , we have

$$\begin{aligned} f((xy)z) &= f(xy)g(z) + xyf(z) \\ &= (f(x)g(y) + xf(y))g(z) + xyf(z) \\ &= (f(x)g(y) + xf(y))z + xyf(z) \quad (g \text{ is onto}) \end{aligned}$$

On the other hand,

$$\begin{aligned} f(x(yz)) &= f(x)g(yz) + xf(yz) \\ &= f(x)g(yz) + xf(y)g(z) + xyf(z) \\ &= f(x)g(y)z + xf(y)z + xyf(z) \quad (g \text{ is onto}) \end{aligned}$$

Combining both expressions of  $f(xyz)$ , we have

$$(f(x)g(y) + xf(y))z = f(x)g(y)z + xf(y)z$$

for all  $x, y, z \in N$ . □

LEMMA 3.5. ([4]) *Let  $N$  be a 2-torsion free prime near-ring and let  $f$  be a nonzero semiderivation associated with a surjective function  $g : N \rightarrow N$ . If  $f(N) \subseteq Z(N)$ , then  $N$  is commutative.*

PROPOSITION 3.6. *Let  $N$  be a prime near-ring. If  $N$  admits a nonzero semiderivation  $f$  with  $g$  such that  $f(x \circ y) = x \circ y$  for all  $x, y \in N$ , then  $N$  is commutative.*

*Proof.* By hypothesis, we have

$$(3.2) \quad f(x \circ y) = x \circ y, \quad \forall x, y \in N.$$

Replacing  $y$  by  $xy$  in (3.2), we get

$$(3.3) \quad f(x \circ (xy)) = x^2y + xyx, \quad \forall x, y \in N.$$

Since  $x \circ (xy) = x(x \circ y)$ , we obtain

$$f(x \circ (xy)) = f(x)g(x \circ y) + xf(x \circ y), \quad \text{for all } x, y \in N.$$

By the equation (3.2), we get

$$x \circ (xy) = f(x)(x \circ y) + x(x \circ y), \quad \text{for all } x, y \in N.$$

Hence  $f(x)(x \circ y) = 0$  for all  $x, y \in N$ . This implies

$$(3.4) \quad f(x)xy = -f(x)yx, \quad \forall x, y \in N.$$

Substituting  $yz$  for  $y$  in (3.4), we obtain for all  $x, y \in N$ ,

$$(3.5) \quad -f(x)yzx = f(x)xyz = (-f(x)yx)z = f(x)y(-x)z.$$

Since  $-f(x)yzx = f(x)y(-x)z$ , (3.5) becomes

$$(3.6) \quad f(x)yz(-x) = f(x)y(-x)z, \quad \forall x, y \in N.$$

Taking  $-x$  instead of  $x$  in (3.6), we obtain

$$(3.7) \quad f(-x)yzx = f(-x)yxz, \quad \forall x, y \in N.$$

Hence  $f(-x)y(zx - xz) = 0$  and so

$$(3.8) \quad f(-x)N[z, x] = 0, \quad \forall x, y \in N.$$

By primeness, we have either  $x \in Z(N)$  or  $f(-x) = 0$ . That is,

$$(3.9) \quad f(x) = 0 \text{ or } [x, z] = 0, \quad \forall x, y \in N.$$

From (3.9), it follows that for each fixed  $x \in N$ , we have

$$(3.10) \quad f(x) = 0 \text{ or } x \in Z(N).$$

But  $x \in Z(N)$  also implies that  $f(x) \in Z(N)$ , and so the equation (3.8) forces to

$$(3.11) \quad f(x) \in Z(N), \quad \forall x, y \in N.$$

In light of (3.11),  $f(N) \subset Z(N)$  and using Proposition 3.5, we conclude that  $N$  is commutative.  $\square$

**THEOREM 3.7.** *Let  $N$  be a prime near-ring. If  $N$  admits a nonzero semiderivation  $f$  with  $g$  such that  $f(x \circ y) = -(x \circ y)$  for all  $x, y \in N$ , then  $N$  is commutative.*

*Proof.* By hypothesis,

$$(3.12) \quad f(x \circ y) = -(x \circ y), \quad \forall x, y \in N.$$

Replacing  $y$  by  $xy$  in (3.12), we have, for all  $x, y \in N$ ,

$$f(x \circ xy) = -(x \circ xy)$$

$$f(x(x \circ y)) = -x(x \circ y).$$

Hence we have  $f(x)g(x \circ y) + xf(x \circ y) = -x(x \circ y)$ . Using (3.12) in the above equation (3.12) and  $g$  is onto, we get

$$f(x)(x \circ y) = 0$$

for all  $x, y \in N$ . The rest of the proof is as in the proof Theorem 3.6.  $\square$

**THEOREM 3.8.** *Let  $N$  be a prime near-ring. If  $N$  admits a nonzero semiderivation  $f$  with  $g$  such that  $f(x) \circ y = x \circ y$  for all  $x, y \in N$ , then  $N$  is commutative.*

*Proof.* Suppose that

$$(3.13) \quad f(x) \circ y = x \circ y, \quad \forall x, y \in N.$$

Replacing  $x$  by  $xy$  in (3.13), we get

$$f(xy) \circ y = xy \circ y = (x \circ y)y.$$

Using the equation (3.13), we get  $f(xy) \circ y = (f(x) \circ y)y$ , and so

$$f(xy)y + yf(xy) = f(x)y^2 + yf(x)y$$

and

$$f(x)g(y)y + xf(y)y + yf(x)g(y) + yxf(y) = f(x)y^2 + yf(x)y.$$

Hence we have

$$f(x)y^2 + xf(y)y + yf(x)y + yxf(y) = f(x)y^2 + yf(x)y$$

since  $g$  is onto. Therefore we obtain  $xf(y)y + yxf(y) = 0$  for all  $x, y \in N$ , and so

$$(3.14) \quad yxf(y) = -xf(y)y, \quad \forall x, y \in N.$$

Replacing  $x$  by  $xz$  in the equation (3.14), we get for all  $x, y \in N$ ,

$$\begin{aligned} yxz f(y) &= -xz f(y)y = -x(zf(y)y) \\ &= -x(-yzf(y)) = -x(-y)zf(y). \end{aligned}$$

The last expression reduced to

$$(3.15) \quad yxz f(y) = -x(-y)zf(y), \quad \forall x, y \in N.$$

Since  $-yxzf(y) = (-y)xzf(y)$ , we have

$$(3.16) \quad (-y)xzf(y) = x(-y)zf(y), \quad \forall x, y \in N.$$

Taking  $-y$  instead of  $y$  in (3.16), we get  $yxzf(-y) = xyzf(-y)$  for all  $x, y \in N$ . Hence  $(yx - xy)zf(-y) = 0$ , and so  $[y, x]zf(-y) = 0$ , that is,

$$(3.17) \quad [y, x]Nf(-y) = 0, \quad \forall x, y \in N.$$

By primeness, we have either  $y \in Z(N)$  or  $f(-y) = 0$  for all  $y \in N$ . Accordingly,

$$(3.18) \quad f(y) = 0 \text{ or } y \in Z(N), \quad \forall y \in N.$$

But  $y \in Z(N)$  also implies that  $f(y) \in Z(N)$ , and so the equation (3.17) forces to

$$(3.19) \quad f(y) \in Z(N), \quad \forall y \in N.$$

In light of (3.19),  $f(N) \subset Z(N)$  and using Proposition 3.5, we conclude that  $N$  is commutative.  $\square$

**THEOREM 3.9.** *Let  $N$  be a prime near-ring. If  $N$  admits a nonzero semiderivation  $f$  with  $g$  such that  $f(x \circ y) = [x, y]$  for all  $x, y \in N$ , then  $N$  is commutative.*

*Proof.* Let

$$(3.20) \quad f(x \circ y) = [x, y], \quad \forall x, y \in N.$$

Replacing  $y$  by  $yx$  in (3.20), we obtain

$$f((x \circ y)x) = [x, y]x.$$

Hence we have

$$f(x \circ y)g(x) + (x \circ y)f(x) = [x, y]x.$$

Since  $g$  is onto and by (3.20), we get

$$(3.21) \quad (x \circ y)f(x) = 0, \quad \forall x, y \in N.$$

Replacing  $y$  by  $zy$  in (3.21), we obtain  $(x(zy) + (zy)x)f(x) = 0$  for all  $x, y, z \in N$ . Now, application of (3.21) yields  $yx f(x) = -xy f(x)$ . Combining this fact with the latter relation, we have  $xz + z(-x))yf(x) = 0$  for all  $x, y, z \in N$ . This implies that  $[x, z]yf(x) = 0$  for all  $x, y, z \in N$ . That is,

$$(3.22) \quad [x, z]Nf(x) = 0, \quad \forall x, y \in N.$$

Since  $N$  is a prime near-ring, for each  $x \in N$ , we have either  $f(x) = 0$  or  $x \in Z(N)$  for all  $x \in N$ . But  $x \in Z(N)$  also implies that  $f(x) \in Z(N)$ , and so the equation (3.22) forces to

$$(3.23) \quad f(x) \in Z(N), \quad \forall x, y \in N.$$

In light of (3.23),  $f(N) \subset Z(N)$  and using Proposition 3.5, we conclude that  $N$  is commutative.  $\square$

**THEOREM 3.10.** *Let  $N$  be a prime near-ring. If  $N$  admits a nonzero semiderivation  $f$  with  $g$  such that  $f[x, y] = x \circ y$  for all  $x, y \in N$ , then  $N$  is commutative.*

*Proof.* Let

$$(3.24) \quad f([x, y]) = x \circ y, \quad \forall x, y \in N.$$

Replacing  $y$  by  $yx$  in (3.24), we get, for all  $x, y \in N$ ,

$$f([x, y]x) = (x \circ y)x,$$

and so we obtain

$$f([x, y])g(x) + [x, y]f(x) = (x \circ y)x.$$

Since  $g$  is onto and using the equation (3.24), we get

$$(3.25) \quad [x, y]f(x) = 0,$$

Replacing  $y$  by  $yz$  in (3.25), we obtain

$$(3.26) \quad [x, y]Nf(x) = 0, \quad \forall x, y \in N.$$

Since  $N$  is a prime near-ring, for each  $x \in N$ , we have either  $f(x) = 0$  or  $x \in Z(N)$  for all  $x \in N$ . But  $x \in Z(N)$  also implies that  $f(x) \in Z(N)$ , and so the equation (3.26) forces to

$$(3.27) \quad f(x) \in Z(N), \quad \forall x, y \in N.$$

In light of (3.27),  $f(N) \subset Z(N)$  and using Proposition 3.5, we conclude that  $N$  is commutative.  $\square$

**PROPOSITION 3.11.** *Let  $N$  be a 2-torsion free prime near-ring and let  $f$  is a semiderivation associated with a surjective function  $g : N \rightarrow N$ . If  $f^2(x) = 0$  for all  $x \in N$ , then  $f = 0$ .*

*Proof.* By hypothesis, we have for all  $x \in N$ ,

$$f^2(x) = 0.$$

Replacing  $x$  by  $xy$  in the above equation, we obtain  $f^2(xy) = 0$  for all  $x, y \in N$ . Hence, for any  $x, y \in N$ ,

$$\begin{aligned} 0 &= f(f(xy)) \\ &= f(f(x)g(y) + xf(y)) \\ &= f^2(x)g(g(y)) + f(x)f(g(y)) + f(x)g(f(y)) + xf^2(y) \\ &= 2f(x)f(g(y)). \end{aligned}$$

Since  $N$  is 2-torsion free and  $g$  is surjective, we have  $f(x)f(y) = 0$  for all  $x, y \in N$ . Replacing  $y$  by  $yz$ , we get for all  $x, y, z \in N$ ,

$$\begin{aligned} 0 &= f(x)f(yz) \\ &= f(x)f(y)g(z) + f(x)yf(z) \\ &= f(x)yf(z). \end{aligned}$$

Hence we obtain  $f(x)sf(z) = 0$  for all  $x, z \in N$ . since  $N$  is prime,  $f(x) = 0$  or  $f(z) = 0$  for all  $x, z \in N$ . That is, in both cases,  $f = 0$ .  $\square$

### References

- [1] H. E. Bell. and W. S. Martindale III, *Semiderivations and commutativity in prime rings*, Canad. Math. **31** (1988), no. 4, 500-5008.
- [2] J. Bergen, *Derivations in prime rings*, Canad. Math. **26** (1983), 267-270.
- [3] J. Bergen and P. Grzeszczuk, *Skew derivations with central invariants*, J. London Math. Soc. **59** (1999), no. 2, 87-99.
- [4] A. Boua and L. Oukhtite, *Semiderivations satisfying certain algebraic identities on prime rings*, Asian-European Journal of Mathematics, **6** (2013), no. 3, 8.
- [5] M. Bresar, *On a generalization of the notion of centralizing mappings*, Proc. Amer. Math. Soc. **114** (1992), 641-649.
- [6] M. Bresar, *Semiderivations of prime rings*, Proc. Amer. Math. Soc. **108** (1990), no. 4, 859-860.
- [7] J. C. Chang, *On semiderivations of prime rings*, Chinese J. Math Soc. **12** (1984), 255-262.
- [8] C. L. Chuang, *On the structure of semiderivations in prime rings*, Proc. Amer. Math. Soc. **108** (1990), no. 4, 867-869.
- [9] A. Firat, *Some results for semiderivations of prime rings*, International J. of pure and Applied Mathematics **28** (1990), no. 3, 363-368.

\*

Department of Mathematics  
Korea National University of Transportation  
Chungju 380-702, Republic of Korea  
*E-mail:* ghkim@ut.ac.kr