

Kimchi Packaging Technology: An Overview

Suyeon Jeong and SeungRan Yoo*

World Institute of Kimchi, 86 Kimchi-ro, Gwangju, Korea

Abstract This paper provides an overview of kimchi packaging technology, focusing on packaging materials, package design, and active/intelligent packaging technology for kimchi. From a packaging-material standpoint, although various materials have been used to ensure customer satisfaction and convenience, plastic is the most widely used material, in the form of bags, trays, pouches, and rigid containers. Additionally, recent efforts in the kimchi packaging industry have allowed companies to differentiate their products by using different packaging materials and technologies, while simultaneously improving product safety and quality. On the other hand, the biggest problem in kimchi packaging is excess CO₂ production, leading to package expansion and leakage. To alleviate this problem, the use of CO₂ absorbers, high CO₂-permeable films, and degassing valves, in addition to the use of different packaging systems, has been investigated. Active and/or intelligent packaging systems have been developed, to include active functions beyond simply inert, passive containment and protection of the kimchi product. However, most such approaches are not yet adequately effective to be useful on a commercial scale. Therefore, further studies are needed to resolve the limitations of each technology.

Keywords Kimchi, Fermentation, Packaging, Technology

Introduction

Kimchi is the most well-known, traditional Korean food prepared from lactic acid fermentation of brined vegetables with the addition of various kinds of seasoning, giving it a unique flavor that is sour, spicy, salty, and carbonic. Kimchi has recently been demonstrated to promote health, by enhancing immunity as well as exhibiting anti-carcinogenic and anti-oxidative activities. Kimchi was listed among the “World’s Healthiest Foods” in 2006¹⁾, resulting in a rapid increase in kimchi popularity and consumption worldwide²⁾.

Preparing kimchi involves several complicated processes as shown in Fig. 1, including salting in brine for a couple of hours. Of note, the scarcity of ingredients in winter poses a challenge to the preparation of kimchi during this season. Therefore, a large amount of kimchi is prepared before the onset of winter and stored in kimchi jars buried in the ground, which is the traditional method for its long-term storage.

The concept of kimchi packaging and distribution began with its commercialization. The first commercial kimchi appeared in local grocery stores in the mid-1980s and quickly became a favorite of consumers because, despite its place as

an essential part of the cuisine in Korean households, preparation of high-quality kimchi requires significant effort, time, and good quality ingredients. Since 2010, Korea annually exports approximately 100 million USD worth of kimchi to over 60 countries. With the growth of the commercial kimchi market, kimchi packaging has gained importance. Today, commercial kimchi is packaged in plastic film bags or rigid plastic trays with a film lid for domestic consumption and export.

Fermentation of kimchi is brought about by various microorganisms and enzymes associated with the ingredients/raw materials, producing a unique, fresh sour flavor by generating various organic acids³⁾. Thus, the taste and texture of kimchi are affected over time by microorganisms during storage⁴⁾. Additionally, these microbiological and enzymatic activities continue even in well-aged kimchi, producing excess acid that results in a sour and bitter taste, offensive odor, and softening due to the deterioration of kimchi^{3,5)}. Excess acid production also leads to CO₂ accumulation in the packaging headspace, as shown in Fig. 2⁴⁾, which may cause damage owing to bloating and result in leakage⁶⁾. Therefore, in addition to its limited shelf life, the number one priority for kimchi manufacturers is a quick turnover to prevent kimchi package damage from CO₂ accumulation³⁾. Hence, the control of metabolic processes is key to preserving the quality of kimchi and extending its shelf life.

To solve the leakage problem, various processing methods such as canning, bottling, pasteurization, adding preservatives, use of food additives, irradiation, pinhole packaging systems,

*Corresponding Author : SeungRan Yoo
World Institute of Kimchi, 86 Kimchi-ro, Gwangju, Korea
Tel : +82-62-610-1738, Fax: +82-62-610-1850
E-mail : sryoo@wikim.re.kr




Fig. 1. Flow chart of kimchi preparation.




Fig. 2. Lactic acid fermentation of kimchi.

selective CO_2 -permeable films, and CO_2 absorbents have been introduced to improve product quality and extend the shelf life of kimchi products. However, while these new methods have improved kimchi packaging technology, some are not feasible for commercialization. In this review, we investigate the recent trends in kimchi packaging and technology development in the commercial kimchi industry.

General Trends in Kimchi Packaging

As for most other food packaging, the principal roles of kimchi packaging are to contain and distribute kimchi easily, to protect kimchi products from the external environment and damage, to decelerate kimchi product deterioration, to retain

the beneficial effects of fermentation, to extend shelf life, to maintain/increase the quality and safety of kimchi, and to provide consumers with ingredient and nutritional information. Prolonging the shelf life of kimchi requires retardation of enzymatic, microbial, and biochemical reactions using various strategies such as temperature control; moisture control; addition of chemicals such as salt, sugar, or natural acids; removal of oxygen; or a combination of these with effective packaging⁵⁻⁷.

The most commonly used kimchi packaging materials are plastic, metal, glass, and paper. Functional kimchi packaging materials were recently introduced in the market. These often combine several packaging materials to exploit each material's functional merits, as shown in Table 1^{8,9}. Each material has its own benefits and limitations in maintaining kimchi quality

Table 1. General package types for kimchi products in Korean markets⁹⁾

Package type	Construction	Photo
Plastic pouch	PET+Al+CPP, PET+Al+PE, PET+Al+LLDPE, PET+PE, PET+LLDPE, Nylon+PE, PET+PE+LLDPE, PE	
Plastic container and tray	PET (body)+PE, PET (body)+PP, PP (body)+PE	
Glass bottle	Glass (body)+metal, Glass (body)+PP, Glass (body)+PE	
Can	Metal, PET (body)+Al (top)	

and freshness during distribution and storage; therefore, careful consideration is required in the choice of packaging materials with respect to distribution and storage conditions.

Plastic is the most commonly used material for kimchi packaging, in the form of plastic bags, trays, pouches, and rigid containers. Plastic was first introduced in the late 1800s and the first major commercial plastic spray bottle was developed

in 1946 by Dr. Jules Montenier. Since then, it has been widely used in the food packaging industry. Owing to its superior cost effectiveness for mass production, moisture/grease barrier properties, and ease of deformation, plastic is currently one of the most popular packaging materials⁷⁾. However, there are growing concerns for adopting plastic in packaging materials, mainly because of its lack of degradability, the closing of landfill

sites, and growing water and land pollution issues.

Kimchi capacity is important for the consumer purchasing intention. Kimchi products are packaged in different sizes ranging from 40 g to 10 kg, and most are packaged in 500 g and 1 kg units⁹). Bulk kimchi products above 1 kg (family size) are available at grocery stores in plastic bags or rigid containers and are commonly packaged in plastic bags with cable ties. Kimchi products below 1 kg target single-person households and are most widely found at convenience stores and small local supermarkets in plastic pouches with aluminum foil, plastic trays with lids, or metal cans consisting of a tin plate (steel with a thin coating of tin) or aluminum and steel coated with a lacquer. Polyethylene terephthalate (PET) bodies with aluminum covers are increasingly used in the kimchi packaging market. Table 1 shows the various types of kimchi packaging described above.

With an increase in the kimchi market and sales, competitors attempt to differentiate themselves by changing their product packaging. One example of this is the development of a packaging method that mimics the traditional kimchi storage technique. Briefly, traditional kimchi storage involves putting freshly made kimchi in a special ceramic jar called an Onggi. A heavy stone is placed on top of the kimchi so that it is immersed in the fermentation liquid (Fig. 3). Several kimchi package technologies mimicking this traditional technique have been introduced in the market and are popular with customers because they improve the quality of taste and flavor during distribution and storage. The heavy object presses down on the kimchi packaging container, which maintains the quality of kimchi and prevents discoloration, bleaching, and contamination. Other materials used by manufacturers as packaging materials for market differentiation include paper and fabric. In a niche market, an Onggi designed by a master artisan is also used as a packaging material.

Beside the improvement of kimchi quality and extension of its shelf life, another great advancement in kimchi packaging is improved convenience in handling kimchi. A plastic container with a handle or a zippered bag that can be re-sealed makes the product more easy to use, thus giving a boost to the kimchi



Fig. 3. Plastic packaging mimicking an Onggi (Stone pressing down on the kimchi).

industry.

Packaging Technology to Prevent Product Expansion

Several kinds of lactic acid bacteria (LAB) are known to be involved in kimchi fermentation, including *Lactobacillus* and *Leuconostoc* species⁴). Moreover, some LAB grow under aerobic conditions in the initial fermentation stage, producing a series of metabolic products including lactic acid, acetic acid, and CO₂ that are responsible for kimchi's distinctive flavor⁴). As noted above, the CO₂ gas produced in kimchi during storage can result in damage to the packaging and cause leakage⁶). To solve the leakage problem, CO₂ absorbents, selective CO₂-permeable films, and pinhole packaging systems were introduced to prevent bloating of kimchi pouches⁶).

1. CO₂ absorber

Silica gel is widely used as a gas absorbent and some manufacturers use KOH, NaOH, Ca(OH)₂, MgCl₂, and CaCl₂ to absorb CO₂¹⁰). For this approach, a sachet of CO₂ absorbent is attached inside the kimchi packaging, as shown in Fig. 4, to remove the CO₂ that causes bloating⁸). Recently, Lee *et al.*¹¹) suggested using zeolite as a reversible CO₂ absorbent to relieve pressure buildup and/or bloating while maintaining an adequate level of equilibrated CO₂ partial pressure in the kimchi packaging. Additionally, Shin *et al.*⁸) demonstrated that Na₂CO₃ shows good potential for absorbing CO₂ under adequate water supply and used variations in the water vapor content and CO₂ transmission rate of the sachet and plastic sheet containing Na₂CO₃ to control the CO₂ absorption rate. Yoon *et al.*¹⁰) compared the utility of gas-absorbent materials such as KOH, NaOH, Ca(OH)₂, MgCl₂, and CaCl₂ to prevent package bloating and prolong the shelf life of kimchi. Their results showed that KOH and NaOH tend to retard the decrease in pH of kimchi. However, these materials fail to maintain suitable CO₂ levels in kimchi packaging and this can lead to depletion of CO₂ in the kimchi juice, thus causing loss of the product's characteristic fresh carbonic taste⁸).

2. Gas-permeable packaging

Traditionally, fermented foods including kimchi, are stored in Korean earthenware called an Onggi^{12,13}), which is made of yellow, pulverized clay that leaves small, air-permeable pores¹⁴). This inherent breathability is one of the unique characteristics of Onggi pottery and positively contributes to the quality of fermented foods¹²).

Using this principle, packaging materials have been designed with high-permeability films or micro-pores that eliminate the gas generated during storage and distribution. Laser perforation is the newest method to create micro-pores to increase gas permeability. Son *et al.*¹⁵) introduced femtosecond laser




Fig. 4. Attachment of gas absorbers inside kimchi pouches.

micro-perforated films for modified atmosphere packaging. In an alternate approach, Lee et al.¹²⁾ suggested the use of permeability-controlled polyethylene containers. Kimchi fermented in these containers showed a stable fermentation pattern with respect to pH and acidity, which was not observed for kimchi fermented in the other containers tested.

3. Degassing valve

Gas generated from the storage and distribution process can be released through a degassing valve. Yu et al.¹⁶⁾ designed one-way degassing valves with ring-type rubber disks, called SP valves, to allow pressure release from an airtight package while preventing air from reentering the package. They showed that the SP valve can be applied to fermented food packaging and it is expected to contribute to kimchi packaging. Other degassing valves used for products such as coffee or fresh-cut have been difficult to apply to kimchi packaging because of the risk of leakage. Therefore, further studies in this area are needed.

4. pH indicator

The pH and acidity are major indicators of the degree of

kimchi fermentation and can be used to detect the optimum point for ripeness of kimchi³⁾. Such indicators would change color as the pH of the aqueous solutions drops, reflecting the equilibrium between dissolved CO₂ and carbonic acid⁴⁾. Previous studies showed that the optimum conditions for kimchi were pH of 4.2 to 4.4 and acidity between 0.65 and 0.8%³⁾.

5. CO₂ indicator

In packaged kimchi, the partial pressure of CO₂ is an important factor indicating product quality since LAB continually produce CO₂ as a by-product of fermentation during storage and distribution³⁾. The development of a visual indicator for the partial pressure of CO₂ would be the easiest way to provide untrained consumers with an easy method to monitor kimchi quality⁴⁾. A color-indicating sachet consisting of a CO₂ absorbent and a chemical dye has been developed and evaluated for its application to kimchi packaging¹⁷⁾. Under packaged conditions, the partial pressure profile of CO₂ changes with storage temperature and time, and an indicator reflecting changes in CO₂ levels in the package headspace can provide indirect information about kimchi quality³⁾. Jung et al.³⁾ and Meng et al.⁴⁾ developed an irreversible chitosan-based

Table 2. Active packaging technologies for kimchi packaging to control gas production

Technology	Methods	Materials	Reference
CO ₂ absorbent	A gas absorber was attached inside flexible pouches.	Commercial CO ₂ absorbent (Ca(OH) ₂)	Lee et al. ⁶⁾
	Na ₂ CO ₃ -based CO ₂ absorbents were packed in sachets.	Calcium oxide, Magnesium Hydroxide, Calcium hydroxide	Shin et al. ⁸⁾
	A sachet of the CO ₂ absorbent was attached inside kimchi packages (Performance comparison of gas-absorbing materials).	Potassium hydroxide, Calcium hydroxide, Sodium hydroxide, Magnesium chloride, Calcium chloride	Yoon et al. ¹⁰⁾
Gas permeable packaging	Micro-perforated film was prepared for modified atmosphere packaging.	Femtosecond laser	Son et al. ¹³⁾
	Perforated films was used as gas permeable packaging.	Laser perforated BOPP films	Golestan et al. ¹⁸⁾
Degassing valve	Ring-type rubber disk valve was used for degassing.	Ring-type rubber disk	Yu et al. ¹⁷⁾

Table 3. Intelligent packaging technology (CO₂ indicator) to detect fermentation of kimchi

Methods	Materials	Reference
Brilliant blue-incorporated chitosan-based CO ₂ indicator solution was poured into and sealed in the LDPE sachet. The indicator sachet was then placed in the headspace of the package.	Chitosan, oxalic acid, Coomassie brilliant blue G-250, Hydrochloric acid, Acetic acid, Sodium hydroxide, Methanol	Meng <i>et al.</i> ⁴⁾
The chitosan-based indicator solution for CO ₂ detection was sealed into square plastic sachets made of LDPE film. The indicator sachet bag was placed in the headspace of the kimchi package.	Chitosan powder, 2-methyl 1-propanol (AMP), Hydrogen chloride, Sodium hydroxide	Jung <i>et al.</i> ³⁾
LDPE sachet attached	Calcium hydroxide and bromocresol purple or methyl red	Hong <i>et al.</i> ¹⁹⁾

**Fig. 5.** Indicator technology to detect fermentation degree of kimchi.

indicator to monitor kimchi quality corresponding to the partial pressure of CO₂ in the packaging headspace (Table 3). Thus, CO₂ indicators can indirectly identify the starting point of optimum ripening of kimchi since this parameter shows good correlation with the quality of packaged kimchi³⁾. Overall, these studies suggest that color indicators can be employed as an effective technique to detect the ripeness of packaged kimchi products without destroying the package^{3,4,17)}. In order to detect product quality in the package, there must be direct contact between the food product/headspace and the quality detector. However, a major cause for concern is that most of intelligent packaging systems require that food be in direct contact with the sensor, and substances from the sensor may migrate into the food. Additionally, there have been no studies to date on holistic quality indicators and very little work has been done to develop gas indicators that monitor fermentation degree of kimchi⁴⁾.

Conclusions

This paper provides an overview of the current state of kimchi packaging technology, focusing on packaging materials, package design, and active/intelligent packaging technology for kimchi. It summarizes the current state of the field and underscores the need for further research and investment into

the development of advanced technologies for industrial production, storage, and distribution of kimchi. Overall, we demonstrate the growth potential of the kimchi packaging industry. Additionally, we emphasize the need for more in-depth studies and greater investment in the development of advanced technologies for industrial applications, because most such approaches are not yet sufficiently effective for commercial application. Therefore, further studies are needed to resolve the limitations of each technology.

Acknowledgements

This research was supported by High Value-added Food Technology Development Programs(114087-03&316070-02), Ministry of Agriculture, Food and Rural Affairs, Republic of Korea.

References

- Choi, I. H., Noh, J. S., Han, J. S., Kim, H. J., Han, E. S. and Song, Y. O. Kimchi, a fermented vegetable, improves serum lipid profiles in healthy young adults: Randomized clinical trial. *J Med Food*. 16: 223-229.
- Kim, J. H., Park, J. G., Lee, J. W., Kim, W. G., Chung, Y. J. and Byun, M. W. 2008. The combined effects of N2-packag-

- ing heating and gamma irradiation on the shelf-stability of kimchi, Korean fermented vegetable. *Food Control*. 19: 56-61.
3. Jung, J. H., Lee, K. H., Puligundla, P. and Ko, S. H. 2013. Chitosan-based carbon dioxide indicator to communicate the onset of kimchi ripening. *LWT-Food Sci Technol*. 54: 101-106.
 4. Meng, X., Lee, K. H., Kang, T. Y. and Ko, S. H. 2015. An irreversible ripeness indicator to monitor the CO₂ concentration in the headspace of packaged kimchi during storage. *Food Sci Biotechnol*. 24: 91-97.
 5. Kim, Y. W., Jung, J. K., Cho, Y. J., Lee, S. J., Kim, S. H., Park, K. Y. and Kim, S. A. 2009. Quality changes in brined Baechu cabbage using different types of polyethylene film, and salt content during storage. *Korean J Food Preserv*. 16: 605-611.
 6. Lee, J. W., Cha, D. S., Hwang, K. T. and Park, H. J. 2003. Effects of CO₂ absorbent and high-pressure treatment on the shelf-life of packaged kimchi products. *Int J Food Sci technol*. 38: 519-524.
 7. Marsh, K. and Bugusu, B. 2007. Food packaging - Roles, materials, and environmental issues. *J Food Sci*. 72: 39-55.
 8. Shin, D. H., Cheigh, H. S. and Lee, D. S. 2002. The use of Na₂CO₃-based CO₂ absorbent systems to alleviate pressure buildup and volume expansion of kimchi packages. *J Food Eng*. 53: 229-235.
 9. Lee, D. S., Kwon, H. R., An, D. S., Chung, M., Lee, K. S. and Yang, D. J. 2012. Technology innovation in kimchi packaging for marketing in food supply chain. *Korean J Packaging Sci Technol*. 18: 1-8.
 10. Yoon, K. Y., Kang, M. J., Lee, K. H., Youn, K. S. and Kim, K. S. 1997. Studies of prevention of gas production during manufacturing and circulation of kimchi. *J Food Sci Technol*. 9: 145-151.
 11. Lee, D. S., Shin, D. H., Lee, D. U., Kim, J. C. and Cheigh, H. S. 2001. The use of physical carbon dioxide absorbents to control pressure buildup and volume expansion of kimchi packages. *J Food Eng*. 48: 183-188.
 12. Lee, E. J., Park, S. E., Choi, H. S., Han, G. J. and Kang, S. A. 2010. Quality characteristics of kimchi fermented in permeability-controlled polyethylene containers. *Korean J Food Preserv*. 17: 793-799.
 13. Jeong, J. K., Kim, Y. W., Choi, H. S., Lee, D. S., Kang, S. A., and Park, K. Y. 2011. Increased quality and functionality of kimchi when fermented in Korean earthenware (onggi). *Int J Food Sci Technol*. 46: 2015-2021.
 14. Kim, S., No, H., Kim, U. and Cho, W. S. 2014. A study on sources of pore formation in Onggi via the comparison with porcelain. *J Korean Ceram Soc*. 51: 11-18.
 15. Son, I. B., No, Y. C., Choi, S. C., Ko, D. K., Lee, J. M. and Choi, Y. J. 2007. *J. Opt. Soc. Korea* 2007. 2: 8-9.
 16. Yu, H. K., Lee, K. H. and O, J. Y. 2014. Development of degassing valves for food packaging using ring type rubber disk. *Korean J Packaging Sci Technol*. 20: 35-39.
 17. Hong, S. I. and Park, W. S. 1999. Development of color indicators for kimchi packaging. *J Food Sci*. 64: 255-257.
 18. Golestan, M. N., Ghosta, Y., Pourmiza, A. A. and Valizadegan, O. 2015. Study on laser perforated films as gas permeable packaging for confused flour beetle (*Tribolium confusum* Jacquelin du Val.) control inside food packaging. *J Stored Prod Res*. 60: 54-59.
 19. Hong, S. I. and Park, W. S. 1997. Sensitivity of color indicators to fermentation products of kimchi at various temperatures. *Korean J Food Sci Technol*. 29: 21-25.
- 투고: 2016.08.08 / 심사완료: 2016.11.02 / 게재확정: 2016.11.10