



# In Vitro Scolicidal Effects of *Salvadora persica* Root Extract against Protoscolices of *Echinococcus granulosus*

Abdel-Azeem S. Abdel-Baki<sup>1,2</sup>, Esam Almalki<sup>1,3</sup>, Lamjed Mansour<sup>1,4</sup>, Saleh Al-Quarishi<sup>1,\*</sup>

<sup>1</sup>Zoology Department, College of Science, King Saud University, Riyadh, Saudi Arabia; <sup>2</sup>Zoology Department, Faculty of Science, Beni-Suef University, Beni-Suef, Egypt; <sup>3</sup>Department of Medical Laboratory, College of Applied Medical Sciences, Majmaah University, Saudi Arabia;

<sup>4</sup>Unité de Recherche de Biologie intégrative et Ecologie évolutive et Fonctionnelle des Milieux Aquatiques, Département de Biologie, Faculté des Sciences de Tunis, Université De Tunis El Manar, Tunis, Tunisia

**Abstract:** It has been known that Arak, *Salvadora persica*, has a number of medicinal properties. We tried to investigate in vitro scolicidal effect of root extracts of this plant against protoscolices from hydatid cysts of *Echinococcus granulosus*. Protoscolices were aseptically collected from sheep livers containing hydatid cysts. *S. persica* root extract was used in 10, 30, and 50 mg/ml concentration for 10, 20, and 30 min. The viability of protoscolices was ascertained by 0.1% eosin staining. Scolicidal activity of *S. persica* extract at a concentration of 10 mg/ml was 36.3%, 50.3%, and 70.8% after 10, 20, and 30 min of exposure, respectively. The scolicidal effect of this extract at a concentration of 30 mg/ml was 52.9%, 86.7%, and 100% after 10, 20, and 30 min of exposure, respectively. *S. persica* extract at a concentration of 50 mg/ml, meanwhile, killed 81.4%, 100%, and 100% of protoscolices after 10, 20, and 30 min, respectively. Also, the cytotoxic potential of *S. persica* was assessed on human liver cells (HepG2) using trypan blue exclusion test. No cytotoxic effect was observed on HepG2 cell line. The present study confirmed for the first time that the ethanolic extract of *S. persica* has high scolicidal power in vitro. However, in vivo effect of this material remains to be studied for treatment of echinococcosis in humans and herbivorous animals.

**Key words:** *Echinococcus granulosus*, hydatidosis, sheep, liver, Arak, scolicidal agent, cytotoxicity

## INTRODUCTION

Echinococcosis or hydatidosis is a serious zoonotic infection with a worldwide distribution, caused by larval stages of cestodes belonging to the genus *Echinococcus* [1]. Infection with *Echinococcus granulosus*, the causative agent of cystic echinococcosis, is found on all continents, with the highest prevalence in parts of Eurasia (especially Mediterranean countries, the Russian Federation and adjacent independent states, and China), north and east Africa, Australia, and South America [2]. Hydatid cysts of *E. granulosus* develop in internal organs (mainly the liver and lungs) of humans and other intermediate hosts as unilocular fluid-filled bladders [3]. Although the initial phase of the infection is always asymptomatic for many years or even permanently [2]. Without efficient treatment, the continuous

development of the cyst will ultimately result in organ malfunction and even death in many cases [4].

Currently, there are 3 treatment choices for hydatidosis: surgery, which remains the foremost effective treatment, percutaneous aspiration, and medicinal treatment [5]. Surgery is still the preferred method of treatment but one of the major surgical complications of hydatidosis is the recurrence after operation for primary hydatid disease [6]. The spillage of protoscolices-rich fluid during surgery is a major cause of recurrence and multiple secondary hydatidosis [7]. The use of effective scolicidal agents, while avoiding spilling the cyst contents is an integral part of the surgical technique for this intervention, helping to reduce the risk of spilling viable protoscolices [8]. Many scolicidal agents have been used to inactivate the content of cysts, but most are not safe due to their side effects such as sclerosane cholangitis (biliary tract fibrosis), liver necrosis, and methemoglobinemia [9]. The development of safe and effective new scolicidal agents, especially from natural sources, is therefore of great interest [10].

Arak (*Salvadora persica*) is one of the most commonly used medicinal plants for treatment of rheumatism, leprosy, gonorr-

•Received 6 March 2015, revised 30 September 2015, accepted 1 November 2015.

\*Corresponding author (guraishi@yahoo.com)

© 2016, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

rhea, ulcers, scurvy, tumors, and dental diseases [11]. Arak has been used for centuries as a natural toothbrush (as Mesiwak or Siwak), especially in Muslim countries as a sunnah of the prophet, while its fibrous branches have been promoted by the World Health Organization for use in oral hygiene [12].

No laboratory studies have documented the effect of *S. persica* on protoscolices of hydatid cysts. Therefore, the present study was conducted to investigate the in vitro scolicidal effect of root extracts of this plant against protoscolices of *E. granulosus* at different exposure times. Also, the cytotoxic effect of *S. persica* extracts was evaluated using trypan blue exclusion test on human liver cells (HepG2).

## MATERIALS AND METHODS

### Collection of protoscolices and viability test

Hydatid cysts of *E. granulosus* naturally infected in the liver of sheep were obtained from a slaughterhouse in Riyadh city, Saudi Arabia. The intact cysts were immediately placed in an ice-box and transported to the Parasitology Laboratory, Zoology Department, College of Science, King Saud University, Riyadh, Saudi Arabia. Hydatid fluid along with protoscolices was collected according to Smyth and Barret [13]. The collected fluid with protoscolices was transferred into glass cylinders and left to set for 30 min to allow the protoscolices to settle to the bottom of the cylinders. The supernatant was then discarded, and the yielded protoscolices were washed 3 times in nor-

mal saline.


The viability of protoscolices (Fig. 1A) was confirmed from their motility characteristics under light microscopy. The viability was also assessed using 0.1% aqueous solution of eosin stain (1 g of eosin powder in 1,000 ml distilled water). Five min after exposure to the stain, unstained protoscolices were considered as viable (Fig. 1B), while stained protoscolices were considered as non-viable (Fig. 1C) [14]. When 95% or more viable protoscolices were present in the sediments, the sample was considered to be suitable for further experiments, and these were transferred into a dark container containing normal saline and stored at 4°C until use.

### Preparation of *S. persica* extract

*S. persica* L. (Salvadoraceae) roots (Arak) were collected from Jizan, Saudi Arabia. The identification was confirmed by a local expert at the herbarium of the Botany and Microbiology Department, College of Science, King Saud University. Roots were cut into small pieces, and ground into a fine powder using an electrical blender. About 100 g of root powder was extracted using 400 ml of 70% ethanol. The mixture was agitated for 30 min [15], and then maintained at rest for 24 hr. The resulting extract was filtered using a sterile filter paper, and then the solvent was completely removed using a rotary evaporator.

### In vitro scolicidal tests

We tested 3 concentrations (10, 30, and 50 mg/ml) of *S. per-*



**Fig. 1.** Protoscolices of *E. granulosus* collected from naturally infected livers of sheep. Scale bar = 100  $\mu$ m. (A) Live protoscolices without staining. (B) Live protoscolices after staining with 0.1% eosin. (C) Dead protoscolices after treatment with extracts and staining with 0.1% eosin.

*sica* extract for 10, 15, and 30 min. To prepare these concentrations, respectively, 0.1, 0.3, and 0.5 g of dried extract was dissolved in 10 ml of distilled water. Then, 2.5 ml of each concentration was placed in a test tube to which about 10,000 washed protoscolices was added. The contents of the tube were mixed gently, and the tube was then incubated at 37°C for 10, 15, or 30 min. At the end of each incubation period, the supernatant was discarded carefully to avoid disturbing the settled protoscolices. One ml of 0.1% eosin stain was then added to the remaining settled protoscolices and mixed gently. After 5 min, the upper portion of the solution was again discarded. The remaining pellets of protoscolices were smeared on a glass slide, covered with a cover glass, and examined microscopically for viability. The percentages of dead protoscolices were determined by counting at least 450 protoscolices. Non-treated protoscolices in normal saline were considered as a control group. Each experiment was performed in triplicate.

## Cytotoxicity assessment

### Cell culture

Human live cell line (HepG2) that used in the experiment was obtained from American Type Culture Collection (ATCC) (Manassas, Virginia, USA). HepG2 cells were cultured in DMEM, supplemented with 10% FBS, 0.2% sodium bicarbonate, and antibiotic/antimycotic solution (1 ml/100 ml of medium). Cells were grown in 5% CO<sub>2</sub> at 37°C in high humid atmosphere.

### Trypan blue dye exclusion test

The test was based on the principles that live cells with an intact cellular membrane that exclude the trypan blue dye while dead cells uptake the dye and stained blue. The test was performed following the protocol of Siddiqui et al. [16]. In brief, HepG2 cells were exposed to various concentrations (10, 30, 50 mg/ml) of ethanol extracts of *S. persica* for different time intervals (10, 20, and 30 min). After the respective exposure, cell suspensions were aspirated, centrifuged at 2,000 rpm for 3 min and washed with sterile PBS twice immediately after the incubation. The cells were stained with trypan blue (0.4% solution at a ratio of 1:5, dye: cell suspension) and counted using a hemocytometer. The counting for live (unstained) and dead (blue stained) cells was made at 100× magnification under a phase-contrast inverted microscope. The untreated cells were also run simultaneously under the identical conditions and served as the control. Each experiment was performed in triplicate.

### Statistical analysis


Differences between the test and control groups were analyzed with chi-square test using a statistical package program (Sigma Plot version 11.0). *P*-values less than 0.01 were considered to be significant.

## RESULTS

The results of the present experiment revealed highly signifi-

**Table 1.** Scolicidal effect of Arak (*Salvadora persica*) extracts

Concentration	Experiment no.	10 min			20 min			30 min		
		Total proto-scolices	Dead proto-scolices	Mortality rate (%)	Total proto-scolices	Dead proto-scolices	Mortality rate (%)	Total proto-scolices	Dead proto-scolices	Mortality rate (%)
10 mg/ml	1	500	175	35	480	240	50	470	320	68.1
	2	480	182	37.9	500	262	52.4	490	343	70
	3	500	180	36	480	233	48.5	500	370	74
Total		1,480	537	36.3	1,460	735	50.3	1,460	1,033	70.8
30 mg/ml	1	500	270	54	490	433	88.4	500	500	100
	2	480	243	50.6	500	425	85	500	500	100
	3	480	260	52	460	399	86.7	480	480	100
Total		1,460	773	52.9	1,450	1,257	86.7	1,480	1,480	100
50 mg/ml	1	500	403	80.6	480	480	100	500	500	100
	2	500	410	82	490	490	100	500	500	100
	3	470	383	81.5	500	500	100	480	480	100
Total		1,470	1,196	81.4	1,470	1,470	100	1,480	1,480	100
Control	1	500	26	5.2	480	41	8.5	500	60	12
	2	460	25	5.4	500	44	8.8	460	52	10.4
	3	500	25	5	500	39	7.8	480	53	11
Total		1,460	76	5.2	1,480	124	8.4	1,440	165	11.5



**Fig. 2.** Assessment of the percent (%) of cell viability in HepG2 cells following exposure to various concentrations of ethanolic extract of *S. persica* using trypan blue exclusion assay. All values are mean.

cant ( $P < 0.001$ ) scolicidal effects against protoscolices of *E. granulosus* for all of the various concentrations of *S. persica* ethanolic extract, compared to the control results, both within the same period of time and different periods as shown in Table 1. While the maximum death rate in the control group was 11.5%, scolicidal activity of *S. persica* extract at a concentration of 10 mg/ml was 36.3%, 50.3%, and 70.8% after application for 10, 20, and 30 min, respectively. The scolicidal effect of the extract at a concentration of 30 mg/ml was 52.9%, 86.7%, and 100% after application for 10, 20, and 30 min, respectively. *S. persica* extract at a concentration of 50 mg/ml, meanwhile, killed 81.4%, 100%, and 100% of protoscolices after 10, 20, and 30 min, respectively.

The results of the cytotoxicity assay using trypan blue exclusion test are shown in Fig. 2. The cell viability was 100%, 99%, and 98% after exposure to 10 mg/ml for 10, 20, and 30 min, respectively, while it was 98%, 96%, and 95% after exposure to 30 mg/ml for the same time intervals. Application of 50 mg/ml extract for 10, 20, and 30 min showed cell viabilities of 96%, 94%, and 90%, respectively.

## DISCUSSION

Hydatid disease (hydatidosis) is a parasitic disease that remains a clinical problem worldwide, especially in areas where animal husbandry and subsistence farming form an integral part of community life [17]. Seventy-five percent of all hydatid cysts are found in the liver [18], and these are usually managed either by surgery or medicinal treatment [19]. Surgery is associated with considerable mortality, morbidity, and recur-

rence rate. It is also expensive, and, clearly, needs expertise [20]. Chemotherapy can be used as treatment for patients considered not fit for surgery, or as an adjuvant to surgical treatment preoperatively or postoperatively, or both [19]. Various degrees of success have been claimed for many drugs that are now being used in hydatidosis treatment [21]. However, the metabolites of these drugs, including albendazole, albendazole sulfoxide, benzimidazole, and mebendazole, are potentially toxic in some subjects and are associated with severe hepatobiliary complications [22]. Natural scolicidal agents offer a safe alternative with no adverse associated effects [9].

Researchers around the world have begun to explore the real effect of many plants traditionally used as medicines which have not hitherto been validated by rigorous scientific experimentation. In this context, recently, many plants have been screened for their in vitro scolicidal activities, including *Ocimum bacilicum* and *Allium cepa* [6], *Sambucus ebulus* [9], *Trachyspermum ammi* [10], *Allium sativum* [14], *Zingiber officinale* [23], and *Zataria multiflora* [24].

With respect to *S. persica*, Reuben et al. [25] confirmed very high anthelmintic effects (86.7% and 98.9%) for aqueous extracts of *S. persica* at concentrations of 25 mg/ml and 50 mg/ml, respectively, against strongylid nematodes. These results urged us to test the in vitro scolicidal effect of ethanolic extract of *S. persica*. It was shown in this study that the ethanolic extract of *S. persica* exhibited the strongest scolicidal effect (100% and 100%) after 20 min at a concentration of 50 mg/ml and after 30 min at a concentration of 30 mg/ml. The ethanolic extract of *S. persica* is reported to contain a number of important phytoconstituents, such as indole alkaloids (e.g., salvadoricine), flavonoids (e.g., quercetin), the sulphur-containing compound tropaeidin, triterpenes, phytosterols, and isothiocyanates (e.g., benzyl isothiocyanate) [26]. Sofrata et al. [26] proved that *S. persica* root sticks consist of more than 98% benzyl isothiocyanate, while another report has confirmed that benzyl isothiocyanate is the predominant or sole anthelmintic agent in papaya seed extracts against *Caenorhabditis elegans* [27]. Our results showed a high scolicidal effect of *S. persica* root extracts against protoscolices of *E. granulosus*, which might be due to the presence of anthelmintic isothiocyanates. The present study proved that the ethanolic extract of *S. persica* have no cytotoxic effect on HepG2 cells. Similarly, previous studies have found the ethanol extract of this plant to be completely devoid of cytotoxic effects on HGFs and on L929 consistently in 3 cytotoxic assays [28]. Studies have also shown that the extract is safe

in respect to liver and kidney functions and hematological parameters of rats [29]. Also, Ahmed et al. [30] confirmed that the ethanolic extract of *S. persica* did not show any untoward effect up to a dose of 5,000 mg/kg body weight and did not cause death in any tested albino mice.

In conclusion, the root ethanolic extract of *S. persica* is a safe and potent protoscolicide with the potential to be used in hydatid cyst treatment and pre-surgery to prevent secondary cyst recurrence. However, in vivo efficacy of this extract remains to be explored, and further studies are necessary to identify and isolate the active compounds. This is the first report on the scolical activity of *S. persica*.

## ACKNOWLEDGMENTS

We extend our appreciation to the Dean of Scientific Research, King Saud University, for funding the work through the research group project no. RG-004.

## CONFLICT OF INTEREST

The authors declare that the present study followed the ethical standards and the ethical rules applicable for this journal and no conflict of interest.

## REFERENCES

- McManus DP, Zhang W, Li J, Bartley PB. Echinococcosis. *Lancet* 2003; 362: 1295-1304.
- Gavara CG, López-Andújar R, Ibáñez TB, Ángel JMR, Herraiz AM, Castellanos FO, Ibars EP, Rodríguez FSJ. Review of the treatment of liver hydatid cysts. *World J Gastroenterol* 2015; 7: 124-131.
- Thompson RC, McManus DP. Towards a taxonomic revision of the genus *Echinococcus*. *Trends Parasitol* 2002; 18: 452-457.
- Liu C, Zhang H, Lei W, Zhang C, Jiang B, Zheng Q, Yin J, Han X. An alternative mebendazole formulation for cystic echinococcosis: the treatment efficacy, pharmacokinetics and safety in mice. *Parasit Vectors* 2014; 7: 589-598.
- Adas G, Arıkan S, Kemik O, Oner A, Sahip N, Karatepe O. Use of albendazole sulfoxide, albendazole sulfone and combined solutions as scolical agents on hydatid cysts (in vitro study). *World J Gastroenterol* 2009; 15: 112-116.
- Haghani A, Roozitalab A, Safi SN. Low scolical effect of *Ocimum bacilicum* and *Allium cepa* on protoscolices of hydatid cyst: an in vitro study. *Comp Clin Pathol* 2014; 23: 847-853.
- Moro P, Schantz PM. Echinococcosis: a review. *Int J Infect Dis* 2009; 13: 125-133.
- Puryan K, Karadayi K, Topcu O, Canbay E, Sumer Z, Turan M, Karayalcin K, Sen M. Chlorhexidine gluconate: an ideal scolical agent in the treatment of intraperitoneal hydatidosis? *World J Surg* 2005; 29: 227-230.
- Gholami SH, Rahimi-Esboei B, Ebrahimzadeh MA, Pourhajibagher M. In vitro effect of *Sambucus ebuluson* scolices of hydatid cysts. *Eur Rev Med Pharmacol Sci* 2013; 17: 1760-1765.
- Moazeni M, Saharkhiz MJ, Hosseini AA. In vitro lethal effect of ajowan (*Trachyspermum ammi* L.) essential oil on hydatid cyst protoscolices. *Vet Parasitol* 2012; 187: 203-208.
- Kumar S, Dhankhar S, Arya VP, Yadav S, Yadav JP. Antimicrobial activity of *Salvadora oleoides* Decne against some microorganisms. *J Med Plant Res* 2012; 6: 2754-2760.
- Almas AK, Almas K. Miswak (*Salvadora persica* chewing stick): the natural toothbrush revisited. *Odontostomatol Trop* 2014; 37: 27-39.
- Smyth JD, Barret NJ. Procedures for testing the viability of human hydatid cyst following surgical removal, especially after chemotherapy. *Trans R Soc Trop Med Hyg* 1980; 74: 649-652.
- Moazeni M, Nazer A. In vitro effectiveness of garlic (*Allium sativum*) extract on scolices of hydatid cyst. *World J Surg* 2010; 34: 2677-2681.
- Mau J, Chen C, Hsieh P. Antimicrobial effect of extracts from Chinese chive, cinnamon, and comi fructus. *J Agricult Food Chem* 2001; 49: 183-188.
- Siddiqui MA, Singh G, Kashyap MP, Khanna VK, Yadav S, Chandra D, Pant AB. Influence of cytotoxic doses of 4-hydroxynoneal on selected neurotransmitter receptors in PC-12 cells. *Toxicol In Vitro* 2008; 22: 1681-1688.
- Shaw JM, Bornman PC, Krige JEJ. Hydatid disease of the liver. *SAJS* 2006; 44: 70-77.
- Malik AA, Bari SU, Amin R, Jan M. Surgical management of complicated hydatid cysts of the liver. *World J Gastrointest Surg* 2010; 2: 78-84.
- Shams-Ul-Bari, Arif SH, Malik AA, Khaja AR, Dass TA, Naikoo ZA. Role of albendazole in the management of hydatid cyst liver. *Saudi J Gastroenterol* 2011; 17: 343-347.
- Nari G, Ponce O, Cirami M. Five years experience in surgical treatment of liver hydatidosis. *Int Surg* 2003; 88: 194-198.
- Barzinji AKR, Mothana RA, Nasher AK. Effect of leaf extracts of *Dendrosicyos socotrana* and *Jatropha unicostata* on the viability of *Echinococcus granulosus* protoscolices. *EurAsia J BioSci* 2009; 3: 122-129.
- Topcu O, Aydin C, Arici S, Duman M, Koyuncu A, Sen M. The effects of various scolical agents on the hepatopancreatic biliary system. *Viszeralmedizin* 2006; 22: 185-190.
- Moazeni M, Nazer A. In vitro lethal effect of *Zingiber officinale* R. on protoscolices of hydatid cyst from sheep liver. *Microbiol Res* 2011; 2: 91-94.
- Moazeni M, Roozitalab A. High scolical effect of *Zataria multiflora* on protoscolices of hydatid cyst: an in vitro study. *Comp Clin Pathol* 2012; 21: 99-104.
- Reuben DK, Aji SB, Andrew W, Abdulrahman FI. Preliminary

- phytochemical screening and in vitro anthelmintic effects of aqueous extracts of *Salvadora persica* and *Terminalia avicennoides* against Strongyline nematodes of small ruminants in Nigeria. *J Animal Vet Adv* 2011; 10: 437-442.
26. Sofrata AH, Claesson RL, Lingstram PK, Gustafsson AK. Strong antibacterial effect of miswak against oral microorganisms associated with periodontitis and caries. *J Periodontol* 2008; 79: 1474-1479.
27. Kermanshai R, McCarry BE, Rosenfeld J, Summers PS, Weretilnyk EA, Sorger GJ. Benzyl isothiocyanate is the chief or sole anthelmintic in papaya seed extracts. *Phytochemistry* 2001; 57: 427-435.
28. Balto HA, Al-Manei KK, Bin-Mohareb TM, Shakoor ZA, AlHadlaq SM. Cytotoxic effect of *Salvadora persica* extracts on human gingival fibroblast cells. *Saudi Med J* 2014; 35: 810-815.
29. Ibrahim AY, El-Gengaihi SE. Safety profile of meswak root extract on liver, kidney, sexual hormones and hematological parameters of rats. *Not Sci Biol* 2012; 4: 18-23.
30. Ahmad M, Imran H, Yaqeen Z, Rehman Z, Rahman A, Fatima N, Sohail T. Pharmacological profile of *Salvadora persica*. *Pak J Pharm Sci* 2011; 24: 323-330.