

# Improvement of Natamycin Production by Cholesterol Oxidase Overexpression in *Streptomyces gilvosporeus*<sup>S</sup>

Miao Wang<sup>1,2†</sup>, Shaohua Wang<sup>1,2†</sup>, Gongli Zong<sup>1,2</sup>, Zhongwen Hou<sup>1,2</sup>, Fei Liu<sup>1,2</sup>, D. Joshua Liao<sup>3</sup>, and Xiqiang Zhu<sup>1,2\*</sup>

<sup>1</sup>Institute of Biopharmaceuticals of Shandong Province, Shandong Key Laboratory of Biological Drugs, Jinan 250101, P.R. China

<sup>2</sup>Shandong Academy of Pharmaceutical Science, Polysaccharide Drugs Engineering Laboratory of Shandong Province, Jinan 250101, P.R. China

<sup>3</sup>Hormel Institute, University of Minnesota, Austin, MN55912, USA

Received: May 11, 2015  
Revised: September 11, 2015  
Accepted: October 27, 2015

First published online  
October 27, 2015

\*Corresponding author  
Phone: +86-53181213080;  
Fax: +86-53181213083;  
E-mail: article13256745998@126.com

†These authors contributed  
equally to this work.

**S**upplementary data for this  
paper are available on-line only at  
<http://jmb.or.kr>.

pISSN 1017-7825, eISSN 1738-8872

Copyright© 2016 by  
The Korean Society for Microbiology  
and Biotechnology

Natamycin is a widely used antifungal antibiotic. For natamycin biosynthesis, the gene *pimE* encodes cholesterol oxidase, which acts as a signalling protein. To confirm the positive effect of the gene *pimE* on natamycin biosynthesis, an additional copy of the gene *pimE* was inserted into the genome of *Streptomyces gilvosporeus* 712 under the control of the *ermE*\* promoter (*p<sub>ermE</sub>\**) using intergeneric conjugation. Overexpression of the target protein engendered 72% and 81% increases in the natamycin production and cell productivity, respectively, compared with the control strain. Further improvement in the antibiotic production was achieved in a 1 L fermenter to 7.0 g/l, which was a 153% improvement after 120 h cultivation. Exconjugants highly expressing *pimE* and *pimM* were constructed to investigate the effects of both genes on the increase of natamycin production. However, the co-effect of *pimE* and *pimM* did not enhance the antibiotic production obviously, compared with the exconjugants highly expressing *pimE* only. These results suggest not only a new application of cholesterol oxidase but also a useful strategy to genetically engineer natamycin production.

**Keywords:** Natamycin, cholesterol oxidase, *Streptomyces gilvosporeus*, intergeneric conjugations

## Introduction

Natamycin, also known as pimaricin, is a widely used polyene antifungal antibiotic [19]. Its antifungal activity is achieved by its ability to bind to steroidal components in the fungal cell membrane, leading to the leakage of cellular materials [23]. Natamycin is a very potent inhibitor of yeast and mold growth as well as of aflatoxin synthesis. Therefore, natamycin plays an important role in the treatments of many fungal diseases, such as bronchopulmonary aspergillosis [6] and mycotic keratitis.

Natamycin is considered to be effective on a broad spectrum of fungi. It is highly effective at low doses against yeast and molds, but displays low toxicity to mammalian cells. Globally, natamycin is also a widely permitted food additive that is being legally used in over 50 countries,

including the USA and Europe, for the safe preservation of a wide variety of food products, such as cheese, fermented meat, yogurt, beverages, wine, and baked foods, against yeasts and molds [4, 5, 17]. It is one of the very few antibiotics that have been recommended as food additives and belong to the Generally Regarded As Safe list of compounds by the US Food and Drugs Administration. Hence, this antibiotic has considerable commercial value.

Natamycin is produced in submerged culture under aerobic conditions by *Streptomyces natalensis* [14], *Streptomyces chattanoogensis* [7], and *Streptomyces gilvosporeus* [13]. Current industrial concerns include a significant improvement of its production to meet increasing commercial demand. Recently, the polyene macrolide antibiotic biosynthetic gene cluster was characterized [3]. Consequently, increasing attention has been given to genetic manipulation as a good



1.0 kb *pimM* fragment including the whole ORF and its native promoter was inserted into the EcoRI and XbaI double-digested pSET152, generating the plasmid pSET152-*pimM*. Then a 1.9 kb *p<sub>ermE</sub>\*-pimE* fragment was brought into the XbaI-digested pSET152, generating the plasmid pSET152-*pimM-pimE*. Then, pSET152-*pimM-pimE* was introduced by conjugation into the original strain, *S. gilvosporeus* 712 (yielding *S. gilvosporeus-pimE-pimM*), as described above.

### Production of Natamycin in Fermentation Culture

Spores of *Streptomyces* were harvested on MS (20 g/l mannitol, 20 g/l soya flour) agar medium. For fermentation, ~1 cm<sup>2</sup> agar pieces of *S. gilvosporeus* 712, *S. gilvosporeus*-pSET152, and *S. gilvosporeus* swjs-801 were activated in the seed medium (10 g/l glucose, 5 g/l peptone, 3 g/l yeast extract, and 3 g/l malt extract) for 20 h at 29°C. The obtained cells were inoculated into 30 ml of fermentation medium (20 g/l peptone, 4.5 g/l yeast extract, 2 g/l NaCl, 1 g/l MgSO<sub>4</sub>, and 60 g/l glucose) in a 250 ml Erlenmeyer flask at a rate of 5% (v/v), and cultured for 120 h at 29 °C and 220 ×g. Aliquots were transferred to 700 ml of fermentation medium in a 1 L fermenter (INFORS, Switzerland), and grown at 29°C for 120 h with the aeration rate of 1 vvm and at least 30% dissolved oxygen, relevant to the agitation rate adjustment. Antifoaming agent was added once needed at a rate of 0.5% (v/v), and the glucose level was monitored at 20–30 g/l throughout the fermentation process by glucose feeding.


Comparison of natamycin production between *S. gilvosporeus* swjs-801 and *S. gilvosporeus-pimE-pimM* was conducted in the shake flask cultures as described above.

### Detection of Protein Expression with His<sub>6</sub>-tag

A His<sub>6</sub>-tag sequence was added in the gene *pimE* for the detection of the protein. *S. gilvosporeus* 712 and *S. gilvosporeus* swjs-801 were cultured in shake flasks for 3 days. Subsequently, 10 ml of the culture broth was centrifuged for 20 min at 10,000 ×g. The supernatant was concentrated 100 times as an SDS-PAGE sample, which was then fractioned on a 10% polyacrylamide gel. Protein was detected using a western blot assay, with a mouse anti His<sub>6</sub>-tag antibody (CW0286; CWBIO, China) and goat anti-mouse IgG conjugated with horseradish peroxidase (CW0102; CWBIO).

### Analytical Methods

Preliminary screening of exconjugants was bioassayed using *Candida albicans* as the test fungi. Clarified supernatants (1 ml) from shake flask or bioreactor cultures (obtained by centrifugation at 10,000 ×g for 20 min) were used for the determination of the PimE activity. The PimE activity was determined using the hydrogen peroxide colorimetric method of Allain *et al.* [1]. Dry cell weight was assayed by washing the remaining cell pellets twice with distilled water, followed by drying to constant weight at 80°C. For natamycin determination, the culture broth (1 ml) was ultrasonically extracted for 20 min using 9 ml of absolute



**Fig. 1.** Effect of commercial cholesterol oxidase addition on natamycin production by *S. gilvosporeus* 712 (values are the mean ± standard deviation, *n* = 3).

methanol and then centrifuged at 10,000 ×g for 10 min. HPLC analysis of natamycin was carried out using the method of Wang *et al.* [24].

## Results and Discussion

### Effects of Supplementation with Escalating Concentrations of PimE on Natamycin Production by *S. gilvosporeus* 712

Three parallel cultures were conducted to investigate the effects of supplementation with varying (1, 4, 7, 10, and 50 mg/l) PimE levels on natamycin production by *S. gilvosporeus* 712. As shown in Fig. 1, main culture supplementation with PimE stimulated natamycin production compared with the control. Initially, antibiotic production rose with increasing levels of PimE supplementation, thereby achieving peak natamycin production of 1.58 g/l at 7 mg/l PimE, which indicated 56.4% more natamycin production compared with the control. Further supplementation with PimE resulted in a reduced antibiotic production, although the accumulation of natamycin as well as cell productivity was still higher than that in the control. Cell growth was not affected by PimE supplementation at 1–7 mg/l, but higher levels resulted in progressively lower cell growth. These results collectively confirmed that the expression of PimE was insufficient for natamycin biosynthesis in *S. gilvosporeus* 712, and prompted us to investigate whether the homologous PimE can produce better effects.

### Construction of *S. gilvosporeus-pimE* and *S. gilvosporeus-pimM-pimE*

Exconjugants were selected after at least three cycles of

single colony purification, and PCR verified the successful insertion in the plasmids pSET152-*pimE* and pSET152-*pimM-pimE*. Furthermore, the exconjugants have remained stable throughout several generations. The strain with the largest bacteriostatic ring during bioassay was then used in the experiments.

#### Detection of PimE-His<sub>6</sub>


To distinguish the endogenous PimE from the recombinant PimE in *S. gilvosporeus*, the inserted gene (*pimE*) was tagged with a His<sub>6</sub>-Tag. The *pimE* gene encompasses a 1,650 bp DNA and encodes 549 amino acids. As shown in Fig. 2, the PimE-specific band at the predicted molecular mass (60 kDa) was detectable in lane 2, but not in the control line, thereby confirming the expression of PimE in *S. gilvosporeus* swjs-801.

#### Effects of *pimE* Overexpression on Natamycin Production in Shake-Flask Cultures

To investigate the effects of *pimE* overexpression on natamycin production and kinetics of cell growth, *S. gilvosporeus* 712, *S. gilvosporeus*-pSET152, and *S. gilvosporeus* swjs-801 were cultured in 250 ml shake flasks of a 30 ml working volume for 120 h. *S. gilvosporeus*-pSET152, loaded with null plasmid, did not display any significant difference in cell growth and natamycin production compared with the control strain, which presented similarly as *Streptomyces* sp. US 24 and *S. gilvosporeus*-*vgb* [22, 24] (data not shown). Cell growth (Fig. 3B) and sugar consumption (Fig. 3C) curves were also similar to the counterparts in PimE-overexpressing and control strains, suggesting that the integration of the gene *pimE* did not affect growth or antibiotic biosynthesis. In the late stage of fermentation, natamycin production soared beyond expectation probably because of the tardy growth of the cells. However, the recombinant strain displayed a noteworthy increase in natamycin production (72% ( $p < 0.05$ ); Fig. 3A) and bioproductivity (81% ( $p < 0.05$ ); Fig. 3D).

#### Effects of *pimE* Overexpression on Natamycin Production in 1 L Fermenter


To further investigate the effects of *pimE* overexpression on the natamycin production and the kinetics of cell growth, *S. gilvosporeus* 712 and *S. gilvosporeus* swjs-801 were cultured in 1 L fermenters of a 700 ml working volume, with dissolved oxygen and glucose levels controlled. As shown in Fig. 4, in parallel with cell growth, natamycin accumulated in the culture. A significant increase in the




**Fig. 2.** Western blot analysis of PimE protein in the supernatant of *S. gilvosporeus* 712 (lane 1) and *S. gilvosporeus* swjs-801 (lane 2), with the Thermo Scientific PageRuler prestained protein ladder as the marker (lane M).

natamycin yield was observed (Fig. 4A), with the concentration reaching 7.0 g/l, which was 153% ( $p < 0.05$ ) higher than that in the control. Generally, cell productivity was higher (Fig. 4D) in the fermenters, probably because of the continuous oxygen supply and better pH control. Remarkably, PimE activity was higher in the recombinant strain than that in the control (Fig. 4C), specifically in the late fermentation process. As reported by Mendes *et al.* [15], PimE is a signalling protein in natamycin production that relies on its enzymatic activity. Compared with heterogenous PimE, the homologous one strongly triggered natamycin production with a slightly higher enzymatic activity. Thus, PimE plays a key role in natamycin biosynthesis.

The PimE-encoding gene appears to work as a marker for the polyene antibiotic gene cluster that has 26–28 members [16, 21]. We infer that this gene could play a similar role as in the biosynthesis of polyene antibiotics. Our results provide a novel application of PimE in the biosynthesis of polyenes. However, medium composition, pH value, and dissolved oxygen also need to be optimized. Researchers are currently considering the insertion of *pimE*, together with *vgb* (*Vitreoscilla* hemoglobin gene), into the genome of *S. gilvosporeus* 712 to further improve natamycin productivity. However, the exact mechanisms underlying the role of PimE as a signalling protein in polyene antibiotic biosynthesis remains to be understood. Further experiments will provide the answer to this question.



**Fig. 3.** Kinetics of cell growth and natamycin production by *S. gilvosporeus* 712 and *S. gilvosporeus* swjs-801 in shake-flask cultures. Natamycin production (A), dry cell weight (B), residual glucose (C), and cell productivity (D) were measured (values are the mean  $\pm$  standard deviation,  $n = 3$ ).



**Fig. 4.** Kinetics of cell growth and natamycin production by *S. gilvosporeus* 712 and *S. gilvosporeus* swjs-801 in 1 L fermenters. Natamycin production (A), dry cell weight (B), enzyme activities (C), and cell productivity (D) were determined (values are the mean  $\pm$  standard deviation,  $n = 3$ ).



### Comparison of Natamycin Production Between *S. gilvosporeus* swjs-801 and *S. gilvosporeus-pimE-pimM*

To investigate the effects of PimE and PimM on natamycin production, *S. gilvosporeus* swjs-801 and *S. gilvosporeus-pimE-pimM* were cultured in 250 ml shake flasks of a 30 ml working volume for 120 h. The growth characteristics were similar in both exconjugants (data not shown), and otherwise exconjugants highly expressing PimE and PimM did not enhance natamycin production obviously (only 6.2% higher, shown in Fig. S1). PimM controls multiple natamycin biosynthetic promoters directly; however, it does not control that of *pimH* or *pimE* [2]. PimE acts as a signalling protein, which is being investigated [15]. Thus, we infer that PimM and PimE may act on the same genes of the natamycin biosynthetic gene cluster via different regulator circuits, consequently making the regulatory effects less obvious.

In summary, in response to increasing demands for natamycin, an additional copy of the gene *pimE* was integrated into the chromosome of *S. gilvosporeus*, a well-known natamycin producer. As a result of the gene insertion, PimE was overexpressed in *S. gilvosporeus* as an active form and significantly enhanced the natamycin yield. This effect was even more significant on the fermenter level. We believe that this study provides a better natamycin-producing industrial strain, as well as an excellent strategy to raise the yields of other secondary metabolites in industrial strains.

### Acknowledgments

We would like to thank Dr. Fred Bogott at Austin Medical Center, Austin of Minnesota, for his excellent English edition of this manuscript. This work was financially supported by Shandong Province High-Tech Industry Major Projects (No. 2014CGZH1306) and Jinan High-Tech Industry Major Projects (No. 201403009).

### References

- Allain CC, Poon LS, Chan CS, Richmond W, Fu PC. 1974. Enzymatic determination of total serum cholesterol. *Clin. Chem.* **20**: 470-475.
- Anton N, Aberturas JS, Mendes MV, Guerra SM, Martin JF, Aparicio JF. 2007. PimM, a PAS domain positive regulator of pimaricin biosynthesis in *Streptomyces natalensis*. *Microbiology* **153**: 3174-3183.
- Aparicio JF, Colina AJ, Ceballos E, Martin JF. 1999. The biosynthetic gene cluster for the 26-membered ring polyene macrolide pimaricin. *J. Biol. Chem.* **274**: 10133-10139.
- Arroyo-López FN, Bautista-Gallego J, Romero-Gil V, Rodríguez-Gómez F, Garrido-Fernández A. 2012. Growth/no growth interfaces of table olive related yeasts for natamycin, citric acid and sodium chloride. *Int. J. Food Microbiol.* **155**: 257-262.
- Bierhalz ACK, Silva MAD, Kieckbusch TG. 2012. Natamycin release from alginate/pectin films for food packaging applications. *J. Food Eng.* **110**: 18-25.
- Currie DC, Lueck C, Milburn HJ, Harvey C, Longbottom JL, Darbyshire JH, et al. 1990. Controlled trial of natamycin in the treatment of allergic bronchopulmonary aspergillosis. *Thorax* **45**: 447-450.
- Du YL, Chen SF, Cheng LY, Shen XL, Tian Y, Li YQ. 2009. Identification of a novel *Streptomyces chattanoogensis* L10 and enhancing its natamycin production by overexpressing positive regulator ScnRII. *J. Microbiol.* **47**: 506-513.
- Enríquez LL, Mendes MV, Antón N, Tunca S, Guerra SM, Martín JF, Aparicio JF. 2006. An efficient gene transfer system for the pimaricin producer *Streptomyces natalensis*. *FEMS Microbiol. Lett.* **257**: 312-318.
- Flegg HM. 1973. An investigation of the determination of serum cholesterol by an enzymic method. *Clin. Biochem.* **10**: 79-87.
- Flett F, Mersinias V, Smith CP. 1997. High efficiency intergeneric conjugal transfer of plasmid DNA from *Escherichia coli* to methyl DNA-restricting streptomycetes. *FEMS Microbiol. Lett.* **155**: 223-229.
- Horton RM, Ho SN, Pullen JK, Hunt HD, Cai ZL, Pease LR. 1993. Gene splicing by overlap extension. *Methods Enzymol.* **217**: 270-279.
- Jang BY, Hwang YI, Choi SU. 2011. Effects of *pimM* and *pimR* on the increase of natamycin production in *Streptomyces natalensis*. *J. Kor. Soc. Appl. Biol. Chem.* **54**: 141-144.
- Liang J, Xu Z, Liu T, Cen P. 2008. Effects of cultivation conditions on the production of natamycin with *Streptomyces gilvosporeus* LK-196. *Enzyme Microb. Tech.* **42**: 145-150.
- Mendes MV, Recio E, Fouces R, Luiten R, Martin JF, Aparicio JF. 2001. Engineered biosynthesis of novel polyenes: a pimaricin derivative produced by targeted gene disruption in *Streptomyces natalensis*. *Chem. Biol.* **8**: 635-644.
- Mendes MV, Recio E, Anton N, Guerra SM, Santos AJ, Martin JF, Aparicio JF. 2007. Cholesterol oxidases act as signaling proteins for the biosynthesis of the polyene macrolide pimaricin. *Chem. Biol.* **14**: 279-290.
- Omura S, Ikeda H, Ishikawa J, Hanamoto A, Takahashi C, Shinose M, et al. 2001. Genome sequence of an industrial microorganism *Streptomyces avermitilis*: deducing the ability of producing secondary metabolites. *Proc. Natl. Acad. Sci. USA* **98**: 12215-12220.
- Pintado CMBS, Ferreira MASS, Sousa I. 2010. Control of pathogenic and spoilage microorganisms from cheese surface by whey protein films containing malic acid, nisin and natamycin. *Food Control* **21**: 240-246.

18. Purcell JP, Greenplate JT, Jennings MG, Ryerse JS, Pershing JC, Sims SR, *et al.* 1993. Cholesterol oxidase: a potent insecticidal protein active against boll weevil larvae. *Biochem. Biophys. Res. Commun.* **196**: 1406-1413.
19. Raab WP. 1974. Natamycin (Pimaricin): its properties and possibilities in medicine. *J. Basic Microbiol.* **14**: 172.
20. Sambrook J, Russell DW. 2001. *Molecular Cloning: A Laboratory Manual*, 3<sup>rd</sup> Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
21. Seco EM, Pérez-Zúñiga FJ, Rolón MS, Malpartida F. 2004. Starter unit choice determines the production of two tetraene macrolides, rimocidin and CE-108 in *Streptomyces diastaticus* var. 108. *Chem. Biol.* **11**: 357-366.
22. Sioud S, Aigle B, Karray-Rebai I, Smaoui S, Bejar S, Mellouli L. 2009. Integrative gene cloning and expression system for *Streptomyces* sp. US 24 and *Streptomyces* sp. TN 58 bioactive molecule producing strains. *Biomed. Res. Int.* **1**: 83-99.
23. te Welscher YM, ten Napel HH, Balagué MM, Souza CM, Riezman H, de Kruijff B, Breukink E. 2008. Natamycin blocks fungal growth by binding specifically to ergosterol without permeabilizing the membrane. *J. Biol. Chem.* **283**: 6393-6401.
24. Wang SH, Liu F, Hou ZW, Zong GL, Zhu XQ, Ling PX. 2014. Enhancement of natamycin production on *Streptomyces gilvosporeus* by chromosomal integration of the *Vitreoscilla* hemoglobin gene (*vgb*). *World J. Microbiol. Biotechnol.* **30**: 1369-1376.